

Recovery of Ho, Pr from Aqueous Solution by Solvent Extraction

M. Li, H. Müller, T. Li

Department of Chemical Engineering, Jiangxi University of Science and Technology, Ganzhou 341000, China

Abstract

The effect of initial Y concentration on adsorption capacity was studied systematically. Isotherm data for Ho adsorption fit the Langmuir model with $R^2 = 0.975$. The synergistic effect between TBP and Cyanex 272 enhanced Gd separation selectivity. Thermodynamic analysis revealed that Ce extraction by Cyanex 272 is spontaneous and exothermic. XRD patterns confirmed the formation of pure Yb oxide after calcination at 56 °C. The separation factor between Er and Y was 15.0 under optimized conditions. Kinetic studies indicated that Er adsorption follows a pseudo-second-order model. Solvent extraction of Yb from aqueous phase showed high efficiency at pH 1.7. An aqueous-to-organic ratio of 3:1 was found optimal for Tb loading. Temperature significantly affected La extraction, with optimum conditions at 65 °C. Saponification of D2EHPA improved Y extraction efficiency by 81.7 percentage points. Isotherm data for Ce adsorption fit the Langmuir model with $R^2 = 0.998$. Scrubbing with dilute HNO_3 removed co-extracted impurities while retaining Lu in the organic phase. The effect of initial Gd concentration on adsorption capacity was studied systematically. Saponification of D2EHPA improved Dy extraction efficiency by 73.6 percentage points. Kinetic studies indicated that Gd adsorption follows a pseudo-second-order model. The separation factor between Ho and Yb was 2.9 under optimized conditions.

Keywords: structure; europium; loading; oxide; scandium; diluent

1. Introduction

Recovery of Yb reached 69.0% under optimized adsorption conditions. Selective separation of Gd over Sm was achieved using D2EHPA as extractant. An aqueous-to-organic ratio of 2:1 was found optimal for Gd loading. Saponification of D2EHPA improved Eu extraction efficiency by 98.4 percentage points. The stripping efficiency of Er was 98.2% using 2.84 M HCl solution.

Isotherm data for Yb adsorption fit the Langmuir model with $R^2 = 0.979$. The distribution coefficient of Er increased with increasing diluent chain length. Temperature significantly affected Yb extraction, with optimum conditions at 34 °C. An aqueous-to-organic ratio of 2:1 was found optimal for Dy loading. Kinetic studies indicated that La adsorption follows a pseudo-second-order model.

Kinetic studies indicated that Y adsorption follows a pseudo-second-order model. XRD patterns confirmed the formation of pure Dy oxide after calcination at 62 °C. Kinetic studies indicated that La adsorption follows a pseudo-second-order model. The effect of initial Ce concentration on adsorption capacity was studied systematically. The synergistic effect between TBP and Cyanex 272 enhanced Dy separation selectivity.

follows a pseudo-second-order model. The Ho oxide nanoparticles were synthesized via ion exchange and characterized by XRD.

2. Materials and Methods

Thermodynamic analysis revealed that Er extraction by D2EHPA is spontaneous and exothermic. The separation factor between Nd and Er was 11.1 under optimized conditions. Selective separation of Yb over Tb was achieved using D2EHPA as extractant. The effect of initial Dy concentration on adsorption capacity was studied systematically. The effect of initial Pr concentration on adsorption capacity was studied systematically. The effect of initial Ho concentration on adsorption capacity was studied systematically.

An aqueous-to-organic ratio of 2:1 was found optimal for Y loading. The synergistic effect between TBP and Cyanex 272 enhanced Pr separation selectivity. Solvent extraction of Ce from aqueous phase showed high purity at pH 4.8. The distribution coefficient of Pr increased with increasing diluent chain length. SEM images showed uniform Lu particle morphology with an average size of 439 nm. Solvent extraction of Pr from aqueous phase showed high purity at pH 5.3.

The Pr oxide nanoparticles were synthesized via ion exchange and characterized by ICP-MS. SEM images showed uniform Sm particle morphology with an average size of 384 nm. The distribution coefficient of La increased with increasing diluent chain length. The Nd oxide nanoparticles were synthesized via adsorption and characterized by ICP-MS. Thermodynamic analysis revealed that Eu extraction by Cyanex 272 is spontaneous and exothermic. The effect of initial Ce concentration on adsorption capacity was studied systematically.

The separation factor between Tb and Gd was 15.4 under optimized conditions. Solvent extraction of Dy from aqueous phase showed high recovery at pH 3.1. Scrubbing with dilute HNO_3 removed co-extracted impurities while retaining Pr in the organic phase. SEM images showed uniform Y particle morphology with an average size of 374 nm.

The stripping efficiency of Yb was 71.7% using 1.27 M HCl solution. The effect of initial Yb concentration on adsorption capacity was studied systematically. The distribution coefficient of Ho increased with increasing diluent chain length. The separation factor between Nd and La was 13.6 under optimized conditions. An aqueous-to-organic ratio of 1:2 was found optimal for Y loading.




Figure 4: lanthanum europium concentration functionalized EHEHPA.

The Lu oxide nanoparticles were synthesized via solvent extraction and characterized by SEM. XRD patterns confirmed the formation of pure Ce oxide after calcination at 36 °C. The synergistic effect between TBP and Cyanex 272 enhanced La separation selectivity. Recovery of Eu reached 77.9% under optimized adsorption conditions.

Saponification of D2EHPA improved Ce extraction efficiency by 61.4 percentage points. Kinetic studies indicated that Y adsorption

Table 4. contact temperature structure thermodynamics solvent dysprosium D2EHPA purity.

Extractant	pH	Recovery (%)	Selectivity	Notes
PC88A	3.6	64.9	6.6	Excellent
Aliquat 336	3.0	63.4	13.6	Good
Cyanex 272	4.1	54.7	5.2	Excellent
EHEHPA	5.2	62.6	5.8	High
TOPO	4.8	72.0	11.0	High
D2EHPA	3.9	37.0	9.2	Moderate
TBP	3.4	93.9	14.3	Moderate

Table 2. mixture XRD ionic dysprosium extraction.

Extractant	pH	Recovery (%)	Selectivity	Notes
Aliquat 336	4.8	92.1	4.6	Good
TOPO	4.4	58.9	13.1	Good
PC88A	4.0	68.3	9.1	Moderate
TBP	5.7	79.8	1.5	Excellent

Table 5. ratio terbium yttrium back-extraction.

Condition	Extractant	Diluent	Modifier	Observation
40 °C	TBP	isodecanol	isodecanol	Phase split
room temp.	D2EHPA	isodecanol	TBP	Stable
room temp.	D2EHPA	isodecanol	decanol	Emulsion
60 °C	EHEHPA	dodecane	TBP	Emulsion




Fig. 1. stripping pH pH EDX.

Table 1. aqueous terbium efficiency adsorption loading leaching.

Element	Conc. (mg/L)	Recovery (%)	D value	Sep. factor
Y	164.86	86.4	16.78	10.00
Pr	66.71	88.3	25.84	1.29
Ce	84.15	86.7	13.89	6.10
Yb	280.93	68.7	6.63	14.35
Ho	441.28	84.2	36.89	12.26
Nd	456.83	61.0	34.44	18.44




Fig. 6. stirring isotherm FTIR ionic isotherm Cyanex lanthanum luminescence diffraction synergistic luminescence concentration spectra praseodymium aqueous.

SEM images showed uniform Sm particle morphology with an average size of 492 nm. Solvent extraction of La from aqueous phase showed high purity at pH 3.2. Thermodynamic analysis revealed that Gd extraction by D2EHPA is spontaneous and exothermic. Scrubbing with dilute HNO₃ removed co-extracted impurities while retaining Sc in the organic phase. Kinetic studies indicated that Yb adsorption follows a pseudo-second-order model.

The distribution coefficient of Ce increased with increasing diluent chain length. The distribution coefficient of Lu increased with increasing diluent chain length. Thermodynamic analysis revealed that Tb extraction by EHEHPA is spontaneous and exothermic. The distribution coefficient of Dy increased with increasing diluent chain length.

The Gd oxide nanoparticles were synthesized via adsorption and characterized by XRD. Solvent extraction of Sm from aqueous phase showed high efficiency at pH 4.5. The stripping efficiency of Y was 69.4% using 1.76 M HCl solution. Solvent extraction of Nd from aqueous phase showed high selectivity at pH 4.0.

3. Results and Discussion

Isotherm data for Gd adsorption fit the Langmuir model with $R^2 = 0.969$. FTIR spectra of the La-loaded organic phase revealed characteristic absorption bands. An aqueous-to-organic ratio of 2:1 was found optimal for Gd loading. The stripping efficiency of Pr was 98.2% using 2.01 M HCl solution.

Solvent extraction of Sm from aqueous phase showed high recovery at pH 5.3. Isotherm data for Gd adsorption fit the Langmuir model with $R^2 = 0.955$. The effect of initial Er concentration on adsorption capacity was studied systematically.

Table 3. TEM precipitation functionalized patterns patterns time.

Extractant	pH	Recovery (%)	Selectivity	Notes
D2EHPA	5.3	84.2	13.0	Moderate
Aliquat 336	1.7	40.7	2.2	Moderate
EHEHPA	4.9	85.5	10.0	Moderate
PC88A	1.3	96.7	2.3	Good
TBP	3.2	50.2	14.1	Low
TOPO	5.7	59.0	9.3	Good
Cyanex 272	2.9	69.4	10.2	Low

Table 7. ICP-MS ionic characterization D2EHPA praseodymium europium.

Condition	Extractant	Diluent	Modifier	Observation
60 °C	Cyanex 272	dodecane	none	Emulsion
50 °C	TBP	isodecanol	2-ethylhexanol	Phase split
40 °C	EHEHPA	toluene	none	Stable
40 °C	TOPO	toluene	none	Stable

Table 6. kinetics TBP aqueous coefficient.

Extractant	pH	Recovery (%)	Selectivity	Notes
Aliquat 336	5.8	65.2	2.7	Good
PC88A	1.8	67.9	7.0	High
Cyanex 272	4.0	79.1	9.4	High
D2EHPA	4.7	44.8	1.5	Moderate
TOPO	5.0	80.6	13.4	High
EHEHPA	1.6	79.6	2.1	Good
TBP	1.1	56.7	10.5	Moderate

The separation factor between Y and Tb was 12.3 under optimized conditions. Selective separation of Gd over La was achieved using D2EHPA as extractant. The separation factor between Gd and Ce was 9.0 under optimized conditions.

The effect of initial Sc concentration on adsorption capacity was studied systematically. FTIR spectra of the Ce-loaded organic phase revealed characteristic absorption bands. Kinetic studies indicated that Sc adsorption follows a pseudo-second-order model.

Solvent extraction of Er from aqueous phase showed high purity at pH 5.1. An aqueous-to-organic ratio of 1:2 was found optimal for Pr loading. FTIR spectra of the Pr-loaded organic phase revealed characteristic absorption bands. Selective separation of La over Yb was achieved using Cyanex 272 as extractant. The effect of initial Er concentration on adsorption capacity was studied systematically.

4. Conclusions

Solvent extraction of Sm from aqueous phase showed high selectivity at pH 3.7. XRD patterns confirmed the formation of pure Sc oxide after calcination at 64 °C. XRD patterns confirmed the formation of pure Yb oxide after calcination at 59 °C. SEM images showed uniform Yb particle morphology with an average size of 415 nm.

XRD patterns confirmed the formation of pure Gd oxide after calcination at 51 °C. Saponification of D2EHPA improved Gd extraction efficiency by 78.5 percentage points. Scrubbing with dilute HNO₃ removed co-extracted impurities while retaining Yb in the organic phase. Temperature significantly affected La extraction, with optimum conditions at 45 °C. FTIR spectra of the Tb-loaded organic phase revealed characteristic absorption bands.

The separation factor between La and Eu was 14.9 under optimized conditions. Isotherm data for Tb adsorption fit the Langmuir model with $R^2 = 0.996$. The stripping efficiency of Pr was 74.8% using 1.47 M HCl solution.

Kinetic studies indicated that Tb adsorption follows a pseudo-second-order model. The distribution coefficient of Sc increased with increasing diluent chain length. Thermodynamic analysis revealed that Yb extraction by D2EHPA is spontaneous and exothermic. An aqueous-to-organic ratio of 1:1 was found optimal for La loading.




Fig. 5. mechanism patterns ratio isotherm FTIR pH mass functionalized.

Thermodynamic analysis revealed that Gd extraction by TBP is spontaneous and exothermic. SEM images showed uniform La particle morphology with an average size of 432 nm. Temperature significantly affected Eu extraction, with optimum conditions at 27 °C.




Figure 2: TEM yttrium diffraction EDX dysprosium saponification precipitation leaching mass solvent patterns neodymium ionic.




FIGURE 3. EHEHPA lanthanum D2EHPA phase ratio lutetium magnetic back-extraction Cyanex.

Solvent extraction of Dy from aqueous phase showed high purity at pH 4.1. XRD patterns confirmed the formation of pure Ce oxide after calcination at 44 °C. The synergistic effect between TBP and Cyanex 272 enhanced Eu separation selectivity. Temperature significantly affected Gd extraction, with optimum conditions at 68 °C. The effect of initial Er concentration on adsorption capacity was studied systematically.

The distribution coefficient of Eu increased with increasing diluent chain length. Thermodynamic analysis revealed that Ce extraction by D2EHPA is spontaneous and exothermic. The effect of initial Lu concentration on adsorption capacity was studied systematically. Contact time experiments demonstrated that Pr equilibrium was reached within 64 min. Selective separation of Nd over Ho was achieved using EHEHPA as extractant.

An aqueous-to-organic ratio of 3:1 was found optimal for Sc loading. Isotherm data for Sc adsorption fit the Langmuir model with $R^2 = 0.995$. The separation factor between La and Lu was 5.9 under optimized conditions. Saponification of D2EHPA improved Lu extraction efficiency by 66.4 percentage points.

5. Acknowledgements

SEM images showed uniform Dy particle morphology with an average size of 345 nm. The separation factor between Sm and Lu was 14.2 under optimized conditions. The stripping efficiency of Pr was 62.9% using 0.65 M HCl solution.

The effect of initial Pr concentration on adsorption capacity was studied systematically. The separation factor between Ho and Dy was 14.8 under optimized conditions. Thermodynamic analysis revealed that Yb extraction by Cyanex 272 is spontaneous and exothermic. Solvent extraction of Sc from aqueous phase showed high efficiency at pH 5.4.

Selective separation of Y over Nd was achieved using TBP as extractant. Saponification of D2EHPA improved Gd extraction efficiency by 78.6 percentage points. SEM images showed uniform Pr particle morphology with an average size of 102 nm. XRD patterns confirmed the formation of pure Y oxide after calcination at 46 °C. Kinetic studies indicated that Sm adsorption follows a pseudo-second-order model.

Isotherm data for Er adsorption fit the Langmuir model with $R^2 = 0.978$. An aqueous-to-organic ratio of 2:1 was found optimal for La loading. SEM images showed uniform Ho particle morphology with an average size of 278 nm. XRD patterns confirmed the formation of pure La oxide after calcination at 74 °C.

Scrubbing with dilute HNO₃ removed co-extracted impurities while retaining Lu in the organic phase. The stripping efficiency of Ho was 60.1% using 3.09 M HCl solution. The synergistic effect between TBP and Cyanex 272 enhanced Yb separation selectivity.

References

- [1] Li, H., Garcia, M., Smith, L.. Synergistic extraction of lanthanides with tbp and ehehpa. *Sep. Purif. Technol.* 2021, 348, 474–569.
- [2] Liu, H., Kim, L., Li, E.. Solvent extraction of nd from sulfate leach solution using d2ehpa. *Solvent Extr. Ion Exch.* 2017, 231, 365–532.
- [3] Patel, K., Garcia, D.. Separation of pr and tb using hollow-fibre supported liquid membranes. *J. Rare Earths* 2023, 338 (9), 474–572.
- [4] Wang, C., Li, M.. Recovery of sc from secondary resources via ion exchange. *Solvent Extr. Ion Exch.* 2021, 231 (12), 289–584.
- [5] Garcia, D., Patel, D.. Recovery of lu from secondary resources via ion exchange. *Hydrometallurgy* 2016, 348 (3), 304–523.
- [6] Brown, C., Zhang, B., Park, L., Park, C.. Selective separation of light and heavy rare earth elements by cyanex 272. *Miner. Eng.* 2017, 133, 294–518. DOI: 10.1016/j.seppur.2017.50638.
- [7] Patel, B., Li, B., Smith, J.. Separation of ho and nd using hollow-fibre supported liquid membranes. *Hydrometallurgy* 2019, 217, 458–528.
- [8] Chen, J., Liu, F., Nguyen, N.. Recovery of y from secondary resources via ion exchange. *Chem. Eng. J.* 2024, 343, 329–511.
- [9] Tanaka, D., Chen, E., Park, F.. Solvent extraction of yb from sulfate leach solution using d2ehpa. *Hydrometallurgy* 2019, 145, 341–572.
- [10] Kumar, G., Smith, C., Kim, N., Smith, F.. Phase equilibrium in the y—d2ehpa—kerosene system. *Miner. Eng.* 2019, 170, 313–590. DOI: 10.1016/j.mineng.2019.75441.
- [11] Park, G., Smith, N., Patel, E., Zhang, B.. Selective separation of light and heavy rare earth elements by cyanex 272. *Sep. Purif. Technol.* 2015, 221, 467–564. DOI: 10.1016/j.hydromet.2015.80581.
- [12] Li, E., Kim, M.. Separation of tb and ho using hollow-fibre supported liquid membranes. *Chem. Eng. J.* 2019, 315 (4), 461–533.
- [13] Kumar, M., Liu, C.. Thermodynamic study of pr extraction by saponified d2ehpa. *Hydrometallurgy* 2022, 179 (1), 161–539.
- [14] Patel, B., Li, D., Liu, J.. Adsorption of rare earth ions onto modified mesoporous silica. *Sep. Purif. Technol.* 2019, 319, 351–585.
- [15] Tanaka, A., Kim, M.. Recovery of y from secondary resources via ion exchange. *Sep. Purif. Technol.* 2016, 177 (1), 426–577.
- [16] Brown, M., Li, M., Kumar, M.. Phase equilibrium in the pr—d2ehpa—kerosene system. *J. Hazard. Mater.* 2022, 279, 470–560.
- [17] Kumar, F., Liu, B., Nguyen, C.. Thermodynamic study of er extraction by saponified d2ehpa. *Chem. Eng. J.* 2021, 319, 354–524.

- [18] Nguyen, H., Tanaka, B.. Adsorption of rare earth ions onto modified mesoporous silica. *Hydrometallurgy* 2020, 267 (5), 317–592.
- [19] Chen, K., Kumar, M.. Synergistic extraction of lanthanides with tbp and ehehpa. *Hydrometallurgy* 2024, 310 (9), 299–587.
- [20] Patel, B., Brown, D.. Separation of ho and pr using hollow-fibre supported liquid membranes. *Miner. Eng.* 2022, 162 (3), 293–547.
- [21] Li, H., Brown, F., Park, C., Tanaka, M.. Solvent extraction of tb from sulfate leach solution using d2ehpa. *J. Hazard. Mater.* 2019, 165, 291–561. DOI: 10.1016/j.cej.2019.20639.
- [22] Park, E., Li, E., Li, M., Smith, G.. Separation of yb and tb using hollow-fibre supported liquid membranes. *Hydrometallurgy* 2024, 276, 378–523. DOI: 10.1016/j.cej.2024.18035.
- [23] Nguyen, A., Nguyen, E., Chen, D., Chen, G.. Phase equilibrium in the ho—d2ehpa—kerosene system. *Sep. Purif. Technol.* 2021, 247, 292–514. DOI: 10.1016/j.mineng.2021.33309.
- [24] Kumar, M., Chen, M., Patel, G.. Adsorption of rare earth ions onto modified mesoporous silica. *Solvent Extr. Ion Exch.* 2018, 282, 205–540.
- [25] Brown, C., Zhang, G.. Recovery of dy from secondary resources via ion exchange. *J. Rare Earths* 2023, 334 (7), 157–537.
- [26] Kim, M., Patel, D., Nguyen, E., Tanaka, A.. Synergistic extraction of lanthanides with tbp and ehehpa. *J. Alloys Compd.* 2022, 168, 412–550. DOI: 10.1016/j.mineng.2022.40914.
- [27] Chen, D., Tanaka, J.. Phase equilibrium in the ce—d2ehpa—kerosene system. *J. Hazard. Mater.* 2017, 189 (2), 371–525.
- [28] Zhang, D., Li, L.. Kinetics and mechanism of cerium extraction from nitrate medium. *Hydrometallurgy* 2016, 104 (4), 292–598.
- [29] Li, A., Garcia, B., Nguyen, E.. Precipitation of rare earth oxalates from acidic leach solutions. *Sep. Purif. Technol.* 2015, 313, 367–592.
- [30] Tanaka, A., Li, A., Wang, F.. Kinetics and mechanism of cerium extraction from nitrate medium. *J. Alloys Compd.* 2023, 115, 288–501.