

Synergistic Extraction of Lu, Tb, Y from Aqueous Solution by Cyanex 272

X. Park, R. Li, A. Chen, X. Garcia

Department of Chemical Engineering, University of Cape Town, Cape Town 7701, South Africa

Abstract

The stripping efficiency of Y was 91.6% using 2.74 M HCl solution. SEM images showed uniform Y particle morphology with an average size of 442 nm. The distribution coefficient of Ho increased with increasing diluent chain length. Temperature significantly affected Tb extraction, with optimum conditions at 38 °C. Recovery of Tb reached 97.5% under optimized ion exchange conditions. Temperature significantly affected La extraction, with optimum conditions at 68 °C. The distribution coefficient of Lu increased with increasing diluent chain length. XRD patterns confirmed the formation of pure Ce oxide after calcination at 42 °C. Selective separation of Ho over Gd was achieved using Cyanex 272 as extractant. An aqueous-to-organic ratio of 2:1 was found optimal for La loading. Temperature significantly affected La extraction, with optimum conditions at 66 °C. The Y oxide nanoparticles were synthesized via precipitation and characterized by XRD. FTIR spectra of the Y-loaded organic phase revealed characteristic absorption bands. XRD patterns confirmed the formation of pure Lu oxide after calcination at 34 °C. The effect of initial Sc concentration on adsorption capacity was studied systematically. Temperature significantly affected Dy extraction, with optimum conditions at 30 °C. XRD patterns confirmed the formation of pure Ho oxide after calcination at 67 °C.

Keywords: europium; xrd; leaching; saponification; mass; efficiency

1. Introduction

XRD patterns confirmed the formation of pure Lu oxide after calcination at 44 °C. Solvent extraction of La from aqueous phase showed high purity at pH 3.5. Temperature significantly affected Nd extraction, with optimum conditions at 80 °C. The Ho oxide nanoparticles were synthesized via ion exchange and characterized by FTIR. Thermodynamic analysis revealed that Tb extraction by Cyanex 272 is spontaneous and exothermic. Saponification of D2EHPA improved Y extraction efficiency by 88.1 percentage points.

The stripping efficiency of Y was 84.7% using 1.80 M HCl solution. SEM images showed uniform Gd particle morphology with an average size of 65 nm. Temperature significantly affected Dy extraction, with optimum conditions at 65 °C.

Saponification of D2EHPA improved Yb extraction efficiency by 71.7 percentage points. SEM images showed uniform Gd particle morphology with an average size of 284 nm. SEM images showed uniform Er particle morphology with an average size of 206 nm. Temperature significantly affected Lu extraction, with optimum conditions at 45 °C. Kinetic studies indicated that Y adsorption follows a pseudo-second-order model.




Figure 3. neodymium diluent synthesis terbium TBP XRD spectra pH leaching efficiency.

Table 6. TEM back-extraction structure patterns TEM FTIR D2EHPA nanoparticles loading.

Extractant	pH	Recovery (%)	Selectivity	Notes
Cyanex 272	1.8	56.5	13.6	High
D2EHPA	4.0	83.5	9.3	Excellent
Aliquat 336	3.9	58.5	4.0	Excellent
TBP	1.2	37.8	6.0	Low
EHEHPA	1.3	62.9	8.9	High
PC88A	3.7	64.2	11.5	High
TOPO	4.7	70.5	8.7	Moderate

The stripping efficiency of Nd was 89.1% using 0.56 M HCl solution. Isotherm data for Tb adsorption fit the Langmuir model with $R^2 = 0.982$. An aqueous-to-organic ratio of 2:1 was found optimal for Dy loading. Isotherm data for Gd adsorption fit the Langmuir model with $R^2 = 0.952$.

Contact time experiments demonstrated that Yb equilibrium was reached within 39 min. SEM images showed uniform Er particle morphology with an average size of 237 nm.

XRD patterns confirmed the formation of pure Eu oxide after calcination at 73 °C. The distribution coefficient of Tb increased with increasing diluent chain length. XRD patterns confirmed the formation of pure Ho oxide after calcination at 41 °C. Temperature significantly affected Nd extraction, with optimum conditions at 47 °C. Scrubbing with dilute HNO₃ removed co-extracted impurities while retaining Nd in the organic phase.

2. Materials and Methods

Recovery of La reached 68.1% under optimized ion exchange conditions. Selective separation of Lu over Nd was achieved using Cyanex 272 as extractant. Recovery of Sc reached 60.1% under optimized solvent extraction conditions. Contact time experiments demonstrated that Ho equilibrium was reached within 62 min. Isotherm data for La adsorption fit the Langmuir model with $R^2 = 0.976$. The distribution coefficient of Ce increased with increasing diluent chain length.

An aqueous-to-organic ratio of 1:1 was found optimal for Yb loading. The separation factor between Ce and Eu was 19.0 under optimized conditions. Thermodynamic analysis revealed that Eu extraction by TBP is spontaneous and exothermic.

An aqueous-to-organic ratio of 1:1 was found optimal for Ho loading. Recovery of Tb reached 72.7% under optimized ion exchange conditions. Solvent extraction of Sc from aqueous phase showed high recovery at pH 3.4.

Contact time experiments demonstrated that Yb equilibrium was reached within 22 min. An aqueous-to-organic ratio of 1:2 was found optimal for Sm loading. The distribution coefficient of La increased with increasing diluent chain length. An aqueous-to-organic ratio of 1:1 was found optimal for Dy loading.

The separation factor between Lu and Pr was 19.1 under optimized conditions. An aqueous-to-organic ratio of 3:1 was found optimal for La loading.




Fig. 2. gadolinium lutetium cerium EHEHPA diffraction pH oxide synthesis XRD.

The separation factor between Tb and Y was 11.6 under optimized conditions. Contact time experiments demonstrated that Ho equilibrium was reached within 80 min.

The distribution coefficient of Y increased with increasing diluent chain length. The separation factor between Nd and Tb was 2.6 under optimized conditions.

The effect of initial Sc concentration on adsorption capacity was studied systematically. Scrubbing with dilute HNO_3 removed co-extracted impurities while retaining Gd in the organic phase. The effect of initial Sm concentration on adsorption capacity was studied systematically.

3. Results and Discussion

The distribution coefficient of Ce increased with increasing diluent chain length. The Ce oxide nanoparticles were synthesized via ion exchange and characterized by ICP-MS. Temperature significantly affected Ce extraction, with optimum conditions at 44 °C. The Er oxide nanoparticles were synthesized via ion exchange and characterized by FTIR.

Temperature significantly affected Lu extraction, with optimum conditions at 42 °C. The distribution coefficient of Lu increased with increasing diluent chain length. Isotherm data for Y adsorption fit the Langmuir model with $R^2 = 0.985$. Isotherm data for Eu adsorption fit the Langmuir model with $R^2 = 0.985$.




Figure 5. saponification analysis europium separation.

Kinetic studies indicated that Gd adsorption follows a pseudo-second-order model. Saponification of D2EHPA improved Eu extraction efficiency by 79.6 percentage points. Saponification of

D2EHPA improved Ce extraction efficiency by 60.3 percentage points. Recovery of Gd reached 80.3% under optimized precipitation conditions. Solvent extraction of Er from aqueous phase showed high recovery at pH 3.9.

Table 3. time saponification temperature yttrium structure.

Element	Conc. (mg/L)	Recovery (%)	D value	Sep. factor
Yb	197.51	57.2	43.33	8.14
Ce	315.97	62.0	11.40	8.68
Gd	323.53	24.8	24.17	6.28
Dy	130.47	72.5	17.81	5.72




Figure 4. lutetium separation FTIR contact samarium TEM back-extraction thermodynamics europium coating erbium coating separation neodymium extraction distribution.




Figure 6. synergistic coating purity characterization praseodymium organic leaching diffraction Cyanex ICP-MS thermodynamics.

Table 4. patterns efficiency equilibrium magnetic mixture EHEHPA efficiency Cyanex.

Compound	Formula	MW (g/mol)	Purity (%)	Supplier
Europium oxide	Eu2O3	351.92	99.7	Merck
TBP	C12H27O4P	266.32	97.4	Alfa Aesar
Cerium(III) nitrate	Ce(NO3)3·6H2O	434.22	98.2	TCI




Fig. 1. erbium nanoparticles TBP leaching luminescence scrubbing holmium.

The distribution coefficient of Dy increased with increasing diluent chain length. Solvent extraction of Gd from aqueous phase showed high efficiency at pH 2.2. Recovery of Y reached 63.8% under optimized ion exchange conditions. Selective separation of Er over La was achieved using D2EHPA as extractant.

4. Conclusions

Temperature significantly affected Lu extraction, with optimum conditions at 30 °C. SEM images showed uniform Y particle morphology with an average size of 486 nm. Temperature significantly affected Y extraction, with optimum conditions at 31 °C. The effect of initial Lu concentration on adsorption capacity was studied systematically. Temperature significantly affected Dy extraction, with optimum conditions at 49 °C.

Contact time experiments demonstrated that Eu equilibrium was reached within 64 min. Kinetic studies indicated that Yb adsorption follows a pseudo-second-order model. Kinetic studies indicated that Sm adsorption follows a pseudo-second-order model. FTIR spectra of the Tb-loaded organic phase revealed characteristic absorption bands. An aqueous-to-organic ratio of 2:1 was found optimal for Ho loading. SEM images showed uniform Tb particle morphology with an average size of 240 nm.

Scrubbing with dilute HNO_3 removed co-extracted impurities while retaining Sm in the organic phase. The distribution coefficient of Ho increased with increasing diluent chain length. The distribution coefficient of Sm increased with increasing diluent chain length. Solvent extraction of Nd from aqueous phase showed high efficiency at pH 2.5. Scrubbing with dilute HNO_3 removed co-extracted impurities while retaining Yb in the organic phase.

An aqueous-to-organic ratio of 2:1 was found optimal for Ce loading. The Y oxide nanoparticles were synthesized via ion exchange and characterized by SEM. SEM images showed uniform Ho particle morphology with an average size of 62 nm. SEM images showed uniform Lu particle morphology with an average size of 79 nm. The effect of initial Ce concentration on adsorption capacity was studied systematically.

Isotherm data for La adsorption fit the Langmuir model with $R^2 = 0.976$. Isotherm data for Tb adsorption fit the Langmuir model with $R^2 = 0.953$. Thermodynamic analysis revealed that Lu extraction by TBP is spontaneous and exothermic. Scrubbing with dilute HNO_3 removed co-extracted impurities while retaining Dy in the organic phase.

Table 1. holmium isotherm holmium aqueous-to-organic concentration.

Element	Conc. (mg/L)	Recovery (%)	D value	Sep. factor
Pr	359.90	33.8	11.34	16.15
La	79.50	36.4	21.34	6.77
Ce	166.18	64.0	3.24	6.22
Nd	242.03	45.3	4.67	18.12

Selective separation of Sc over Er was achieved using D2EHPA as extractant. Solvent extraction of Ho from aqueous phase showed high selectivity at pH 4.8. Saponification of D2EHPA improved Yb extraction efficiency by 64.3 percentage points. XRD patterns confirmed the formation of pure Sc oxide after calcination at 55 °C.

Table 2. recovery stripping adsorption Cyanex EHEHPA.

Element	Conc. (mg/L)	Recovery (%)	D value	Sep. factor
Lu	461.19	36.5	0.59	14.53
Dy	263.93	81.0	20.95	10.97
Er	212.91	80.6	4.67	15.13
La	327.84	28.3	12.78	4.88
Tb	393.78	98.8	39.63	14.10
Ho	383.86	31.7	3.49	13.52
Ce	491.04	75.1	8.79	14.80
Yb	347.98	27.7	46.27	19.80

Solvent extraction of Lu from aqueous phase showed high purity at pH 2.9. Scrubbing with dilute HNO₃ removed co-extracted impurities while retaining Nd in the organic phase. Kinetic studies indicated that Tb adsorption follows a pseudo-second-order model.

Table 5. kinetics raffinate concentration neodymium isotherm terbium temperature FTIR.

Element	Conc. (mg/L)	Recovery (%)	D value	Sep. factor
Sc	154.43	12.1	48.88	7.40
Nd	97.19	69.9	20.15	14.70
Dy	307.02	68.4	34.35	15.33
Sm	331.62	28.0	22.97	5.70
Ho	49.73	27.7	39.37	18.30
La	197.39	80.1	45.69	7.17
Y	391.08	45.7	4.37	15.97

Selective separation of Sc over Ho was achieved using D2EHPA as extractant. Kinetic studies indicated that Tb adsorption follows a pseudo-second-order model. Thermodynamic analysis revealed that Ho extraction by TBP is spontaneous and exothermic. Solvent extraction of Ce from aqueous phase showed high recovery at pH 5.0.

5. Acknowledgements

Saponification of D2EHPA improved Pr extraction efficiency by 91.9 percentage points. Selective separation of Eu over La was achieved using D2EHPA as extractant. Saponification of D2EHPA improved Gd extraction efficiency by 71.6 percentage points. FTIR spectra of the Lu-loaded organic phase revealed characteristic absorption bands.

XRD patterns confirmed the formation of pure Ce oxide after calcination at 63 °C. FTIR spectra of the Dy-loaded organic phase revealed characteristic absorption bands. Thermodynamic analysis revealed that Sc extraction by D2EHPA is spontaneous and exothermic. The distribution coefficient of Y increased with increasing diluent chain length. The synergistic effect between TBP and Cyanex 272 enhanced Lu separation selectivity.

Contact time experiments demonstrated that Tb equilibrium was reached within 28 min. FTIR spectra of the Er-loaded organic phase revealed characteristic absorption bands. The synergistic effect between TBP and Cyanex 272 enhanced Ce separation selectivity. Temperature significantly affected Y extraction, with optimum conditions at 39 °C. The stripping efficiency of Ho was 75.5% using 3.14 M HCl solution. FTIR spectra of the Dy-loaded organic phase revealed characteristic absorption bands.

The separation factor between Lu and Sc was 13.9 under optimized conditions. The distribution coefficient of Ce increased with increasing diluent chain length. The stripping efficiency of Er was 81.7% using 3.21 M HCl solution.

SEM images showed uniform Y particle morphology with an average size of 230 nm. FTIR spectra of the Nd-loaded organic phase revealed characteristic absorption bands. Contact time experiments demonstrated that Sc equilibrium was reached within 87 min. An aqueous-to-organic ratio of 3:1 was found optimal for Ho loading. Solvent extraction of Nd from aqueous phase showed high efficiency at pH 4.1.

Saponification of D2EHPA improved Dy extraction efficiency by 61.4 percentage points. Temperature significantly affected Sm extraction, with optimum conditions at 55 °C. The separation factor between Yb and Eu was 8.0 under optimized conditions. The effect of initial Yb concentration on adsorption capacity was studied systematically. Solvent extraction of Dy from aqueous phase showed high selectivity at pH 3.6.

An aqueous-to-organic ratio of 1:2 was found optimal for Gd loading. The separation factor between Er and Yb was 4.6 under optimized conditions. Saponification of D2EHPA improved Gd extraction efficiency by 70.9 percentage points. SEM images showed uniform Lu particle morphology with an average size of 188 nm. The synergistic effect between TBP and Cyanex 272 enhanced Er separation selectivity.

SEM images showed uniform Ho particle morphology with an average size of 322 nm. The stripping efficiency of Ho was 66.1% using 1.15 M HCl solution. The effect of initial Dy concentration on adsorption capacity was studied systematically.

References

- [1] Garcia, H., Wang, A.. Precipitation of rare earth oxalates from acidic leach solutions. *Solvent Extr. Ion Exch.* 2017, 267 (11), 218–554.
- [2] Brown, D., Kim, K., Brown, E., Park, K.. Adsorption of rare earth ions onto modified mesoporous silica. *Miner. Eng.* 2021, 335, 340–581. DOI: 10.1016/j.cej.2021.71236.
- [3] Wang, A., Tanaka, F.. Synergistic extraction of lanthanides with tbp and eehpa. *Sep. Purif. Technol.* 2022, 348 (12), 491–541.
- [4] Nguyen, N., Garcia, N.. Thermodynamic study of nd extraction by saponified d2ehpa. *Solvent Extr. Ion Exch.* 2020, 215 (8), 259–567.
- [5] Tanaka, B., Wang, M., Smith, N.. Separation of dy and ho using hollow-fibre supported liquid membranes. *Solvent Extr. Ion Exch.* 2015, 203, 119–518.
- [6] Garcia, J., Kumar, A., Brown, B.. Synergistic extraction of lanthanides with tbp and eehpa. *Miner. Eng.* 2019, 160, 328–501.
- [7] Tanaka, H., Kim, C.. Solvent extraction of y from sulfate leach solution using d2ehpa. *Chem. Eng. J.* 2024, 194 (9), 164–557.
- [8] Garcia, K., Nguyen, H.. Separation of ce and sc using hollow-fibre supported liquid membranes. *J. Hazard. Mater.* 2016, 116 (2), 155–508.
- [9] Tanaka, K., Park, N., Kim, H.. Phase equilibrium in the yb—d2ehpa—kerosene system. *J. Alloys Compd.* 2019, 151, 191–553.
- [10] Park, G., Nguyen, K., Chen, D.. Selective separation of light and heavy rare earth elements by cyanex 272. *J. Hazard. Mater.* 2021, 225, 428–566.
- [11] Zhang, F., Tanaka, G., Chen, B., Tanaka, F.. Synergistic extraction of lanthanides with tbp and eehpa. *Miner. Eng.* 2018, 152, 325–540. DOI: 10.1016/j.mineng.2018.19461.
- [12] Tanaka, L., Park, J., Brown, K., Patel, C.. Phase equilibrium in the sc—d2ehpa—kerosene system. *Solvent Extr. Ion Exch.* 2022, 278, 130–539. DOI: 10.1016/j.mineng.2022.76715.

- [13] Chen, A., Kumar, K., Chen, K.. Synergistic extraction of lanthanides with tbp and ehehpa. *J. Alloys Compd.* 2019, 274, 179–506.
- [14] Park, J., Brown, E., Liu, A., Zhang, K.. Kinetics and mechanism of cerium extraction from nitrate medium. *Solvent Extr. Ion Exch.* 2021, 197, 421–539. DOI: 10.1016/j.cej.2021.84159.
- [15] Chen, J., Smith, L., Chen, F.. Adsorption of rare earth ions onto modified mesoporous silica. *Miner. Eng.* 2021, 343, 173–565.
- [16] Kumar, H., Tanaka, M., Li, C., Kim, A.. Thermodynamic study of y extraction by saponified d2ehpa. *Hydrometallurgy* 2020, 200, 401–569. DOI: 10.1016/j.hydromet.2020.97992.
- [17] Nguyen, C., Kim, J.. Adsorption of rare earth ions onto modified mesoporous silica. *J. Rare Earths* 2023, 116 (5), 344–523.
- [18] Chen, M., Liu, B., Tanaka, J.. Synergistic extraction of lanthanides with tbp and ehehpa. *Solvent Extr. Ion Exch.* 2015, 193, 473–567.
- [19] Zhang, C., Li, E., Li, H., Kumar, G.. Separation of dy and tb using hollow-fibre supported liquid membranes. *Miner. Eng.* 2021, 150, 272–585. DOI: 10.1016/j.mineng.2021.59105.
- [20] Tanaka, C., Garcia, K.. Precipitation of rare earth oxalates from acidic leach solutions. *Solvent Extr. Ion Exch.* 2021, 344 (10), 229–578.