

1 Measurement of the neutrino magnetic moment at the
2 NOvA experiment

3 Technical note

4 Robert Kralik¹

5 ¹University of Sussex, Brighton, UK

6 January 20, 2024

7 **Abstract**

8 This is the abstract

9 Contents

| | | |
|----|---|-----------|
| 10 | 1 Introduction | 3 |
| 11 | 2 Analysis overview | 3 |
| 12 | 2.1 Datasets and Event Reconstruction details | 3 |
| 13 | 2.2 Analysis weights | 6 |
| 14 | 2.2.1 Neutrino magnetic moment signal as a weight | 8 |
| 15 | 2.3 Event selection | 8 |
| 16 | 2.4 Resolution and binning | 10 |
| 17 | 2.5 Systematic uncertainties | 10 |
| 18 | 2.6 Fitting framework | 12 |
| 19 | 3 Conclusion | 12 |

1 Introduction

(TO DO: Describe the main motivations for the analysis. Briefly mention that there was a previous study by Biao, what were the results there and what limitations (or maybe talk about this in the Experimental overview?))

2 Analysis overview

(TO DO: Describe the motivations for this analysis) What are we trying to achieve? Are we aiming for purity of the final sample or efficiency of the selection? We are trying to see the low energy excess of very forward neutrino-on-electron events. We can either do this via a counting experiment (possibly with various control samples/regions to control backgrounds and systematics), or with a fit to the energy spectrum of well-selected neutrino-on-electron events.

We are trying to select nu-on-e events with low electron recoil energies.

(TO DO: Briefly introduce what are we going to do with these events after the selection (fitting)) Are we just going to compare the event counts of signal and background (and possibly correct the background based on some other "sideband" selection?), or are we doing a fit to some spectra - either electron energy, angle or ETh2. - We do not know yet...

(TO DO: Describe what I'm talking about in this section (datasets, weights, selection, resolution, fitting framework).)

(TO DO: Describe already here that we're dividing the signal/background into four due to ...) Here on forward I'm going to describe the differences between these (definitions, weights, signal def, systematics. What is the same: event selection and binning. They're joint together in the fitting framework, where the ν_e CC MEC and the other backgrounds are simply summed together and scaled together. The ν -on-e background (also called the irreducible background by the LDM analysis) is treated/scaled separately.

(TO DO: Say here already that this is the same/similar events that are studied by the ND beam constraint and the LDM analyses)

(TO DO: Should I describe the NOvA Near Detector here? Specifically its capabilities for detecting electrons?)

(TO DO: List things that could be improved for this analysis over the presented technical note. Namely revisiting the event selection specifically for the neutrino magnetic moment events, especially the energy cut. Also including the anti-neutrino events and doing a joint fit.)

2.1 Datasets and Event Reconstruction details

For this analysis we are using near detector CAF samples with the standard Production 5.1 reconstruction.

The SAMWEB definition for the data sample is:

```
prod_caf_R20-11-25-prod5.1reco.1_nd_numi_fhc_full_v1_goodruns.
```

We are following the standard data blinding procedure and have not looked at any data events until the analysis passes the full collaboration review. The Near Detector group has validated **(TO DO: Figure out where did Yiwen and Wenjie actually look at the data)** using this data sample.

The exposure of the data sample is approximately 1.3848600×10^{21} POT. This is the exposure we use for all the following studies shown in this technical note. The Prod5.1 ND data sample contains data from run 10391 in epoch1a (2014-08-22 21:08:40) until run 14010 in epoch 11a (2021-02-03 15:48:21) (from the period and epoch naming wiki page <https://cdcvs.fnal.gov/redmine/projects/novaart/wiki>

(TO DO: Briefly describe the MC details. Versions of the individual simulation software) To tackle the low number of ν -on-e and ν_e CC MEC events in the nominal simulation sample we are using a suit of nominal and enhanced simulation samples for four different signal and background components. Each one contains its nominal sample and special systematically shifted samples for the detector systematics. The use of the samples is summarised in table 6 and described in detail below.

The GENIE tune is GENIE N1810j 02 11a (from the Prod5.Frankenstein docdb: 53360). The Genie release used for this production is R20-08-06-prod5.1genie.h, which has GENIE version V3.0.6 [1]

Also prod5.1 uses Geant4 v4.10.4.p02 [2]

We use the standard NOvA simulation (reference NOvA 3fl paper DOI: 10.1103/PhysRevD.106.032004)

Describe how did we deal with the GENIE skew fix. It was the GSF weights that forces "us" to treat the nueCC MEC background differently than the other background. As I understand it, the GSF is applied simply as the new weight. No change to the systematics is required.

Reference: A. Mislivic, "Genie skew reweight validation." NOvA Internal Document, DocDB: 553811 [from antinueCC IncXSec docdb:53691] "Final state kinematics were predicted by the N1810j 00 000 tune, but total cross section were generated with the intended N1810j 0211a179 tune, which differed in RES and DIS rates tuned to external data. Properly correcting the skew180 would require all simulation to be regenerated, so a temporary solution developed by the NOvA181 Cross-section Tuning Group involves reweighing production 5.1 events to the default N1810j 00 000.182 An additional modification to the GENIE MEC contribution are applied to better agree with NOvA183 ND data.

MC includes simulation in the rock surrounding the ND

(TO DO: Describe here that we're using the nominal ND MC sample for signal utilizing the simple relationship between the Standard Model cross section and the neutrino magnetic moment cross section (ref. theory))

(TO DO: Say already here that the POT inside the enhanced MC samples are not properly accounted for in CAFAna (Loader issue) and so the event counts need to be adjusted post-hoc)

Enhanced ν -on-e sample

(TO DO: Describe the nuone sample) Created by Wenjie Wu (was it just him or also Yiwen?) to do ... and fully described in the technote [3]. Using the overlayed and filematched samples for consistency.

(TO DO: Find a reference and reasoning for why Wenjie hasn't created the other systematics samples) We only have the selected few systematics definitions because ...

Table 1: Overview of the simulation samples corresponding to different signal and background components.

| | |
|---------------------------|--------------------------------|
| Signal | Enhanced ν -on-e sample |
| ν -on-e background | Enhanced ν -on-e sample |
| ν_e CC MEC background | Enhanced ν_e CC MEC sample |
| Other background | Nominal ND CAF sample |

(TO DO: Describe the differences)

- Missing cross section parameters - unable to use cross section weights or so
- Special mode for nu-on-e elastic scattering 10005

The list of the nu-on-e sample definitions is in table 2.

Table 2: SAMWEB definitions for the enhanced ν -on-e samples.

Nominal:

prod_caf_R20-11-25-prod5.1reco.g_nd_genie_N1810j0211a_nonswap_fhc_nova_v08_full_v1_nuone_overlay

Systematically shifted samples:

prod_caf_R20-11-25-prod5.1reco.g_nd_genie_N1810j0211a_nonswap_fhc_nova_v08_full_calibup_v1_nuone_overlay

prod_caf_R20-11-25-prod5.1reco.g_nd_genie_N1810j0211a_nonswap_fhc_nova_v08_full_calibdown_v1_nuone_overlay

prod_caf_R20-11-25-prod5.1reco.g_nd_genie_N1810j0211a_nonswap_fhc_nova_v08_full_ckvup_v1_nuone_overlay

prod_caf_R20-11-25-prod5.1reco.g_nd_genie_N1810j0211a_nonswap_fhc_nova_v08_full_ckvdown_v1_nuone_overlay

prod_caf_R20-11-25-prod5.1reco.g_nd_genie_N1810j0211a_nonswap_fhc_nova_v08_full_lightlevelup_v1_nuone_overlay

prod_caf_R20-11-25-prod5.1reco.g_nd_genie_N1810j0211a_nonswap_fhc_nova_v08_full_lightleveldown_v1_nuone_overlay

Enhance ν_e CC MEC sample

Created by Yiwen Xiao [3] to tackle the low statistics of the ν_e CC MEC background events and subsequently large and unphysical cross section weights.

(TO DO: List all the nueCC MEC sample definitions used. Do this after creating the filematched definitions maybe?)
 (TO DO: List the limitations of the sample in the q_3 - q_0 parameter space)

Table 3: SAMWEB definitions of the ν_e CC MEC background sample.

Nominal:

YiwenXiao_NueCCMEC_Single_NJobs7500_CAF_NonSwap

Systematically shifted samples:

YiwenXiao_NueCCMEC_Single_NJobs7500_CAF_NonSwap_CalibUp

YiwenXiao_NueCCMEC_Single_NJobs7500_CAF_NonSwap_CalibDown

YiwenXiao_NueCCMEC_Single_NJobs7500_CAF_NonSwap_CkvUp

YiwenXiao_NueCCMEC_Single_NJobs7500_CAF_NonSwap_CkvDown

YiwenXiao_NueCCMEC_Single_NJobs7500_CAF_NonSwap_LLP

YiwenXiao_NueCCMEC_Single_NJobs7500_CAF_NonSwap_LLDown

YiwenXiao_NueCCMEC_Single_NJobs7500_CAF_NonSwap_Aging

YiwenXiao_NueCCMEC_Single_NJobs7500_CAF_NonSwap_CalibShape

YiwenXiao_NueCCMEC_Single_NJobs7500_CAF_NonSwap_MCNP

Near Detector filematched CAF sample

(TO DO: describe and list all the ND nominal CAF samples)

The nominal ND MC includes 4x data POT. The systematics are file-matched to remove any statistical bias

2.2 Analysis weights

(TO DO: Describe why do we use weights) What are the weights we are using and why?

To correct for known deficiencies in simulation of neutrino flux or cross sections we apply weights calculated for each event.

Table 5 shows what CAFAna weights are used to simulate what signal/background sample.

PPFX weight

ana::kPPFXFluxCVWgt [4] (TO DO: What does this do (one sentence ish).) Maybe cite Leo's thesis? Or paper? L. Aliaga, "Neutrino Flux Prediction for the NuMI Beamline." PhD Thesis, FERMILAB-1081 THESIS-2016-03

Table 4: SAMWEB definitions of the other background samples.

Nominal:

prod_caf_R20-11-25-prod5.1reco.a_nd_genie_N1810j0211a_nonswap_fhc_nova_v08_full_v1

Systematically shifted samples:

prod_caf_R20-11-25-prod5.1reco.e_nd_genie_N1810j0211a_nonswap_fhc_nova_v08_full_calibup_v1_batch2

prod_caf_R20-11-25-prod5.1reco.e_nd_genie_N1810j0211a_nonswap_fhc_nova_v08_full_calibdown_v1_batch2

prod_caf_R20-11-25-prod5.1reco.f_nd_genie_N1810j0211a_nonswap_fhc_nova_v08_full_ckvup_v1_batch2

prod_caf_R20-11-25-prod5.1reco.f_nd_genie_N1810j0211a_nonswap_fhc_nova_v08_full_ckvdown_v1_batch2

prod_caf_R20-11-25-prod5.1reco.g_nd_genie_N1810j0211a_nonswap_fhc_nova_v08_full_lightlevelup_v1_batch2

prod_caf_R20-11-25-prod5.1reco.g_nd_genie_N1810j0211a_nonswap_fhc_nova_v08_full_lightlevelup_v1_batch2

prod_caf_R20-11-25-prod5.1reco.f_nd_genie_N1810j0211a_nonswap_fhc_nova_v08_full_detectorageing_v1_batch2

prod_caf_R20-11-25-prod5.1reco.f_nd_genie_N1810j0211a_nonswap_fhc_nova_v08_full_calibshape_v1_batch2

prod_caf_R20-11-25-prod5.1reco.g_nd_genie_N1810j0211a_nonswap_fhc_nova_v08_full_mcnp_v1_batch2

Table 5: Overview of CAFAna weights applied to each analysis sample.

| | |
|---------------------------|---|
| Signal | Flux and neutrino magnetic moment weights |
| ν -on-e background | Flux and radiative correction weights |
| ν_e CC MEC background | Flux and cross section weights |
| Other background | Flux and cross section weights |

Prod5.1 GSF XSec weight

ana::kXSecCVWgt2020GSFProd51 (*TO DO: Find the reference: possibly Maria's docdb:53336 together with the official 2020 XSec tuning technote docdb:43962.*)

NOvAReweight reference: J. Wolcott, "NOvARwgt software." <https://github.com/novaexperiment/NOvARwgt>

public.

(TO DO: Briefly describe what does this do. Also mention Yiwen's talk/technote about the large XSec weights that made her create an enhanced nueCC MEC sample.)

We are only using the for the background since we assume that the cross section for the signal is perfect. Also there are not weights for this kind of interaction.

Radiative correction weight

(TO DO: Why are we doing this? (reference Yiwen's talk/technote).)

Mention here where did I get the original GENIE cross section from (reference Yiwen's talk or technote, plus the original paper that was used). nu-on-e technote[3]

(TO DO: Write out the actual version of the weight. Including the original and the corrected XSec constants)

MINERvA paper: <https://journals.aps.org/prd/pdf/10.1103/PhysRevD.100.092001>

Say that we are not using the third part of the correction because it is tiny and it makes no difference. (tried and tested)

(TO DO: correct the equation) Calculated as

$$weight_{\text{Radiative Corr.}} = \frac{d\sigma_{\nu\text{-on-e}}}{dy} \Big|_{\text{Radiative Corr.}} / \frac{d\sigma_{\nu\text{-on-e}}}{dy} \Big|_{\text{GENIE 3}} ; y = \frac{E_e - m_e}{E_\nu} \quad (1)$$

2.2.1 Neutrino magnetic moment signal as a weight

(TO DO: What does this do and why does it work? Reference the theory part as to why is the magnetic moment signal simply a rescaling of the GENIE cross section.)

Using the same tree-level cross section from GENIE as in the rad. corr. weight.

(TO DO: Write the name of the weight in CAFAna/nuone namespace and where it is located)

(TO DO: correct the equation) Calculated as

$$weight_{\nu \text{ Mag. Moment}} = \frac{d\sigma_{\nu\text{-on-e}}}{dy} \Big|_{\nu \text{ Mag. Moment}} / \frac{d\sigma_{\nu\text{-on-e}}}{dy} \Big|_{\text{GENIE 3}} ; y = \frac{E_e - m_e}{E_\nu} \quad (2)$$

2.3 Event selection

Should this be a separate section or is it all right to keep it here? It will have a lot of plots...

(TO DO: Add the link to the LDM group's technote and say what's different (or maybe do this after we discuss the cuts?)) Currently we are using the exact same selection as is used by the ND group [3] and very similar to the Light Dark Matter analysis (cite their technote).

Signal definition

(TO DO: Define the signal of the NuMM. Reference the NuMM weight description above) The signal of the neutrino magnetic moment analysis is just a reweighted signal of the ν -on-e analysis from the near detector group. We are using the same event selection as the near detector group.

152 (TO DO: *Decide and explain what signal definitions we're using (kIsVtxContained VS Fiducial*
 153 *volume)* What is the signal and all the background samples definition? Difference between using
 154 kIsVtxCont and the fiducial volume. Is there a fundamental difference or preference? Or does it
 155 just depend on me? The results/counts are quite different...

Table 6: Overview of signal and background definitions. Mode 10005 denotes ν -on-e events, while mode 5 denotes all electron scattering events, including inverse muon decay interactions. That is why we had to add a requirement of an electron in the final state. Mode 10 denotes all MEC events. (TO DO: *Check that the definitions are correct from the code.*)

| | |
|---------------------------|---|
| Signal | <code>kMode== 10005 && NDNuoneFiducial</code> |
| ν -on-e background | <code>kMode== 10005 && NDNuoneFiducial</code> |
| ν_e CC MEC background | <code>!(kMode== 5 && kElInFinState && NDNuoneFiducial) && (kIsCC && kIsNue && kMode == 10)</code> |
| Other background | <code>!(kMode== 5 && kElInFinState && NDNuoneFiducial) !(kIsCC && kIsNue && kMode == 10)</code> |

156 Pre-selection

157 The pre-selection cuts have been kept from the ν_e CC analysis with loosened cut values (TO DO:
 158 *find a reference for this analysis*). Pre-selection cuts include basic quality cuts (TO DO: *describe*
 159 *the basic quality cuts that are implied from the preselection cuts*). They also remove the obvious
 160 ν CC interactions by requiring that the length of the longest prong is < 800 cm, number of planes
 161 crossed by the longest prong is < 120 , and the summed number of cells for all prongs in the slice is
 162 < 600 . In pre-selection we also include a cut on the time difference between the mean times of the
 163 "current" slice and of the slice closest in time, which should be > 25 ns. This ensures that ... (TO
 164 DO: *describe why do we need the closest slice cut with reference to Yiwen's talk and technote*).

165 Fiducial and containment cuts

166 (TO DO: *Describe what does the fiducial cut do*) We require that the reconstructed vertex is
 167 contained within the following volume: $-185 < V_{txX} < 175, -175 < V_{txY} < 175, 95 < V_{txZ} <$
 168 1095 cm.

169 To ensure all the energy is contained within the detector and to remove events originating
 170 outside of the detector (rock muons), we require that the extreme positions of hits for all prongs
 171 in the slice are within the following volume: $-190 < \min_X, \max_X < 180, -180 < \min_Y, \max_Y <$
 172 $190, 105 < \min_Z, \max_Z < 1275$ cm

173 Single particle requirement

174 To selection events with a single particle we require that the fraction of energy contained in the
 175 most energetic shower is > 0.8 , that the summed energy of all cells (above threshold and within ± 8

planes from the vertex) outside of the most energetic shower is < 0.02 GeV, and that the distance between the vertex and the start of the primary shower is < 20 cm.

Shower energy cut

(TO DO: discuss the energy cut, should this be removed? What is the effect on the event count? Why was this included in the first place (the identifiers are not as strong for lower energies - is this true though? - also there are further unexplored backgrounds that would need to be further studied and explore. Maybe depends on where would we move the cut...)) The calorimetric energy of the primary shower is required to be within $0.5 < E_{cal} < 5$ GeV.

Event classifiers

We are using two event classifiers based on convolution neural network that were developed specifically to identify ν -on-e interactions. The first one (NuoneID) is trained to select ν -on-e events and the second one (Epi0ID) is trained on the events passing the NuoneID to reject the π^0 background. Our selection requires that NuoneID > 0.73 and that Epi0ID > 0.92 .

(TO DO: reference theory for the kinematics of nuone scattering) We require that the product of reconstructed energy of the primary shower and the square of its angle from the Z axis is $E_{cal} \theta^2 < 0.005 \text{ GeV} \times \text{rad}^2$.

(TO DO: Add plots of distributions of the event selection variables with two columns. LHS shows no cuts applied and RHS shows all previous cuts applied)

Using the many plots below that show the effect of each of the cuts on the signal and all background events. (For signal we are showing NuMM=...)

(TO DO: Describe the cutflow tables below) The final event count and efficiency of each of the cuts is shown on the table 7. Table 1 shows the dissemination of background into the individual components.

(TO DO: Add a discussion of possible improvements on the event selection on its limitations - mostly for the analysis review committee) From here we can see that ... Maybe what can be improved is... This can likely be improved upon by specifically selection low energy events and removing the cut on the reconstructed shower energy.

2.4 Resolution and binning

(TO DO: Add the energy resolution and binning plots) The electron energy and angle distributions and resolutions. Are we going to fit in E, Th, or ETh2? Is there something else?

Show plots of Reco V True for both energy and angle. (Should I show it with or without the energy cut?). Also show the resolution plots.

2.5 Systematic uncertainties

(TO DO: Describe the main systematic uncertainties. Add plots showing their effect on the NuMM events. Possibly with different event selection variables as X axes. Also show the final table with the

percentages summed) Plots showing combined uncertainties for signal and backgrounds. Maybe also some interpolations. Table of systematic uncertainties on the event count.

Normalization systematics

(TO DO: Describe the normalization systematics (or just remove this if not using them in the end)) Should we include normalization systematics? Would that make any difference? There's a POT scaling uncertainty which is very small (find out exactly how small).

In the fitting experiment normalization uncertainties would probably not make any difference whatsoever, but in the counting experiment they might be important?

Neutrino flux systematics

(TO DO: Describe the flux uncertainties. Describe the PCA. Describe the difference between what the ND group is doing and what we're doing) Using the PCA vs using the PPFx universes+beam transport separately. Plots of energy showing shifts for signal and backgrounds separately

(TO DO: understand differences with ND and 3F methods)

This is mainly a normalization. Discuss how to use the fact that ν -on-e events can be used (and are used) to constraint the beam uncertainty. Would the counting experiment still be valid then?

Table 7: Event selection cutflow table

| Selection | ν Mag. Moment signal | | | ν -on-e background | | | Other background | | |
|------------------|--------------------------|------------------|----------------|------------------------|------------------|----------------|--------------------|------------------|-----------------------|
| | N_{sig} | ϵ^{N-1} | ϵ (%) | N_{IBkg} | ϵ^{N-1} | ϵ (%) | N_{Bkg} | ϵ^{N-1} | ϵ (%) |
| No Cut | 269.77 | 100 | 100 | 3.43×10^3 | 100 | 100 | 2.96×10^8 | 100 | 100 |
| Vtx Is Valid | 180.58 | 66.94 | 66.94 | 3.33×10^3 | 96.94 | 96.94 | 2.34×10^8 | 79.09 | 79.09 |
| N Prongs | 174.69 | 96.74 | 64.76 | 3.23×10^3 | 96.99 | 94.02 | 8.66×10^7 | 37.00 | 29.27 |
| Png Length | 174.67 | 99.99 | 64.75 | 3.22×10^3 | 99.64 | 93.68 | 7.67×10^7 | 88.56 | 25.92 |
| N Planes | 174.67 | 100 | 64.75 | 3.22×10^3 | 99.98 | 93.67 | 7.67×10^7 | 99.98 | 25.92 |
| N Cells | 174.67 | 100 | 64.75 | 3.22×10^3 | 99.98 | 93.65 | 7.42×10^7 | 96.78 | 25.08 |
| Closest Slc | 169.82 | 97.22 | 62.95 | 3.14×10^3 | 97.54 | 91.35 | 6.95×10^7 | 93.68 | 23.49 |
| Fiducial | 167.72 | 98.76 | 62.17 | 3.09×10^3 | 98.41 | 89.89 | 3.59×10^7 | 51.71 | 12.15 |
| Cont. | 159.37 | 95.02 | 59.08 | 2.48×10^3 | 80.43 | 72.30 | 1.38×10^7 | 38.35 | 4.66 |
| ShwE Frac. | 150.37 | 94.35 | 55.74 | 2.42×10^3 | 97.59 | 70.56 | 8.82×10^6 | 63.97 | 2.98 |
| Vtx E | 142.29 | 94.63 | 52.74 | 2.18×10^3 | 90.16 | 63.62 | 4.15×10^6 | 47.07 | 1.40 |
| Shw Gap | 137.96 | 96.96 | 51.14 | 2.09×10^3 | 95.58 | 60.80 | 3.25×10^6 | 78.34 | 1.10 |
| Shw E | 37.13 | 26.92 | 13.76 | 1.36×10^3 | 65.10 | 39.58 | 6.25×10^5 | 19.21 | 0.21 |
| Nuoneid | 29.48 | 79.39 | 10.93 | 940.21 | 69.18 | 27.38 | 2.42×10^4 | 3.88 | 8.19×10^{-3} |
| Epi0id | 22.51 | 76.35 | 8.34 | 749.93 | 79.76 | 21.84 | 1.47×10^4 | 60.75 | 4.97×10^{-3} |
| $E\theta^2$ | 19.74 | 87.73 | 7.32 | 675.02 | 90.01 | 19.66 | 84.15 | 0.57 | 2.84×10^{-5} |
| $E\theta^2$ (sb) | 2.74 | - | 1.01 | 74.30 | - | 2.16 | 1.01×10^3 | - | 3.43×10^{-4} |
| No ShwE | 37.62 | - | 13.94 | 782.67 | - | 22.79 | 238.79 | - | 8.07E-05 |

226 Maybe if we made another sideband sample...

227 **Detector systematics**

228 *(TO DO: Make plots of energy showing shifts for signal and backgrounds separately)* Reference
229 for the Prod5.1 detector systematics is docdb 53225

230 **Cross section systematics**

231 *(TO DO: Describe the XSec syts)* Only for the non nu-on-e background. Assuming the nu-on-e
232 events (including the signal events) are precisely known.

233 Plots of energy showing shifts for signal and backgrounds separately

234 **2.6 Fitting framework**

235 *(TO DO: Describe the fitting framework, ideally with an example plot, maybe with some arrows)*
236 How does the fitting framework work? It's based on the framework developed by Mu Wei for the
237 Light Dark Matter analysis (ref.) which was developed together (is this fair?). Basic description of
238 the framework.

239 Also this framework is used for both LDM and NuMM together. It is trivial to simply switch
240 between including the NuMM or LDM in it. This was done to save space in creating predictions
241 since our backgrounds are exactly the same (or at least they should be...). Theoretically this could
242 be separated into two difference frameworks.

- 243 • <NDPredictionSingleElectron> Prediction class which holds the LDM as a special 2-D spec-
244 trum (not used for NuMM), and NuMM, ν -on-e background, ν_e CC MEC background and
245 other background as simple 1-D spectra. Also scaling each spectra by...
- 246 • <NDPredictionSystSingleElectron> class derived from PredictionInterp that takes in the
247 NDPredictionSingleElectron and applies systematic shifts to it. Includes the interpolation/extrapolation
248 between the systematic shifts.
- 249 • FitVariables and what do they do
- 250 • Fitter which does exactly what... What are the parameters of the fit? What are the re-
251 sults/outputs?

252 **3 Conclusion**

253 *(TO DO: Report the limit with its uncertainty)*

254 *(TO DO: Very briefly discuss differences with current world limit and how the techniques*
255 *differ)*

256 *(TO DO: Very briefly summarise expectations for future measurements.)*

References

- [1] C. Andreopoulos et al. The GENIE Neutrino Monte Carlo Generator. *Nucl. Instrum. Meth. A*, 614:87–104, 2010. arXiv:0905.2517, doi:10.1016/j.nima.2009.12.009.
- [2] S. Agostinelli et al. GEANT4—a simulation toolkit. *Nucl. Instrum. Meth. A*, 506:250–303, 2003. doi:10.1016/S0168-9002(03)01368-8.
- [3] Wenjie Wu and Yiwen Xiao. Neutrino-Electron Elastic Scattering in the NOvA Near Detector - Technote. NOVA Document 56383, October 2023. NOvA technical note. URL: <https://nova-docdb.fnal.gov/cgi-bin/sso/ShowDocument?docid=56383>.
- [4] Leonidas Aliaga Soplin. PPFX tech-note for the 2017 analysis. NOVA Document 23441, November 2017. NOvA technical note. URL: <https://nova-docdb.fnal.gov/cgi-bin/sso/ShowDocument?docid=23441>.

Figure 1: Event selection outflow table for background components

| Selection | $V_e \text{CC MEC}$ | | | $V_e \text{CC Other}$ | | | $V_\mu \text{CC}$ | | | NC | | | Other | | |
|------------------|---------------------|------------------|----------------|-----------------------|------------------|-----------------------|--------------------|------------------|-----------------------|--------------------|------------------|-----------------------|--------------------|------------------|-----------------------|
| | N | ϵ^{N-1} | $\epsilon(\%)$ | N | ϵ^{N-1} | $\epsilon(\%)$ | N | ϵ^{N-1} | $\epsilon(\%)$ | N | ϵ^{N-1} | $\epsilon(\%)$ | N | ϵ^{N-1} | $\epsilon(\%)$ |
| No Cut | 3.50×10^4 | 100 | 100 | 3.23×10^6 | 100 | 100 | 2.24×10^8 | 100 | 100 | 3.40×10^7 | 100 | 100 | 3.49×10^7 | 100 | 100 |
| Vtx Is Valid | 3.27×10^4 | 93.58 | 93.58 | 2.62×10^6 | 81.14 | 81.14 | 1.99×10^8 | 89.02 | 89.02 | 2.57×10^7 | 75.55 | 75.55 | 6.53×10^6 | 18.70 | 18.70 |
| N Prongs | 2.74×10^4 | 83.76 | 78.39 | 1.39×10^6 | 53.05 | 43.05 | 6.75×10^7 | 33.89 | 30.17 | 1.51×10^7 | 58.57 | 44.25 | 2.65×10^6 | 40.51 | 7.58 |
| Png Length | 2.73×10^4 | 99.79 | 78.22 | 1.37×10^6 | 98.53 | 42.42 | 5.77×10^7 | 85.56 | 25.81 | 1.49×10^7 | 99.07 | 43.84 | 2.64×10^6 | 99.87 | 7.57 |
| N Planes | 2.73×10^4 | 99.99 | 78.22 | 1.37×10^6 | 99.99 | 42.41 | 5.77×10^7 | 99.98 | 25.81 | 1.49×10^7 | 100 | 43.84 | 2.64×10^6 | 100 | 7.57 |
| N Cells | 2.73×10^4 | 99.99 | 78.21 | 1.28×10^6 | 93.49 | 39.65 | 5.59×10^7 | 96.82 | 24.98 | 1.44×10^7 | 96.34 | 42.24 | 2.64×10^6 | 100 | 7.57 |
| Closest Slc | 2.73×10^4 | 99.79 | 78.05 | 1.21×10^6 | 94.25 | 37.37 | 5.33×10^7 | 95.40 | 23.84 | 1.35×10^7 | 94.17 | 39.77 | 1.43×10^6 | 54.22 | 4.10 |
| Fiducial | 1.39×10^4 | 51.12 | 39.90 | 6.30×10^5 | 52.10 | 19.47 | 2.60×10^7 | 48.77 | 11.62 | 8.25×10^6 | 60.99 | 24.26 | 1.05×10^6 | 73.53 | 3.02 |
| Cont. | 9.32×10^3 | 66.82 | 26.66 | 2.63×10^5 | 41.72 | 8.12 | 7.64×10^6 | 29.38 | 3.42 | 4.96×10^6 | 60.15 | 14.59 | 9.12×10^5 | 86.62 | 2.61 |
| ShwE Frac. | 9.20×10^3 | 98.70 | 26.32 | 1.95×10^5 | 74.39 | 6.04 | 4.82×10^6 | 63.10 | 2.15 | 2.97×10^6 | 59.78 | 8.72 | 8.28×10^5 | 90.81 | 2.37 |
| Vtx E | 5.92×10^3 | 64.33 | 16.93 | 6.05×10^4 | 30.96 | 1.87 | 1.97×10^6 | 40.79 | 0.88 | 1.36×10^6 | 45.75 | 3.99 | 7.62×10^5 | 92.03 | 2.18 |
| Shw Gap | 5.50×10^3 | 92.91 | 15.73 | 4.62×10^4 | 76.40 | 1.43 | 1.58×10^6 | 80.18 | 0.70 | 1.06×10^6 | 77.78 | 3.10 | 5.69×10^5 | 74.61 | 1.63 |
| Shw E | 3.62×10^3 | 65.81 | 10.35 | 1.12×10^4 | 24.15 | 0.35 | 4.38×10^5 | 27.80 | 0.20 | 1.71×10^5 | 16.15 | 0.50 | 1.28×10^3 | 0.23 | 3.68×10^{-3} |
| Nuoneid | 1.40×10^3 | 38.63 | 4.00 | 2.11×10^3 | 18.89 | 0.065 | 1.17×10^4 | 2.66 | 5.21×10^{-3} | 8.99×10^3 | 5.27 | 0.026 | 66.43 | 5.17 | 1.90×10^{-4} |
| Epi0id | 1.14×10^3 | 81.78 | 3.27 | 1.61×10^3 | 76.40 | 0.050 | 7.17×10^3 | 61.52 | 3.20×10^{-3} | 4.76×10^3 | 52.94 | 0.014 | 29.47 | 44.36 | 8.44×10^{-5} |
| $E\theta^2$ | 15.13 | 1.32 | 0.043 | 39.00 | 2.42 | 1.21×10^{-3} | 8.62 | 0.12 | 3.85×10^{-6} | 20.91 | 0.44 | 6.15×10^{-5} | 0.50 | 1.69 | 1.43×10^{-6} |
| $E\theta^2$ (sb) | 386.16 | - | 1.10 | 306.55 | - | 9.48×10^{-3} | 165.59 | - | 7.40×10^{-5} | 149.93 | - | 4.41×10^{-4} | 6.24 | - | 1.79×10^{-5} |
| No ShwE | 15.54 | - | 0.044 | 69.61 | - | 2.15×10^{-3} | 68.48 | - | 3.06×10^{-5} | 75.67 | - | 2.22×10^{-4} | 9.49 | - | 2.72E-05 |