Drimagic

APPLICATION: As growth supplement for better crop output.

GENERAL INFORMATION:

A hydrolysate of nutrients from certain plants produced employing a unique multi-level still fermentation technology. It contains a host of substances that help in uniform growth of plants with substantial increase in crop output. The product is organic.

DRIMAGIC

Description	:Black colour liquid / off white powder.
Odour	:Specific
Specific gravity	:1.03
Solubility in water	:Complete.
pH	:5-8
Storage	:48 months.

COMPOSITION

- Nicotinic acid :24 ppm
- Pyridoxine :28 ppm
- Thiamine :18 ppm
- Amino acids :5234 ppm
- Proteins :1400 ppm
- Carbodyrates :9800 ppm
- Sachharides :9600 ppm

N :0.103%
P :01.10%
K :00.06%
Mn :12 ppm
Cu :60 ppm
Zn :245 ppm

Since the product is a botanical extraction, the content levels may vary. All the contents are derived from plant sources.

MODE OF ACTION

Nicotinic acid, Pyridoxine & Thiamine	These ingredients promote metabolism, causes physiological reactions and protect chlorophyll level. Helps in breakdown of carbohydrates and proteins.
Amino acids – L-Lysine and Proteins	Plants assimilate L lysine amino acid, one of the essential amino acids for pollination. It increases the pollen germination and the length of the pollen tube. Proteins have structural metabolism transport and amino acid stock function.
Carbohydrates / Sachharides	Promotes photosynthesis. Provides energy through metabolism pathways.

DILUTION : 2 ml per liter of water.

Crop	Application	Advantages
Fruit crops	Before flowering. After small fruit setting.	Less flower/fruit drop. Improved shelf life. Better juice quality.
Cotton	Ten days before flowering. At the time of square formation.	Better ball setting. Higher yield and chlorophyll content.
Oil Seeds	40 days after sowing. 55 days after sowing. 70 days after sowing.	Improved yield. Higher oil content in seeds Higher weight of seeds
Pulses	At the time of flowering. At the time of fruit setting	Improved chlorophyll content. Less flower/fruit drop. Higher yield.
Cereals	20-30 days after sowing. 10 days after tillering.	Higher yield. Higher grain weight.
Flowers	Before budding. After bud setting at an interval of 2 days.	Intermode distance improves. Firm petals. Petal fall delayed. Longer life.
Spices	At the time of pruning. 10 days after first spray. 10 days after second spray.	Less flower and fruit drop. Higher yield. Better aroma.