PRODUCT CATALOG OF DBPL

FY 2009-10

DRISHTI BIOTECH

We , Drishti Biotech Pvt. Ltd.(DBPL), are a sister concern of JR Group Of Companies . we are a professionally managed company engaged in multiple activities of manufacturing , production and selling of Agri- Products such as Bio-Pesticides , field crop seeds, fodder seeds , Crop Growth Promoters, Vegetable Seeds , Agro – chemicals , Agri Commodities and Agribusiness Consultancy and services (ABCAS).

Our motto is to offer end to end solution to our clients, we have modern factories , one Godown and three offices in India to well facilitate our clients with our premium products. Currently, we are serving in five states including Gujarat , Maharashtra, Rajasthan, M.P. and Karnataka .Our skilled Agricultural and Biotechnology experts and scientist offer expert solutions for issues related to all agricultural & Agribusiness needs in highly professional & Competent ways to meet the customer expectation. Our mission is to become a world class service provider and supplier in the agri-business sector by offering an end to end solution package for the entire farming community to make them the stakeholders of the Agri business by changing their impression of traditional farmer, so that it can reap more production. This in turn would enable them to become stake holders and part of larger interest groups.

Promoted By

1. **HARIYANT C SHELEAT**- **CHAIRMAN**– A SCIENCE & LAW GRADUATE & A PROVEN PERSONALITY IN BUSINESS AND EXPERIENCE OF 34 YEARS IN BUSINESS IN INDIA.
2. **RAJESH JHA – MANAGING DIRECTOR**– A AGRICULTURE AND MANAGEMENT GRADUATE WITH 16 YEARS PROVEN EXPERTISE IN AGRIBUSINESS INSTITUTIONAL BUSINESS SECTOR.
3. **RAKESH JHA** – **CHIEF EXECUTIVE OFFICER**– A AGRICULTURE AND MANAGEMENT GRADUATE WITH 11 YEARS PROVEN EXPERIENCE OF MARKETING IN AGRIBUSINESS SECTOR IN AGROCHEMICAL, SEED BUSINESS, COMMODITY BUSINESS AND FINANCE BUSINESS
4. **DHIRENDRA VAISHNAV- EXECUTIVE DIRECTOR** – AGRICULTURE AND MANAGEMENT GRADUATE WITH 34 YEARS PROVEN EXPERIENCE IN SEED BUSINESS IN INDIA.
5. **JAIMIN PATEL- EXECUTIVE DIRECTOR** – SCIENCE AND MANAGEMENT GRADUATE WITH 8 YEARS EXPERIENCE IN COMMODITY AND INSURANCE BUSINESS.

Our Associated Companies

- Shubham Agribusiness & Financial Solution – Ahmedabad
- Chinmay Raj Biotech Pvt Ltd- Ahmedabad
- Scorpion Biotech Company – Ahmedabad
- Chandel Agribusiness solutions Pvt Ltd- Delhi
- Ambica Properties- Ahmedabad

Product Range

- **Products from Biotech Division:** DriMagic , DRIMAZER, Amazer, BioMagic ,Akti M3 Liquid, Green Magic ,Akti Gold H, Shaarp MB, Shaarp killer, Shaarp Shakti , Shaarp WF etc.
- **Products from seeds Division:**
  - **Field crop seeds:** Sesamum Seed, Maize, Bajra, Castor, Cotton, Rape Seeds
  - **Fodder Seeds:** Sorghum, African Tall Maize, Lucerne
  - **Vegetable Seeds :** Okra ,Hot pepper, Tomato, Cluster Bean, Coriander, Onion
- **Products of Commodity Division:** Wheat Paddy, Maize, Bajra, Jowar, Rice, Soyabean, Ground Nut, Jeera, Pulses, etc. with F&V like Fresh Potato, Onion and Mango
- **Products for Crop Protection:** Endosulfan, Monocrotophos, Imidacloprid, Chlorpyriphos, Glyphosate etc
- **Services :**
  - Contract Farming for Hi tech Agriculture and High Income Generation
  - Agribusiness Planning, Advisory, Implementation & Support
  - Commodity Handling, Procurement, financing and Disposal Services
  - Agri Project preparation, Implementation, Support and finance
  - Financial Advisory and Management for Corporate and SMEs
  - Need based Farming Projects for NGOs, Corporate and Institutions
  - Market Research and Documentation Support for NGO, Govt, and Industry
  - Market Promotion, Training, Motivation & Supports

DRISHTI BIOTECH PVT LTD

FF6, Maurya Arcade,
Opp SG Sola Highway, Sola Science City Road,
Sola, Ahmedabad 380061-GUJ-INDIA
TEL FAX : +91 79 4005 3080
Email : Group@drishtiagrotech.com

Corporate Office :

FF6, Maurya Arcade,
Opp SG Sola Highway, Sola
Science City Road, Sola,
Ahmedabad 380061-GUJ-INDIA
TEL FAX : +91 79 4005 3080

Web : http://www.drishtiagrotech.com
Email : group@drishtiagrotech.com
info@drishtibio.com
GST TIN No : 240 7360 5802
CST TIN No : 245 7360 5802

Quality Assurance:

Customers buy benefit and not products. Drishti Biotech Pvt Ltd. With its products and services sells benefit to its customers. The hardworking committed and disciplined staff that we have ensures that our clients get quality norms and standards are followed during the production process. We use quality raw materials to manufacture Agri-products so that customers can avail nothing but the best. Our firm belief is that quality is remembered longer. With our apt solutions, we cater almost all kinds of requirements of the market. Thus, we assure customers that our range of Agri- products is of world-class quality.

Infrastructure

Manufacturing bio-products requires big paraphernalia, including labs and dealing centres. We, Drishti Biotech Pvt Ltd, have two factories situated at VU Nagar & Khatraj. One Central Godown situated at Sarkhej and three offices situated at Ahmedabad, Delhi & Indore in India. We are based in the commercial city of Ahmedabad. We have full –challenges manufacturing units and office setups to effectively handle and meet rising challenges in Agri –business.

Product Range:

We manufacture a wide range of bio-products that lead to optimized production of crops. Our quality products incl-

- **Products from Biotech Division:** Drishti Magic , DRISHTIMAZER, Amazer, BioMagic ,Akti M3 Liquid, Green Magic ,Akti Gold H, Shaarp MB, Shaarp killer, Shaarp Shakti , Shaarp WF , Dhan varsha etc.
- **Products from seeds Division:**
  - **Field crop seeds:** Wheat –Drishti Suruchi, paddy-Drishti Shalimar, Drishti Guj-2, Maize –Drishti 121, Bajra –Drishti 4444, Castor- Drishti Avinash Cotton- Drishti 505, Rape Seeds – Drishti Zenith Black etc.
  - **Fodder Seeds:** Sorghum- Drishti Red Seeds – Drishti Red Diamond, African Tall Maize, Lucerne
  - **Vegetable Seeds :** Okra –Drishti -707,Hot pepper- Drishti 555, Tomato-Drishti 111,Cluster Bean-Drishti 999, Coriander S – Drishti Jinnat , Onion Seed- Drishti Selection 1 etc.
- **Products of Commodity Division:** Wheat Paddy, Rice, Soyabean, Ground Nut in Shell, Jeera, Pulses, etc.
- **Products for Crop Protection:** Acetamaprid 20% SP, Carbendazim 50%, Thiram 75% WS, Imidacloprid17SL etc
- **Agribusiness Consultancy & Services:**
  - Bulk Procurement of Agri commodities like Wheat , Paddy, RM Seed ,Castor Seed , Soyabean ,Ground nut, Maize etc. from Gujarat , Rajasthan , MP,CG,MS, Bihar, UP, etc. from Farmers or Trade.
  - Bulk supply of Agri Commodities like pulses, Rice, Wheat, Paddy, RM Seed, Soyabean, Jeera, Fennel,
  - Liasioning for Bulk & institutional Trade of any Agri Commodities as well as Agri inputs.
  - Contract Farming and Process Management of any Agri Produce in the state of Gujarat , RJ, MP, MS

Our Associated Companies:

- Chinmay Raj Biotech Pvt Ltd- Ahmedabad
- Scorpion Biotech Company – Ahmedabad
- Chandel Agribusiness solutions Pvt Ltd- Delhi
- JR Enterprises – Ahmedabad
- Ambica Properties- Ahmedabad

Corporate Office :

FF6, Maurya Arcade,
Opp SG Sola Highway, Sola
Science City Road, Sola,
Ahmedabad 380061-GUJ-INDIA
TEL FAX : +91 79 4005 3080

Web : http://www.drishtiagrotech.com
Email : group@drishtiagrotech.com
info@drishtibio.com
GST TIN No : 240 7360 5802
CST TIN No : 245 7360 5802

1. BIO MAGIC

A revolutionary Plant growth promoter - **Increases growth and productivity of Plants by increasing the Chlorophyll**

In this world more than 95% food products are produced by using the unique natural photosynthesis process of plants. Three main factors of Photosynthesis are CO2, Sunlight & Chlorophyll, out of this Sunlight & CO2 is available in abundantly in the nature but the third factor **Chlorophyll** is lacking in plants many times due to several reasons. This results in retarded growth of root & shoots system in plant / crop. Out of several major problem is chlorosis. BIOMAGIC is made by studying all the major problems of plant and is developed by using the methods and research of Biotechnology to improve the Chlorophyll contents in plant, which ultimately increases the photosynthesis in Crop/ Plants. Main composition of BIOMAGIC is Plant Extracts, Plasticizers, Surfactants and carriers.

Benefits of Using BIOMAGIC:

- Increases Chlorophyll contents in plant
- Improves photosynthesis process in plants
- Helps plants for better intake of all essential micronutrients from soil
- Improves root & shoot development
- Improves vegetative & reproductive growth of the plant as & when required
- Increases flower development & restricts flower shedding from plants/crops
- Improves fruit setting in plants and helps in restricting fruit shedding from plants
- Improves fruit/seed size, weights and numbers
- Improves quality of produce
- Improves Oil, protein & carbohydrate contents in Ground nut
- Improves intake of nitrogen from soil to plants
- Improves Disease & pest resistance in plants by making the plants stronger
- Increases the average yield of crop by 15 to 20%

How to Use: Dilute BIOMAGIC Paste in separate container in water smoothly by your hand

CROP WISE USE OF BIOMAGIC: BIOMAGIC will be used at the dose of 1 gram per litres of water (300 gram per 300 litres of water is recommended per acres for best results)

Groundnut	First Spray 30 DAS, 2nd and 3rd sprays in 12 to 16 days after first spray
Paddy/ Rice	First spray 15 days after transplanting, and second spray just after weeding operations
Potato	1st spray after 15 days of germination, then afterward every 10th day 2nd & 3rd spray to be done
Chilly, Tomato, Brinjal	First spray before flowering, second & third sprays in 15 days interval. For hybrid vegetable 5 sprays are recommended
Cotton	Onset of flowering in interval of 15 days 3 sprays. For Hybrids 4 to 5 sprays are recommended
Roses & All Flowers	After Monsoon (Sept) first spray. Afterward in interval of 15 days 2 to 3 sprays
Sugarcane	After germination, after 2 months on 15 days interval 3 to 4 sprays
Grape	After Pruning on set of new shoots first spray to be done. Total 4 to 5 sprays required
Pomegranate, Mango & other fruits	As per growth of plants. Before onset of monsoon in June months in 12 to 16 days intervals 4 to 6 sprays to be done
Wheat	First spray before weeding, then on every 15 days interval 2 additional sprays recommended
Jowar, Bajri, Maize, Rajko	15 days after germination, total 3 sprays in 12 to 16 days interval
Tea	After Pruning on set of new shoots first spray to be done. Total 4 to 5 sprays required
Cumin, fennel, Fenugreek	10 days after germination, total 3 sprays in 12 to 16 days interval
Sapota, Lime, banana, etc horticultural crops	As per growth of plants. Before onset of monsoon in June months in 12 to 16 days intervals 4 to 6 sprays to be done

Corporate Office :

FF6, Maurya Arcade,
Opp SG Sola Highway, Sola
Science City Road, Sola,
Ahmedabad 380061-GUJ-INDIA
TEL FAX : +91 79 4005 3080

Web : http://www.drishtiagrotech.com
Email : group@drishtiagrotech.com
info@drishtibio.com
GST TIN No : 240 7360 5802
CST TIN No : 245 7360 5802

2. AMAZER

(A high end plant growth promoter)

AMAZER, is a biotechnological technical product developed by Drishti Biotech Pvt Ltd. AMAZER is actually derived from Seaweed *Ascophyllum nodosum* with special fermentation process which contain readly to use natural nutrients like gibbrellines, auxin, enzymes, vitamins, cytokinins and other minerals to be used for liquid growth promoter for foliar spray. AMAZER is a unique product with very low dosage is needed for large crop cultivation area.

Ingredients : *Ascophyllum nodosum* 22 %, Protein hydrolysate (60%) 9 %, Alloe vera (88%) QS with other adjuvants.

AMAZER is eco-friendly, non-toxic, non-chemical, organic storehouse of naturally occurring growth-promoting substances in a very concentrated form. The Cytokinins, Auxins and Gibbellines with other are in natural form and are readily assimilated which helps to improve cell division and cell enlargement thus resulting in better chlorophyll content and increased production. The constitutes of AMAZER induces faster and vigorous development of root mass which in turn improves the suction of previously unavailable nutrients and other substances from soil for the plant to use. Gibberellins, Enzymes and Hydrolysed protein helps in cell growth and break down of complex molecules in to simpler ones which can be utilized by plants and also improves the soil microbe activity.

AMAZER is a natural plant food, which can be applied as a foliar feed, soil application, root dip or seedling dip or as a seed treatment. This type of application enables plant to receive direct benefits when the plant needs them especially at the time of Biomass development, Vegetative growth, C/N ratio differentiation, reproductive stage and Maturity.

Benefits : Useful for all the crops as Plant growth promoter, it increases the root development, nutrient uptake capacity, retention of flowers and fruits, fruits are bigger and shining that produce keeping quality. Very low dose per acre.

Dosage : 3 to 5 ml per 15 lits of water (100 ml per 300 lits of water)

Corporate Office :

FF6, Maurya Arcade,
Opp SG Sola Highway, Sola
Science City Road, Sola,
Ahmedabad 380061-GUJ-INDIA
TEL FAX : +91 79 4005 3080

Web : http://www.drishtiagrotech.com
Email : group@drishtiagrotech.com
info@drishtibio.com
GST TIN No : 240 7360 5802
CST TIN No : 245 7360 5802

3. DRIMAZER:

APPLICATION : As premium growth stimulant.

GENERAL: This formulation is a combination of high quality marine protein, chelated B vitamins and nutrients from rare herbs. The product is a unique and energy packed one that can enhance the growth uniformly resulting in higher crop output.

Description	:	Light cream colour powder / Black liquid
Odour	:	Specific.
Solubility in water	:	Complete.
pH	:	6-8
Shelf life	:	36 months.

COMPOSITION

Soluble solids, containing plant extracts, proteins, carbohydrates and inherent nutrients derived from fish & vegetable source and B vitamins.	:	32%
Media (stabilizers, preservatives and aqueous diluent)	:	68%

BENEFITS

- Promotion of cell division and cell elongation, protein synthesis; root initiation and development.
- Improves yield and shelf life of output.
- Helps to withstand stress conditions.
- Improves rate of fertilizer absorption.
- Helps growth retarded plants to spring into its full potency and yielding level.

DILUTION : 2 ml per liter of water.

Recommendations

Crop	Applications	Advantages
Fruit crops	• Before flowering. • After small fruit setting.	• Less flower/fruit drop. • Improved shelf life. • Better juice quality.
Cotton	• Ten days before flowering. • At the time of square formation.	• Better ball setting. • Higher yield and chlorophyll content.
Oil Seeds	• 40 days after sowing. • 55 days after sowing. • 70 days after sowing.	• Improved yield. • Higher oil content in seeds • Higher weight of seeds
Pulses	• At the time of flowering. • At the time of fruit setting	• Improved chlorophyll content. • Less flower/fruit drop. Higher yield.
Cereals	• 20-30 days after sowing. • 10 days after tillering.	• Higher yield. • Higher grain weight.
Flowers	• Before budding. • After bud setting at an interval of 2 days.	• Intermode distance improves. • Firm petals. • Petal fall delayed. Longer life.
Spices	• At the time of pruning. • 10 days after first spray. • 10 days after second spray.	• Less flower and fruit drop. • Higher yield. • Better aroma.

Corporate Office :

FF6, Maurya Arcade,
Opp SG Sola Highway, Sola
Science City Road, Sola,
Ahmedabad 380061-GUJ-INDIA
TEL FAX : +91 79 4005 3080

Web : http://www.drishtiagrotech.com
Email : group@drishtiagrotech.com
info@drishtibio.com
GST TIN No : 240 7360 5802
CST TIN No : 245 7360 5802

4. DRIMAGIC

APPLICATION : As growth supplement for better crop output.

GENERAL INFORMATION: A hydrolysate of nutrients from certain plants produced employing a unique multi-level still fermentation technology. It contains a host of substances that help in uniform growth of plants with substantial increase in crop output. The product is organic.

Description	:	Black colour liquid / off white powder.
Odour	:	Specific
Specific gravity	:	1.03
Solubility in water	:	Complete.
pH	:	5-8
Storage	:	48 months.

COMPOSITION

Nicotinic acid	:	24 ppm	N	:	0.103%
Pyridoxine	:	28 ppm	P	:	01.10%
Thiamine	:	18 ppm	K	:	00.06%
Amino acids	:	5234 ppm	Mn	:	12 ppm
Proteins	:	1400 ppm	Cu	:	60 ppm
Carbodyrates	:	9800 ppm	Zn	:	245 ppm
Sachharides	:	9600 ppm			

Since the product is a botanical extraction, the content levels may vary. All the contents are derived from plant sources.

MODE OF ACTION

Nicotinic acid, Pyridoxine & Thiamine	These ingredients promote metabolism, causes physiological reactions and protect chlorophyll level. Helps in breakdown of carbohydrates and proteins.
Amino acids – L-Lysine and Proteins	Plants assimilate L lysine amino acid, one of the essential amino acids for pollination. It increases the pollen germination and the length of the pollen tube. Proteins have structural metabolism transport and amino acid stock function.
Carbohydrates / Sachharides	Promotes photosynthesis. Provides energy through metabolism pathways.

DILUTION: 2 ml per liter of water.

Recommendations

Crop	Applications	Advantages
Fruit crops	• Before flowering. • After small fruit setting.	• Less flower/fruit drop. • Improved shelf life. • Better juice quality.
Cotton	• Ten days before flowering. • At the time of square formation.	• Better ball setting. • Higher yield and chlorophyll content.
Oil Seeds	• 40 days after sowing. • 55 days after sowing. • 70 days after sowing.	• Improved yield. • Higher oil content in seeds • Higher weight of seeds
Pulses	• At the time of flowering. • At the time of fruit setting	• Improved chlorophyll content. • Less flower/fruit drop. Higher yield.
Cereals	• 20-30 days after sowing. • 10 days after tillering.	• Higher yield. • Higher grain weight.
Flowers	• Before budding. • After bud setting at an interval of 2 days.	• Intermode distance improves. • Firm petals. • Petal fall delayed. Longer life.
Spices	• At the time of pruning. • 10 days after first spray. • 10 days after second spray.	• Less flower and fruit drop. • Higher yield. • Better aroma.

Corporate Office :

FF6, Maurya Arcade,
Opp SG Sola Highway, Sola
Science City Road, Sola,
Ahmedabad 380061-GUJ-INDIA
TEL FAX : +91 79 4005 3080

Web : http://www.drishtiagrotech.com
Email : group@drishtiagrotech.com
info@drishtibio.com
GST TIN No : 240 7360 5802
CST TIN No : 245 7360 5802

5. DRIMINO

APPLICATION : As a growth regulator.

GENERAL: The amino acids are manufactured employing indigenously developed yet sophisticated fermentation method. The amino acids are in the L form and are biologically pure and are in Free State. The active nature of amino acids helps it to get chelated and absorbed in the plant system. It consists of all essential amino acids. Amino Acid is an organic nutritional product derived from vegetable source (SOYA) required for plant growth promotion and for higher yields. It promotes hormonal activity and induces their synthesis resulting into flowering and fruit setting. It also reduces the flower and fruit drop caused by hormonal imbalance. It enhances nutritional uptake and their utilization which improves the colour, size, luster and taste of fruits. It also improves keeping quality. It is an ideal product to resist from adverse conditions. The foliar application enables the plants to receive direct benefits from naturally balanced nutrients / growth stimulating substances when they are needed by the plants especially at the time of stage differentiation. It is extracted from naturally occurring cereal protein source, the most biologically active for crops as a source of protein.

Description	:	Dark brownish yellow colored powder.
Odour	:	Specific, spicy.
Solubility in water	:	Complete (99%).

COMPOSITION

Total free amino acids	:	50% w/v (+-).
Loss on drying	:	05% max
Organic nitrogen	:	6.0% w/v
NH CL	:	45% w/v max
Chloride	:	1.5% w/v max.
Ash	:	04% w/v max
Storage stability	:	3 years.

Amino Acids	*g/100g protein *(an approximate table)*		
L-Alanine	2.20	L-Arginine	5.20
L-Aspartic Acid	3.45	L-Cysteine	2.45
L-Glutamic	9.12	L-Glysine	2.43
L-Hystidine	0.90	L-Isoleucine	1.70
L-Leucine	2.80	L-Lysine	2.30
L-Methionine	0.23	L-Phenylalanine	1.78
L-Proline	3.50	L-Serine	4.49
L-Threonine	3.56	L-Tyrosine	1.02
L-Valine	3.10		

Mode of Action

- Promotes uptake of macro and micro nutrients.
- Promotes enzymes activity by acting as natural stimulant
- Stimulates protein synthesis as it contains Amino Acids which are Building Blocks.
- Induces Flowering and Fruit set related hormones.
- Improves the efficiency of the pesticides and fungicides with synergism.
- Provides drought resistance by its direct action.

DILUTION : 2-3 gms per litre of water.

Application:

Foliar Application	:	2-3 gms / litre of water
Soil Application	:	Dilute it as 1:3000-4000 in water.

It can be mixed with most nutrient solutions and pesticides. Nevertheless make a miscibility test before first application.

Corporate Office :

FF6, Maurya Arcade,
Opp SG Sola Highway, Sola
Science City Road, Sola,
Ahmedabad 380061-GUJ-INDIA
TEL FAX : +91 79 4005 3080

Web : http://www.drishtiagrotech.com
Email : group@drishtiagrotech.com
info@drishtibio.com
GST TIN No : 240 7360 5802
CST TIN No : 245 7360 5802

6. DRIBT

APPLICATION : As special growth supplement for Bt. Cotton.

GENERAL: DRIBT is a specialty formula made to fulfil the high nutritional demand by Bt cotton. The nutritional requirement by Bt. cotton is high compared to standard cotton. Growth of Bt. cotton is vigorous demanding more nutrition. Additional supply of nutrition is necessary to improve the Bt. gene efficacy. DRIBT is a truly novel formula manufactured by combining botanical nutrients, marine protein and animal protein employing certain scientific as well as traditional methods, to obtain optimum crop output.

Description	:	Dark brown colour liquid / black liquid
Odour	:	Specific
Solubility in water	:	Complete
pH	:	6-8
Shelf life	:	36 months.

COMPOSITION

Soluble solids, containing plant Extract, proteins, carbohydrates And inherent nutrients Contained in products of Vegetable, fish and animal Origin	:	30%
Stabilizers, preservatives and Aqueous diluents	:	70%

BENEFITS

- Improved plant vigor.
- Better Bt gene expression.
- Uniform and increased boll size in cotton.
- Better drought tolerance and resistance to diseases.

DILUTION : 2 ml per litre of water.

RECOMMENDATION : As a special spray for Bt grade cotton.

7. DRIZYME

APPLICATION : To be used to incorporate in soil application bentonite granules.

GENERAL

The formulation is in line with BF1, but in a suitable form to be incorporated in bentonite granules.

Description	:	Black colour liquid.
Odour	:	Specific
Specific gravity	:	1.03
Solubility in water	:	Complete.
pH	:	5-8
Storage	:	48 months.

Note : On purchase, we will provide details of manual method of processing bentonite granules with DRIZYME.

Corporate Office :

FF6, Maurya Arcade,
Opp SG Sola Highway, Sola
Science City Road, Sola,
Ahmedabad 380061-GUJ-INDIA
TEL FAX : +91 79 4005 3080

Web : http://www.drishtiagrotech.com
Email : group@drishtiagrotech.com
info@drishtibio.com
GST TIN No : 240 7360 5802
CST TIN No : 245 7360 5802

8. DRISTICKER

APPLICATION : A wetting and sticking agent.

GENERAL INFORMATION: A unique wetting and sticking agent made of soluble polymers and surfactants to enhance compatibility with a broad spectrum of agricultural chemicals over wide varieties of water qualities

Description	:	White colour liquid.
Odour	:	Specific.
Storage stability	:	Minimum 36 months

COMPOSITION

Soluble Polymers & surfactants	:	90.00% v/v min.
Other ingredients	:	10.00% v/v max.

PROPERTIES

1. Stable formula.
2. With lesser surface tension, it enables free flow of spray liquid by allowing droplets to stretch out on the target surface, thereby giving better coverage.
3. It holds the spray fluid for a longer period of time on leaves and plant surfaces.
4. Non-toxic even at higher concentration.
5. With low vapour pressure, it reduces the evaporation losses.
6. Enhances performance of fungicide and insecticides.
7. Compatible with all commonly used insecticides, acaricides and fungicides.

DOSAGE : 1 ml per litre of water.

USAGE: Prepare the pesticide spray solution followed by addition of this spreading and sticking agent. Ensure that the entire solution is mixed properly to get a homogeneous solution before spraying.

Corporate Office :

FF6, Maurya Arcade,
Opp SG Sola Highway, Sola
Science City Road, Sola,
Ahmedabad 380061-GUJ-INDIA
TEL FAX : +91 79 4005 3080

Web : http://www.drishtiagrotech.com
Email : group@drishtiagrotech.com
info@drishtibio.com
GST TIN No : 240 7360 5802
CST TIN No : 245 7360 5802

9. DRIFUNGI

APPLICATION : An effective alternative to synthetic fungicides.

BIO-EFFICACY : University of Agricultural Sciences at Bangalore has successfully evaluated the performance of this formula. (Report will be made available on demand.)

GENERAL INFORMATION
Alkaloids are synthesized by certain plants, which are 'potential natural *insecticides*'. This formulation contains a complex of alkaloids isolated from adhatoda vasica and lantana camara employing indigenously developed extraction technology. Surface-active agents and carbonates help the formulation enhance its efficacy when applied.

Description	:	Grey/off-white powder / clear liquid
Odour	:	Specific.
Storage stability	:	Minimum 24 months

COMPOSITION

Complex of alkaloids (Min.)	:	01.00%
Surface active agents & Carbonates (Max.)	:	99.00%

MODE OF ACTION
The formulation disperses, spreads and activates. Alkaloids are physiologically active. It blocks the metabolic functions of pathogens, thereby rendering them lifeless. The surface-active agents help the spray spread evenly and activate the functions of alkaloids. It also helps the spray remain active for longer time. Apart from the above, the growth promoting substances in the botanical nutrients work as a tonic rejuvenating the effected plants instantly to a healthy and productive state. It acts as protective as well as curative. Applications should be repeated regularly to replace the formulation that has been washed or braded away and to protect foliage produced since last application.

DILUTION : 2-3 gms / ml per litre of water depending on the infestation rate.

RECOMMENDATION

Crop	Application	Targets
Paddy	2-3 applications 25-30 days after transplanting; repeat after one month	Leaf spot, blast, bacterial leaf blight.
Chilies	3-4 applications 30 days after transplanting; repeat after 30 days.	Leaf spot, leaf curling, powdery mildew.
Tomato	3-4 applications	Leaf spot, damping off.
Potato	2-3 applications.	Late blight.
Grapes	3-4 applications.	Powdery mildew.
Mango	2 applications.	Powdery mildew.
Pomegranate	3-4 applications.	Leaf spot, fruit rot, bacterial wilt.
Vegetables	3-4 applications.	Leaf spot, fruit rot, powdery mildew.
Spices	2-3 applications.	Capsule rot, root rot, wilt.
Floriculture	2-3 applications.	Powdery mildew.

Corporate Office :

FF6, Maurya Arcade,
Opp SG Sola Highway, Sola
Science City Road, Sola,
Ahmedabad 380061-GUJ-INDIA
TEL FAX : +91 79 4005 3080

Web : http://www.drishtiagrotech.com
Email : group@drishtiagrotech.com
info@drishtibio.com
GST TIN No : 240 7360 5802
CST TIN No : 245 7360 5802

10.DRIDM:

APPLICATION : As anti-fungal agent that controls diseases like Downy mildew etc.

BIO-EFFICACY: Satisfactory results were obtained during the bio-efficacy trials by the research wings

GENERAL INFORMATION: It works based on a synergistic effect by various components that provide a methodological control of the causal organisms. The contributive ingredients are alkaloids sourced from plant sources that are physiologically active, soluble polymers, sulphonated natural oils, salt of fatty acids and hydroxyl derivatives of aromatic hydrocarbons.

Description : Dark Red liquid
Odour : Specific.
Storage stability : Minimum 24 months

COMPOSITION
A suspension of alkaloids, soluble polymers, sulphonated natural oils, salt of fatty acids and derivatives of aromatic hydrocarbons. : 100.00%

MODE OF ACTION
The formulation disperses, spreads and activates. Alkaloids are physiologically active. It blocks the metabolic functions of pathogens, thereby rendering them lifeless. The surface-active agents including salt of fatty acids and sulphonated natural oils help the spray spread evenly helping to cover the affected areas and to activate the functions of alkaloids. The hydrocarbon derivatives burn the body tissues of the organisms. It also helps the spray remain active for longer time. It acts as protective as well as curative. Applications should be repeated regularly to replace the formulation that has been washed or braded away and to protect foliage produced since last application.

DILUTION : 2 ml per litre of water depending on the infestation rate.

RECOMMENDATION

Crops	*Targets*
Vegetables, chilli, grapes, beans, potato, tomato, onion, apple etc.	*Downy mildew, Leaf curl, late blight and early blight.*

CONTROL: Application of DRIDM controls DM and other such diseases effectively.

Corporate Office :

FF6, Maurya Arcade,
Opp SG Sola Highway, Sola
Science City Road, Sola,
Ahmedabad 380061-GUJ-INDIA
TEL FAX : +91 79 4005 3080

Web : http://www.drishtiagrotech.com
Email : group@drishtiagrotech.com
info@drishtibio.com
GST TIN No : 240 7360 5802
CST TIN No : 245 7360 5802

11. DRIANTIFID

APPLICATION : As antifeedant for chewing and sucking pests.

BIO-EFFICACY: Satisfactory results were obtained during the bio-efficacy trials by our research wings

GENERAL

It is an effective formula to control chewing and sucking pests on cereal and similar crops, based on natural bitters and castor extract incorporated in surfactant media. There is no toxic ingredient in the formula and as such it is totally safe to human, pets and the environment. It effectively controls BROWN PLANT HOPPER on paddy.

Description	:	Viscous liquid.
Odour	:	Specific.
Storage stability	:	Minimum 24 months

COMPOSITION

Natural bitters suspended in natural oils	:	25.00% (min.)
Salts of unsaturated aliphatic carboxylic acids	:	25.00% (min.)
Emulsifiers & media	:	50.00% (max.)

MODE OF ACTION: Insects do not feed on the crop due to its antifeedent properties and they leave the plants quickly

DILUTION : 3 - 4 ml per liter of water.

RECOMMENDATION

Crop	*Target Pests*
Paddy	Brown Plant Hopper
Cotton	Mealy bugs, Aphids & Ash weevil.

12. DRIBACT

APPLICATION : An effective organic bactericide.

BIO-EFFICACY : The bio-efficacy of DRBACT has been successfully tried and tested in the field some time ago by the company representatives.

GENERAL INFORMATION : Alkaloids are mildly toxic and are effective against a number of bacteria.

Description	:	Grey/off-white powder
Odour	:	Specific.
Storage stability	:	Minimum 24 months

COMPOSITION

Group of alkaloids derived from Lantana Camara, Catharanthus Roseus (sadaphuli)	:	1.00% w/w min.
Associates derivatives, fatty acids based Surface active agents & functional fillers	:	99.00% w/w max.

MODE OF ACTION

This formulation disperses, spreads and activates. Alkaloids from all the above sources have powerful anti-bacaterial properties that physically disrupt the causal bacteria. It may be used in both curative and preventive applications – curative because it kills pathogens on contact, preventive because its applications will effectively lower disease pressure.

DILUTION : 2-3 gms per litre of water depending on the infestation rate.

RECOMMENDATION: Against bacterial speck, bacterial spot, grey mold & white patches on tomato, fruits, onion, chilli, paddy and other crops.

Corporate Office :

FF6, Maurya Arcade,
Opp SG Sola Highway, Sola
Science City Road, Sola,
Ahmedabad 380061-GUJ-INDIA
TEL FAX : +91 79 4005 3080

Web : http://www.drishtiagrotech.com
Email : group@drishtiagrotech.com
info@drishtibio.com
GST TIN No : 240 7360 5802
CST TIN No : 245 7360 5802

13. DRIMITI

APPLICATION: As an alternative to synthetic Acaricide to control red mites and mealy bugs.

BIO-EFFICACY: The bio-efficacy of DRIMITI has been successfully tried and tested by our research wing

GENERAL INFORMATION: DRIMITI is an effective alternative to other synthetic contact pesticide. The formulation is primarily based on alkaloids, salts of unsaturated aliphatic carboxylic acids & natural oils. This product may be used to control targeted pests on crops which include vegetables, grapes and other crops. This product can be applied up to harvest.

Description	:	Viscous liquid
Odour	:	Specific
Storage stability	:	Minimum 24 months.

COMPOSITION

Alkaloids suspended in natural oils	:	25.00% (min.)
Salts of unsaturated aliphatic	:	25.00% (min.)
carboxylic acids Emulsifiers & media	:	50.00% (max.)

MODE OF ACTION: This is a contact formulation. The spray blocks the insects' breathing spores, interfering with respiration; suffocates and kills them. Residual activity is limited once the spray is dried.

DILUTION : 3-4 ml per litre of water.

RECOMMENDATION

Crop	Target Pests
Cotton	Mites
Tomato	Spider mite (t. evansi)
Chilli	Two-spotted spider mite
Rose	(t.urticae), common red spider mite
Egg plants	(t. cinnabarinus)
Grape, apple, pear & peach	Spider mite, Thrips

Corporate Office :

FF6, Maurya Arcade,
Opp SG Sola Highway, Sola
Science City Road, Sola,
Ahmedabad 380061-GUJ-INDIA
TEL FAX : +91 79 4005 3080

Web : http://www.drishtiagrotech.com
Email : group@drishtiagrotech.com
info@drishtibio.com
GST TIN No : 240 7360 5802
CST TIN No : 245 7360 5802

14. DRIKILLER

APPLICATION : An effective organic contact formula to control heliothis and mealy bugs.

BIO-EFFICACY : Satisfactory results were obtained during the bio-efficacy trials by the research wings of some of our patrons.

GENERAL INFORMATION & MODE OF ACTION

It is an effective alternative to synthetic contact pesticides. The formulation is primarily based on botanical extracts and modified natural oils. This product may be used to control targeted pests on crops which include vegetables, grapes and other crops. This product can be applied up to harvest. It does not harm the plants or affect the quality of the crop. Foliar spray, covering entire plants should be done at the beginning of pest infestation for effective control. For better result, sufficient volume spray solution should be used for uniform coverage. It is compatible with commonly used insecticides and fungicides except sulphur based ones. It does not yield any knock down effect. Upon spraying this contact formula, the insects lay weak and do not feed on further due to its antifeedant properties. The spray blocks insect's breathing spores, interfering with respiration; suffocates and kills them slowly. The insect dies due to suffocation and starvation. Residual activity is limited once the spray is dried. Further damage is controlled immediately after spray, as the insects are no more able to feed on the crops.

Description : Highly viscous liquid
Odour : Specific
Storage stability : Minimum 24 months.

COMPOSITION

Colloidal suspension of Natural bitters, natural soluble Polymers in sulphonated oils : 100.00%

DILUTION : 4-5 ml per litre of water.

RECOMMENDATION

It is suggested to spray in the evening so that drying of the sprayed solution is delayed.

Crop	Target Pests
Cotton	Bollworm, pink bollworm, cutworm
Grapes	Thrips, Aphids
Paddy	Stem borer
Beans	Aphid, Pod borer
Gram	Cut worm, Pod borer
Mung / green gram	Pod borer
Tomato	Fruit borer
Chillies	Aphids, Thrips

Corporate Office :

FF6, Maurya Arcade,
Opp SG Sola Highway, Sola
Science City Road, Sola,
Ahmedabad 380061-GUJ-INDIA
TEL FAX : +91 79 4005 3080

Web : http://www.drishtiagrotech.com
Email : group@drishtiagrotech.com
info@drishtibio.com
GST TIN No : 240 7360 5802
CST TIN No : 245 7360 5802

15. DRIBAIT

APPLICATION : As organic rat control bait.

GENERAL : DRIBAIT is a revolutionary product in pellet form for the control of rodent pest species. The product does not contain any toxic substance or poison and works based on a unique water absorption mechanism.

Description	:	Pellets.
Odour	:	Sweet
Shelf life	:	36 months.

COMPOSITION

Activated cellulose ($C_6H_{10}O_5$)	:	60.00%
Edible oil & flavouring agents	:	40.00%

MODE OF ACTION

DRIBAIT does not kill a rodent based on toxic effect of poison. DRIBAIT, on consumption by a rodent, physically disrupts its water absorption mechanism. As a result the rodent becomes lethargic, falls into coma and then dies. Death normally occurs in 4-6 days of regular bait intake. It contains only naturally derived food grade materials. No toxic chemicals are added in the formula. No distress or discomfort by the rodents was observed during the laboratory trials.

DOSAGE : 15-30 gms per rodent.

RECOMMENDATION

Prior to treatment, all likely food sources should be made unavailable. Place the bait at locations where there are signs of activity such as fresh droppings, or near burrows, runways, feeding places and points of entry around building perimeters. Keep the package content tightly sealed. Use 15-30 g of bait at 6 ft apart. For heavy infestation, reduce the distance to 4 ft apart. Protect bait stations from non-target animals and moisture. Inspect bait stations frequently and replace eaten bait. Can be applied by hand. Do not feed to non-target animals. To avoid possible chocking hazard, keep away from children and pets. Unused bait may be disposed with normal household refuse. A rat may carry certain diseases. It is advisable to wear protective gloves to dispose off any rat and or bait trays that have been visited by rates. Always wash hands after handling remains of bait and bait stations.

Corporate Office :

FF6, Maurya Arcade,
Opp SG Sola Highway, Sola
Science City Road, Sola,
Ahmedabad 380061-GUJ-INDIA
TEL FAX : +91 79 4005 3080

Web : http://www.drishtiagrotech.com
Email : group@drishtiagrotech.com
info@drishtibio.com
GST TIN No : 240 7360 5802
CST TIN No : 245 7360 5802

16. Shaarp Shakti:

For effective control of Thrips

Thrips, order Thysanoptera, are tiny, slender insects with fringed wings. They feed by puncturing their host and sucking out the cell contents. In the family Thripidae are plant feeders that scar leaf, flower, or fruit surfaces or distort plant parts. Most adult thrips are slender, minute and have long fringes on the margins of both pairs of their long, narrow wings. nymphs are similarly shaped with a long, narrow abdomen but lack wings. Thrips range in color from translucent white or yellowish to dark brown or blackish. In many species, thrips feed within buds and furled leaves or in other enclosed parts of the plant.

Thrips prefer to feed in rapidly growing tissue. Feeding by thrips typically causes tiny scars on leaves and fruit, called stippling, and can stunt growth. Damaged leaves may become papery and distorted. Infested terminals may discolor, become rolled, and drop leaves prematurely. Thrips cause silvery to brownish, scabby scarring on the avocado Thrips feeding on fruit trees can deform or scar developing fruit. Cotton thrips feeding severely distorts blueberry shoot tips and foliage, reducing yield.

Mites

Mites are common pests in many fruit trees, field crops, vegetables, and ornamental plants. The spider mites, also called web spinning mites, To the naked eye, spider mites look like tiny moving dots; Adult females, the largest forms, are less than 1/20 inch long. Spider mites live in colonies , mostly on the under-surfaces of leaves; a single colony may contain hundreds of individuals. The names "spider mite" and "web spinning mite" come from the silk webbing most species produce on infested leaves. The presence of webbing is an easy way to distinguish them from all other types of mites. Adults have eight legs and an oval body, with two red eyespots near the head end of the body. Females usually have a large, dark blotch on each side of the body and numerous bristles covering the legs and body. Immatures resemble adults, except the newly hatched larvae have only six legs. Eggs are spherical and translucent, like tiny droplets, becoming cream colored before hatching. Mites cause damage by sucking cell contents from leaves. A small number of mites is not usually reason for concern, but very high populations—levels high enough to show visible damage to leaves—can be damaging to plants. At first, the damage shows up as a stippling of light dots on the leaves; sometimes the leaves take on a bronze color. As feeding continues, the leaves turn yellow and drop off. Often leaves, twigs, and fruit are covered with large amounts of webbing. Damage is usually worse when compounded by water stress.

How the Shaarp Shakti works ?

Generally Shaarp Shakti directly act as on nervous system of the every soft bodied pests and manage the thrips, mites and other soft bodied pest populations. Shaarp Shakti is cent percent bio based formulations and eco friendly product. It is non toxic to human beings.

Dosage : 20 ml per pump as per our guideline to regular interval of 15 to 20 days.

Suggestions : For effective management of thrips and mites populations, it should be spray in the time of morning and evening.

Corporate Office :

FF6, Maurya Arcade,
Opp SG Sola Highway, Sola
Science City Road, Sola,
Ahmedabad 380061-GUJ-INDIA
TEL FAX : +91 79 4005 3080

Web : http://www.drishtiagrotech.com
Email : group@drishtiagrotech.com
info@drishtibio.com
GST TIN No : 240 7360 5802
CST TIN No : 245 7360 5802

17. Shaarp WF:

For effective control of White flies :
Whiteflies are tiny, sap-sucking insects that are frequently abundant in field crops, vegetables and ornamental plantings. They excrete sticky honeydew and cause yellowing or death of leaves. Whiteflies usually occur in groups on the undersides of leaves. They derive their name from the mealy, white wax covering the adult's wings and body. Adults are tiny insects with yellowish bodies and whitish wings. Whiteflies normally lay their tiny, oblong eggs on the undersides of leaves. The eggs hatch, and the young whiteflies gradually increase in size through four nymphal stages called instars. The crawlers move around for several hours, then settle and remain immobile. Later nymphal stages are oval and flattened like small scale insects. The legs and antennae are greatly reduced, and older nymphs do not move. The winged adult emerges from the last nymphal stage. All stages feed by sucking plant juices from leaves and excreting excess liquid as drops of honeydew as they feed. Whiteflies suck phloem sap. Large populations can cause leaves to turn yellow, appear dry, or fall off plants. Like aphids, whiteflies excrete honeydew, so leaves may be sticky or covered with black sooty mold. Feeding by the immature whitefly, *Bemisia tabaci,* can cause plant distortion, discoloration and may cause serious losses in cotton and some vegetable crops.

How the Shaarp WF works ?
Generally most of the pesticides or biopesticides have only limited effect on whiteflies. They kill only those whiteflies that come in direct contact with them. So, for these troublesome situations, Shaarp WF directly act as on nervous system of the whiteflies and also kill by means of contact poison. They also manage the thrips, mites and other soft bodied pest populations too. Shaarp WF is cent percent bio based formulations and eco friendly product. It is non toxic to human beings.
Dosage : 20 ml per pump as per our guideline to regular interval of 15 to 20 days.
Suggestions : For effective management of Whiteflies, thrips and mites populations, it should be spray in the time of morning and evening.

18. Shaarp Killer: (Larvicide Sp.)

Generally the eggs of Heliothis, Armiworm, Pink bollworm, Spodoptera etc pests the eggs are spherical and flattened. Newly hatched, first-instar larvae have several rows of dark tubercles along the back, each bearing one or two bristles. Larvae range from olive green to dark reddish brown in color and can be best distinguished from most other caterpillars by the tiny spines, visible under a hand lens, that cover most of the body surface. Cotton bollworm larvae damage bolls and squares. Larvae chew holes into the base of bolls and may hollow out locks. Moist frass usually accumulates around the base of the boll. Larvae may also chew shallow gouges in the boll surface, which can become infected with rot organisms. Squares injured by cotton bollworm usually have a round hole near the base. Fifth-instar larvae are the most destructive; they not only damage more fruit than do earlier instars, but they damage larger fruit that are harder for the plant to replace.

How the Shaarp Killer works?
Generally most of the pesticides or biopesticides have limited effect on cotton bollworm complex. Shaarp Killer directly act as on nervous system of the all bollworm complex of cotton and other field and vegetable crops and also kill by means of contact poison. Shaarp Killer is 100 % bio based formulation and eco friendly product. It is non toxic to human beings.
Dosage: 20 ml per pump as per our guideline to regular interval of 15 to 20 days.
Suggestions: For effective management of bollworm complex, it should be spray in the time of morning and evening.

Corporate Office :

FF6, Maurya Arcade,
Opp SG Sola Highway, Sola
Science City Road, Sola,
Ahmedabad 380061-GUJ-INDIA
TEL FAX : +91 79 4005 3080

Web : http://www.drishtiagrotech.com
Email : group@drishtiagrotech.com
info@drishtibio.com
GST TIN No : 240 7360 5802
CST TIN No : 245 7360 5802

19. Shaarp MB:

Generally mealy bug include in Hemiptera class and sub class is Homoptera. Mealy bug lay their eggs in soil upto the depth of 8-15 cm. they also lay their eggs on stem, cracks on the stem in bunch. So, the mealy bug management is very difficult. Their nymphs are very tiny and difficulty to see by nacked eyes. They generally spread naturally by wind, birds and animals and more commonly by people moving form infested area to non infested areas. Improper disposal of uprooted weeds which helped mealy bug to migrate from weeds to cotton fields By throwing uprooted weeds in canals/water channels helped the pest to migrate to other vegetation and cotton crop. They suck the cell sap from tender leaves and then they converted into adults. Female having white silky layer on the upper surface of the body. So, almost all the pesticides and bio pesticides are very difficult to cut this layer and manage this pest. The major damage by the adult and nymphs is crinkled or twisted leaves and shoots, bunched and unopened leaves, distorted or bushy shoots, white fluffy mass on buds, stem, fruit and roots, presence of honey dew, black sooty mold and ants, unopened flowers which often shrivel and die, small deformed fruits. Mealy bug attacks on Cotton, Egg plant, Okra, Tomato, Cabbage, Mango, Papaya, Pigeonpea, Guava, Cassava etc.

How the Shaarp MB works?

The Specialty bio based formulations which directly penetrate to the white coating layer and kill the adults of mealy bug. This type of formulations which is prepared only for the mealy bug management purpose only. Shaarp MB is cent percent bio based formulations. This product is safe for the natural enemies, human beings and other crops. Shaarp MB definitely increases the yield upto the 25 per cent in cotton crop.

Dosage: 40 ml per pump as per our guideline to regular interval of 15 to 20 days. For better and early results, mix the Shaarp MB with other pesticides upto 15 to 20 ml per 15 lits of water.

Suggestions: For effective management of Shaarp MB, it should be spray in the time of morning and evening.

Corporate Office :

FF6, Maurya Arcade,
Opp SG Sola Highway, Sola
Science City Road, Sola,
Ahmedabad 380061-GUJ-INDIA
TEL FAX : +91 79 4005 3080

Web : http://www.drishtiagrotech.com
Email : group@drishtiagrotech.com
info@drishtibio.com
GST TIN No : 240 7360 5802
CST TIN No : 245 7360 5802

PRICE LIST of BIOPRODUCTS:

PRODUCT	PV per Case	Packing unit (Gr/ ML)	DBP Rs./Unit	Case Size (Lit / Kg)	No of units/ Case	Case Rate (Rs.)	MRP per unit (Rs.)
DRIFUNGI Dose 2 ml/ lt	0.50	250	193	10	40	7,700	275
	0.50	500	368	10	20	7,350	525
	0.50	1000	700	10	10	7,000	1,000
DRIDM Dose 2 ml/ lt	0.50	250	210	10	40	8,400	300
	0.50	500	403	10	20	8,050	575
	0.50	1000	770	10	10	7,700	1,100
DRIANTIFEED Dose 2 to 3 ml/ lt	0.50	250	315	10	40	12,600	450
	0.50	500	613	10	20	12,250	875
	0.50	1000	1,190	10	10	11,900	1,700
DRIBACT Dose 3 ml/ lt	0.50	250	166	10	40	6,650	238
	0.50	500	333	10	20	6,650	475
	0.50	1000	665	10	10	6,650	950
DRIMITI Dose 3 ml/ lt	0.50	250	315	10	40	12,600	450
	0.50	500	613	10	20	12,250	875
	0.50	1000	1,190	10	10	11,900	1,700
DRIKILLER DOSE 2-3 ML/ LT	0.50	250	315	10	40	12,600	450
	0.50	500	613	10	20	12,250	875
	0.50	1000	1,190	10	10	11,900	1,700
DRIMAGIC Dose 2 to 3 ml/ pump	0.20	500	133	10	20	2,660	190
	0.20	1000	231	10	10	2,310	330
	0.20	5000	980	10	2	1,960	1,400
DRIZYME Dose 8 Kgs / Acre	0.20	1000	224	10	10	2,240	320
	0.20	5000	1,050	5	1	1,050	1,500
	0.20	8000	1,624	8	1	1,624	2,320
DRIMINO Dose 2 to 3 ml / lt	0.30	500	210	10	20	4,200	300
	0.30	1000	385	10	10	3,850	550
	0.30	5000	1,750	10	2	3,500	2,500
DRIMAZER Liquid dose 2 ml/ lt	0.20	500	158	10	20	3,150	225
	0.20	1000	280	10	10	2,800	400
	0.20	5000	1,225	10	2	2,450	1,750
DRIBT Liquid dose 2 ml/ lt	0.20	500	166	10	20	3,325	238
	0.20	1000	298	10	10	2,975	425
	0.20	5000	1,313	10	2	2,625	1,875
DRISTICKER (SPL as Milky White- dose 1 ml/ lt)	0.20	500	140	10	20	2,800	200
	0.20	1000	245	10	10	2,450	350
	0.20	5000	1,050	10	2	2,100	1,500
DRIBAIT (PALLETS & Liquid - dose 30 gr/ rodent at 6' dist)	0.40	250	140	10	40	5,600	200
	0.40	500	280	10	20	5,600	400
	0.40	1000	560	10	10	5,600	800

DRISHTI BIOTECH PVT LTD

DRISHTI BIOTECH

Corporate Office :

FF6, Maurya Arcade,
Opp SG Sola Highway, Sola
Science City Road, Sola,
Ahmedabad 380061-GUJ-INDIA
TEL FAX : +91 79 4005 3080

Web : http://www.drishtiagrotech.com
Email : group@drishtiagrotech.com
info@drishtibio.com
GST TIN No : 240 7360 5802
CST TIN No : 245 7360 5802

DBPL - PRICE LSIT OF BIOLOGICAL PRODUCTS FOR FY 2009- 2010

PRODUCT	PV per Case	Packing unit (Gr/ ML)	DBP	Case Size (Lit / Kg)	No of units/ Case	Case Rate (Rs.)	MRP per unit (Rs.)
BIOMAGIC (A revolutionary Chlorophyll Enhancer - dose 15 gram/ 15 Lt. Pump)	2.50	200	280	10	50	14,700	400
	2.50	500	700	10	20	14,700	1,000
AMAZER (A high end plant growth regulator dose 5 ml/ 15 lt Pump)	10.00	50	228	5	100	23,890	350
Shaarp MB (A Mealy Bug Special product dose 50 Ml/ 15 Lt Pump)	0.50	250	135	10	40	5,400	275
	0.50	500	255	10	20	5,100	500
	0.50	1000	480	10	10	4,800	950
Shaarp Shakti (A new generation Acaricide dose 20ml / 15lt pump)	1.00	100	150	5	50	7,500	265
	1.00	250	355	10	40	14,200	625
	1.00	500	690	10	20	13,800	1,200
Shaarp Killer (A Spl Larvaecide dose 20 ml / 15 lt pump)	1.00	100	195	5	50	9,500	280
	2.00	250	465	10	40	18,600	685
	2.00	500	910	10	20	18,200	1,350
Shaarp Arrow (Controls all thrips - dose of 20 ml/ 15 lt pump)	2.00	100	220	5	50	11,000	280
	2.00	250	520	10	40	20,800	685
	2.00	500	1,020	10	20	20,400	1,400
Shaarp WF (Special to control Whitefly - Dose 20 ml/ 15 Lt Pump)	1.00	100	205	5	50	10,250	300
	1.00	250	480	10	40	19,200	735
	1.00	500	940	10	20	18,800	1,450
Royal SC (wide spectrum biopesticide pest dose 35 ml/ 15 lt pump)	0.50	500	110	10	20	2,200	190
	0.50	1000	205	10	10	2,050	360
Shooter SC (Biopesticide dose 20ml / 15Lt pump)	0.50	500	170	10	20	3,400	415
	o.50	1000	325	10	10	3,250	800
Scorpion Kiing (Sea weed based PGR dose 35 ml/ 15Lt Pump)	1.50	500	150	10	20	3,000	230
	1.50	1000	280	10	10	2,800	450
SC Hume Tech 90 % (Dose 5 gr/ 15 Lt Pump)	1.50	50	65	3	60	3,900	190
	1.50	100	110	3	30	3,300	380
SC Hume 12 % (Humic acid dose 30 ml/ 15 Lt Pump)	1.50	500	115	10	20	2,300	310
	1.50	1000	215	10	10	2,150	600
SC Zyme (seaweed + Humic dose 30 ml/ 15 Lt Pump)	2.00	500	185	10	20	3,700	320
	2.00	1000	360	10	10	3,600	620

Corporate Office :

FF6, Maurya Arcade,
Opp SG Sola Highway, Sola
Science City Road, Sola,
Ahmedabad 380061-GUJ-INDIA
TEL FAX : +91 79 4005 3080

Web : http://www.drishtiagrotech.com
Email : group@drishtiagrotech.com
info@drishtibio.com
GST TIN No : 240 7360 5802
CST TIN No : 245 7360 5802

Price List of Vegetable Seeds for FY 2009 -10

Crop	Variety		Attributes	Paking size	Net Billing Price/ kg	MRP
Bitter Gourd	Harithwa		18-20 cm long fruit with dense prickles.	25 gm	2800	82
Hot Pepper	Bhamini		14-15 cm long thick fruit with light green color. Plant height 80-90 cm	10 gm	24000	300
Hot Pepper	SPS 818 (Umiya)		13-14 cm long smooth skinned fruit with dark green color. Dark red color after maturity.	10 gm	24000	320
Hot Pepper	SPS 87		10-12 cm long dark green colored fruit with wrinkles. Deep red color after maturity.	10 gm	30000	350
Hot Pepper	SPS 819		10-12 cm long green colored fruit with smooth skin. Deep red color after maturity.	10 gm	30000	350
Hot Pepper	SPS 817(Chamki)		12-13 cm long green colored fruit with smooth skin.	10 gm	30000	350
Okra	Tender		Green colored fruits with 14-16 cm length. Very good resistance to YVMV. Harvesting starts in 45-48 days after germination.	250 gm	700	400
Marigold	Haridra		70-80 cm tall plant. Flower color deep Yellow. Avg flower weight 10-12 g. Avg no of flowers per plant is 120-150	1000 seed	1350	1600
Marigold	Sunder		70-80 cm tall plant. Flower color Yellow. Avg flower weight 10-12 g. Avg no of flowers per plant is 120-150	1000 seed	1350	1600
Marigold	Sona		80-100 cm tall plant. Flower color deep orange. Avg flower weight 10-12 g. Avg no of flowers per plant is 120-150	1000 seed	1100	1250
Marigold	Sienna		70-80 cm tall plant. Flower color deep Orange. Avg flower weight 12-14gm.vg no of flowers per plant is 120-140	1000 seed	1350	1600
Tomato	Akshaya		Very good tolerence to TLCV. Plant height 80 -90cm. Oval shaped fruit with 80-90g weight. Good for cultivation in summer	10 gm	65000	810
Tomato	Harsha		Good tolerence to TLCV. Plant height 80-90cm. Oval round fruit with 80-90g weight. Good for cultivation in late summer and rainy season	10 gm	32000	450
Tomato	Bhoomika		Plant height 70-80cm.Round fruit with 70-80g weight. Sour in taste. Good for cultivation in rainy season	10 gm	28000	410
Tomato	Sampada		Plant height 70-80cm. Round fruit with 80-90g weight. Good for cultivation in rainy season	10 gm	22000	380
Tomato	Malhar		Plant height 70-80cm. Square Round fruit with 80-90g weight. Good for cultivation in rainy season	10 gm	24000	320
Tomato	Chandrika		Plant height 80-90cm. Very good tolerence to TLCV. Square Round fruit with 80-90g weight. Good for cultivation in rainy season	10 gm	28000	380
Ridge Gourd	Shiva		Vigours plant, Med long, slender, green fruits, high yield	25 gm	6000	240

DRISHTI BIOTECH PVT LTD

DRISHTI BIOTECH

Corporate Office :

FF6, Maurya Arcade,
Opp SG Sola Highway, Sola
Science City Road, Sola,
Ahmedabad 380061-GUJ-INDIA
TEL FAX : +91 79 4005 3080

Web : http://www.drishtiagrotech.com
Email : group@drishtiagrotech.com
info@drishtibio.com
GST TIN No : 240 7360 5802
CST TIN No : 245 7360 5802

DISCOUNT SCHEMES TO AUTHORISED DEALERS FOR BUSINESS PROMOTION (Applicable to Bio Products)

1

CASH DISCOUNT	DISCOUNT
1.1 Advance payment with written Order by Draft/ Scheduled Bank Chq	15%
1.2 Entire Bill Payment within 01 to 15 days on Bill Date by DD/SBC	12%
1.3 Entire Bill Payment within 16 to 30 days on Bill date by DD/ SBC	9%
1.4 Entire Bill Payment within 31 to 45 days on Bill date by DD/ SBC	6%
1.5 Entire Bill Payment within 46 to 60 days on Bill date by DD/ SBC	3%

**** DD= Demand Draft, SBC =Multicity Cheque from Scheduled Bank **** Payment will be treated as cash only on realisation of payment in our bank *** Cash Discount is applicable to Minimum 1 case of any single product in single order*

2

QUANTITY DISCOUNT	DISCOUNT
2.1 For Purchase of 5 to 10 Case of any Single Molecule.	1 %
2.2 For Purchase of 11 to 20 Case of any Single Molecule.	3 %
2.3 For Purchase of 21 to 35 Case of any Single Molecule.	5 %
2.4 For Purchase of 36 to 60 Case of any Single Molecule.	7 %
2.5 For Purchase above 61 Case of any Single Molecule.	10 %

*** QDD will be given at the end of Season on Net Quantity Achieved on realisation of all dues from party as C/N or equitable free goods offer on prevailing Market Price. Value of Goods will be calculated as DBP without VAT*

3

TURN OVER DISCOUNT	DISCOUNT
3.1 For Turnover within Rs. 1.5 lakhs to Rs. 2.99 Lakhs in above products.	1%
3.2 For Turnover within Rs. 3 lakhs to Rs. 5.99 Lakhs in above products.	2%
3.2 For Turnover within Rs. 6 lakhs to Rs. 9.99 Lakhs in above products.	3%
3.3 For Turnover within Rs. 10 Lakh to Rs. 14.99 Lakhs in above products.	4%
3.4 For Turnover above Rs. 15 lakhs with Above products.	5%

***** TOD will be given at the year end on net sales on Net Realised price/ value to the company after settlement of all dues from the party in the way of CN or net equitable Free goods Offer at prevailing market rate (its only applicable for Biomagic & Amazer Sales Jointly) QDD will be allowed on net turnover. The NTO value would be arrived after deducting normal trade discount, CD, QDD etc... This Scheme is applicable from 1st July 2009 to 30th October 2009 only*

Terms & Conditions :

1. All the above prices are on 60 days credit limit against PDC to authorised dealers only as per their internal credit limit set by company.
2. The company reserves the rights to change the price without any prior information.
3. The goods will be supplied on **Freight to Pay** basis to the nearest road booking stations and the freight amount would be deducted from Invoice.
4. Party will earn **points (Point Value = PV)** as per price list. The value per PV will be announced at the end of the season, subject to overall achievement of business by the company and will be communicated to party. An attractive scheme based on PV achieved has been designed for the promotion.
5. Any claims regarding shortage/ leakage due to transporter negligence shall be accepted only on submission of valid proof for the same from the transporter within 10 days of receipt of material.
6. Please note that no fresh supplies will be made till overdue bills are cleared.
7. Goods once sold will not be taken back or exchanged and on no account we shall accept any responsibility for goods returned to us without our written consent.
8. Please inform us regarding complaint if any, as soon as material is received by you.
9. Company will accept account Payee Drafts/ Cheques from scheduled banks only. Payment to be made in favour of " **DRISHTI BIOTECH PVT. LTD.**" payable at **AHMEDABAD** only.
10. Interest @18% PA will be charged from due date on remaining delayed payments and no further supplies are made till overdue payments are cleared.
11. This price list supersedes our all previous price list/ Quotation.
12. All jurisdictions to Ahmedabad Metropolis

Sincerely Yours,

For, Drishti Biotech Pvt Ltd.

Rakesh Jha
Chief Executive Officer

Corporate Office :

FF6, Maurya Arcade,
Opp SG Sola Highway, Sola
Science City Road, Sola,
Ahmedabad 380061-GUJ-INDIA
TEL FAX : +91 79 4005 3080

Web : http://www.drishtiagrotech.com
Email : group@drishtiagrotech.com
info@drishtibio.com
GST TIN No : 240 7360 5802
CST TIN No : 245 7360 5802

HYBRID HOT PEPPER

Tips for Growing Hot pepper

1) Soil: well drained black to medium clay loamy soils is suitable.
2) Sowing Time: As per original practices and timings.
3) Optimum Temp for Germination: 25-30 degree C.
4) Transplanting : 30-35 days after sowing.
5) Spacing: Row to Row: 75-90cm, plant to plant:45-60 cm.
6) Seed Rate: 80-100 g/ acre.

Preparation of main field

- Deep ploughing and harrowing. Add well dcomposed FYM 7-8 tones per acre followed by harrowing to mix in the soil thoroughly. Open the ridges and furrows at required spacing. Irrigate the field one day before transporting to mix in the soil thoroughly. Open the ridges and furrows at required spacing. Irrigate the field one day before transplanting make a hole at required spacing to plant seedling. Transplanting should be done late afternoon, complete transplanting light irrigation should be given for better and quick establishment.

- **Chemical Fertilizer: Fertilizer requirement varies with soil fertility.**
- First dose 6-8 days after transplanting: 50:100:75 NPK kg/acre
- Second dose 20-25 days after first application. 25:50:75 NPK Kg/ acre
- Third dose 20-25 days after second application. 25:0:0 NPK Kg/acre
- At the time of Flowering: Sulphur (Bensulf) 10 kg/acre
- Spray urea & Soluble K (1% solution each) at the time of flowering (To increase fruit set)
- Spray Urea & soluble K (1% solution each) at 15 days interval during harvesting time(to increase no.of pickings.)
- After 15 days or first picking add NP & K as and when required as recommended below.
- Plant protection: follow the plant protection measures as and when required as recommended below.

Disease	Chemical name	Trade Name	Doses per litre of water
Nursery damping –off Thrips /Aphids	Metalaxyl+ Mancozeb 72% wp carbendazim 50%wp Monocrotophos 36% SL imidachloprid 17.8 SL	Ridomyl, Master Bavistin Nuvacron Confidor	2.0g Drench 1.0 g Drench 1.0 g 0.3 ml
Main field powered powdery Mildew	Dinocap 48% EC	Karathane suftaf, Thivit Bayleton	1.0ml 2.0 g 0.3 g
Cercospora Leaf spot	Carvendazim 50% WP Mancozeb 75% wp	Bavistin Dithane M-45, IM-45 Manzate	1.5g 2.0g
Fusarium wilt	Fosetyl Al 80% wp carbandazim 50% Wp	Aliettle+ Bavistin	1.0g+ 1.0g- Drench
Fruit wilt	Profenofos + cypermethrin 4% EC quinolphos 25% EC	Polytrin c44 Ekalux	1.0ml , 2.0ml
Thrips/Aphids/Whiteflies	Monocrotophos 36% SL + Azardactin imidachloprid 17.8, Quinolphos 25% EC, Tirazophos 40% EC + Deltamethrin 2.8% EC	Nuvacorn, Monocil, Bilphos, + Econeem confidor Ekalux	1.5ml , +3.0 ml, 0.3ml, 2.0 ml
Mlites	Dicofol 18.5% EC Ethilon 50% EC sulphur 85% wp Abamectin 1.9% EC	Sparck, kelthane Tafetholin, mit-505, Sultaf, Thivoit vertimec	1.0ml,2.0ml,1.0ml, 2.0g, 0.5ml

Corporate Office :

FF6, Maurya Arcade,
Opp SG Sola Highway, Sola
Science City Road, Sola,
Ahmedabad 380061-GUJ-INDIA
TEL FAX : +91 79 4005 3080

Web : http://www.drishtiagrotech.com
Email : group@drishtiagrotech.com
info@drishtibio.com
GST TIN No : 240 7360 5802
CST TIN No : 245 7360 5802

HYBRID CAULIFLOWER

Tips for growing Cauliflower

1) soil: well drained medium loam and/ or sandy loam soils are suitable.
2) Optimum Temp for germination: 25-30C.
3) Transplanting: 25-30 days after sowing.
4) Spacing: Row to row : 60cm, plant to plant:45cm
5) Seed Rate:100-120 g/acre.

Preparation of main fied

Deep ploughing and harrowing. Add well decomposed FYM 7-8 tones per acre followed by harrowing to mix in the soil thoroughly. Open the ridges and furrows at required required spacing. Apply basal dose of chemical fertilizer before transplanting. Irrigate the field one day before transplanting, make a hole at required spacing to plant seedling. Transplanting should be one late afternoon, after transplanting light irrigation should be given for better and quick establishment.

Chemical Fertilizer: fertilizer requirement depends on the fertility of soil.

- First dose 6-8 days after transplanting: 50:50:60: NPK kg/acre
- Second dose 20-25 days after first application:25:50:60 NPK kg/acre
- Third dose 20-25 days after second application: 25:0:0 NPK kg/acre
- At the time of flowering: Sulphur (Bensulf) 10kg/acre.
- Spray calcium nitrate (1% solution) at the time of flowering (To increase fruit set).
- Spray urea & soluble k (1% solution each) at 15 days interval during harvesting time (To increase no. of . packings)

Plant protection: follow the plant protection measures as and when required as recommended below.

Disease/ insect pest	Chemical name	Trade name	Doses per litre of water
Damping off	Carbendazim 50% wp Metalaxy +Mancozeb 72% wp	Bavistin, Ridomli MZ	1.0g- Drenching , 2.0g- Drenching
Main field- Black Rot	Grow resistant varieties		
Soft Rot	Spray kasugamycin streptocyclin	Kasu-b, streptocyclin	0.5ml 1.0g per 10 liters.
Alternaria Leaf sports	Chlorothalanil 50% WP	Kavach	2.0g
Downy Mildew	Metalaxy+Mancozeb, fosetyl AL 80 wp	Ridomil-MMZ Aliette	2.0g, 1.0g
Diamond Back Moth	Flufenoxuron 10% DC indoxacarb 14.5% sc, Bacillus Thurigenesis	Cascade, Avaunt, deflin , dipel, Hault	0.5ml, 1.0g per 10 liters
Aphids	Monocrotophos 36% Ec Azadirachtin	Nuvacron Econeem, Neemark	1.5ml, 2.5ml
Head Borders/ Caterpillars	Qunolphos 25% EC + cypermethrin 10% EC	Ekalux Jawa, simper	2.0ml,1.0ml

Blanching: to avoid discoloring and yellowing, cover the curd by tying the leaves around on the curd.
Special note: The varietal characters & information given here is based on observations recorded at our trail station. Change in place, session, cultural practice may result in variation.

Corporate Office :

FF6, Maurya Arcade,
Opp SG Sola Highway, Sola
Science City Road, Sola,
Ahmedabad 380061-GUJ-INDIA
TEL FAX : +91 79 4005 3080

Web : http://www.drishtiagrotech.com
Email : group@drishtiagrotech.com
info@drishtibio.com
GST TIN No : 240 7360 5802
CST TIN No : 245 7360 5802

HYBRID TOMATO:

Tips for Growing Tomato

1 soil: A well drained loamy soil is ideal.
2: Sowing Time: As per regional practices and timings.
3: Optimum Temp for germination: 25-30 degree C.
4: Transplanting: 25-30 days after sowing.
5: spacing: Row to row: 90cm, plant to plant.:45-60 cm.
6: Seed Rate: 50-60 g/acre.

Preparation of main filed:

Deep ploughing and harrowing. Add well decomposed FYM 8-10 tones per acre. Make the ridges and furrows at the required spacing. Irrigate the filed and make the holes at the holes at the recommended spacing. Transplanting should be done during afternoon, after transplanting light irrigation should be given for quick and better establishment.

Chemical Fertilizer: Fertilizer requirement depends on the soil fertility.

- First dose 6-8 days after transplanting.: 50:100: NPK kg/acre
- Second dose 20-25 days after second application: 25:50:50: NPK kg/acre
- Third dose 20-25 days after second application:25:0:0 NPK kg/acre.
- At the time of flowering: sulphur (Bensulf) 10kg/acre.
- At the time of fruit setting: Boracol (BSF-12) 50kg/acre.
- Spray calcium nitrate (1% solution) at the time flowering (To increase fruit set).
- Spray Urea & Soluble K(1% solution each) at 15 days interval during harvesting time.(To increase no. of pickings).

Plant protection: Follow the plant protection measures as and when required as recommended below.

Disease/ insect plant	Chemical name	Trade name	Doses per litre
Nursery Aphids/white fly	Monocrotophos 36% SL imidochloprid 200 SL	Nuvacorn, Confidor	1.0ml, 0. 3ml
Damping-off	Carbendazim 50% wp, Metalaxyl + Mancozeb-72 wp	Bavistin, Ridomil-MZ, Master	1.0g-Drenching, 2.0g – Drenching
Main field Fusarium wilt/ Vertiillium wilt	Copper –oxychloride 50% wp fosetyl AL 80% wp+ carbendizam 50% wp	Biltox, Blue copper , Alitette+ Basvistin	3.0g, 1.0g, +1.0g-Directing
Early Blight	Mancozeb 75% wp, Fosetyl Al 80% wp+ Carbendazim 50% wp	Dithane- M-45,IM-45, Manzate, Karvach, Rovral	2.0g, 1.0g,2.0g
Late Blight	Metalaxyl + Mancozeb-72 wp	Ridomil-MZ, master	1.0ml.
Powdery Mildew	Dinocap 48% EC, Triademefon 25% wp	Karathene , Bayleton	0.3g, 1.5ml
Aphids / Thrips	Monocrotophos 36% EC, imdochloprid 200SL	Nuvacorn, Confidor	0.3ml, 1.0ml
Whiteflies/Leaf miner	Traizophos 40% EC, Deltamethrin 2.8% EC, Profenofos+ cypermethrin 4%EC, Quinolphos 25% EC	Spark, polythrin c44, Ekaulx	1.0ml. 2.0ml, 1.0ml
Miltes	Ethlon 50% EC, Sulphur 80% wp	Tarethion , Mit-505, sulfex, Trivit	2.0g. ,1.0ml
Fruit Borers	Prefenophos+ Cypermethrin 4EC, Quinolophos 25% Ec + Azaradactin	Polytrin C44, Ekaluc + Econeem	2.0ml, + 3.0ml

Special note: The varietal & information given here is based on observations recorded at our trial station. Change in place, season, cultural practice may result in variation

Corporate Office :

FF6, Maurya Arcade,
Opp SG Sola Highway, Sola
Science City Road, Sola,
Ahmedabad 380061-GUJ-INDIA
TEL FAX : +91 79 4005 3080

Web : http://www.drishtiagrotech.com
Email : group@drishtiagrotech.com
info@drishtibio.com
GST TIN No : 240 7360 5802
CST TIN No : 245 7360 5802

HYBRID EGGPLANT

Tips for growing Eggplant

1. Soil: A well drained loamy soil is ideal.
2. Sowing time: AS per regional practices and timings
3. Optimum Temp. for germination: 25-30C.
4. Transplanting: 30-32 days after sowing.
5. Spacing: Row to Row: 75-90 cm, plant to plant: 60cm.
6. Seed Rate: 80g/acre.

Preparation of main field

- Deep ploughing and harrowing. Add well decomposed FYM 8-10 tones per acre. Make the ridges and furrows at the required spacing. Irrigate the field and make the holes at the recommended spacing. Transplanting should be done during afternoon, after transplanting light irrigation should be given for quick and better establishment.
- Chemical Fertilizer: Fertilizer requirement depends on the soil fertility.
- First dose 6-8 days after transplanting : 50:50:60 NPK kg/ acre
- Second dose 20-25 days after first application. :25:50:60 NPK kg/acre.
- Third dose 20-25 days after second application: 25:00:00 NPK kg/acre.

Plant protection: follow the plant protection measures as and when required as recommended below.

Disease/ insect pest	Chemical Name	Trade Name	Doses per litre of water
Nursery Damping off	Carbendazim 50% wp, Metalaxyl + Manozeb 72% wp	Bavistin , Ridomyl	1.0g- Drenching, 2.0g- Drenching
Fusarium wilt/ Verticillum wilt	Copper-Oxychloride 50% Wp fosetyl AL 80% WP, Carbendizim 50%wp	Biltox, Blue copper, Aliette + , Bavistin	3.0g, 1.0g, +1.0g. Drenching
Phomopsis fruit rot	Chlorothlonil 75% wp, Manozeb 75% wp	Kavach, Dithane M-45, IM-45	2.0g, 2.0g
Mites	Ethlon 50% EC, sulphur 80% Wp, Abamectin 1.9% Ec	Tafethion, Mit-505, Thiovit, sulfex Vertimec	1.0ml, 2.0g, 0.5ml
Fruit and shoot borer	Quinalphos 25% EC. Profenophos + Cypermethrin 4% EC	Ekalux, polytrin C44,	2.0ml, 1.0ml
Jassids/ Aphids	Imidachloprid 200SL Thimethoxam 25% wg	Confidor, Actara	0.3ml, 0.5ml
White files	Triazophos 40% EC Deltamethrin 2.8% EC, Imidacloprid 200sl	Spark, Confidor	1.0 ml, 0.3 ml

Special note: The varietal characters & information given here is based on observations recorded at our trail station. Change in palace, season, cultural practice may result in variation.

Corporate Office :

FF6, Maurya Arcade,
Opp SG Sola Highway, Sola
Science City Road, Sola,
Ahmedabad 380061-GUJ-INDIA
TEL FAX : +91 79 4005 3080

Web : http://www.drishtiagrotech.com
Email : group@drishtiagrotech.com
info@drishtibio.com
GST TIN No : 240 7360 5802
CST TIN No : 245 7360 5802

SWEET PEPPER

Tips for Growing sweet pepper

1) Soil: Well drained sandy loam soils are suitable.
2) Sowing Time: As per regional practices and timings.
3) Optimum Temp for germination: 25-30 C.
4) Transplanting: 35-40 days after sowing.
5) Spacing: Row to Row : 90cm, Plant to Plant:45cm.
6) Seed Rate : 100-120 g/acre.

Preparation of main field;

Deep ploughing and harrowing. Apply well decomposed FYM @7-8 tones per acre, followed by harrowing to mix in the soil thoroughly. Open the ridges and furrows at required spacing to plant seedling. Transplanting should be done late afternoon, after complete transplanting light irrigation should be given for better and quick establishment.

Chemical Fertilizer: Fertilizer requirement depends on the fertility of soil.

- First dose 6-8 days after transplanting.: 50:100:75 NPK kg/acre
- Second dose 20-25 days after first application. : 25:50:75 NPK kg/acre.
- Third dose 20-25 days after second application.: 25:0:0 NPK kg/acre.
- After 15 days of first picking add NP & K as and when required at the rate of 20:0:30 NPK kg/acre.
- At the time of flowering.: Sulphur (Bensulf) 10 kg/acre.
- At the time fruit setting.: Boracol (BSF-12) 50kg/acre.

Plant protection: follow the plant protection measures as and when required as recommended below.

Disease/pest	Chemical Name	Trade Name	Doses per litre of water
Nursery Damping- off	Metalaxyl+ Mancozeb 72% WP Carbendazim 50% WP	Ridomyl, Master , Bavistin	2.0 g,1.0g
Thirps/ Aphids	Monocrotophos 36% SL, Imidacloprid 17.8% SL	Nuvacron, Conflidor	1.0g , 0.3ml
Main Field Powdery Mildew	Dinocap 48% EC, Sulphur 85% WP, Trademefon 25%WP	Karathane, suftaf, Thiovit, Bayleton	1.0ml, 2.0g, 0.3g
Cercospora Leaf spot	Mancozeb 75% WP, Cabendizim 50% WP	Dithane M-45, IM-45 , Bavistin	2.5 g, 1.0g
Fruit rot/Anthrancnose	Mancozeb 75% WP, Chlorothainil 75% wp.	Dithane –M-45, IM-45 Kavach.	2.5 g, 2.0g
Thrips/ Aphids	Monocrotophos 36% SL + Azardactin	Nuvacorn, Monocil, Bilphos, Econeem	1.5ml, 3.0ml
White files	Imidachloprid 17.8% SL , Trazophos 40% EC + Deltamethrin 2.8% EC	Confldor, spark	0.3ml, 1.0ml
Mites	Dicofol 18.5% EC, Ethion 50% EC, Sulphur 85% WP	Kelthane, Tafethion, Mit-505 Sultaf, Thiovit	2.0ml,1.0ml, 2.0g
Fruit Borers	Quinophos 25% EC, Azaradactin, Profenophos + Cypermethrin 4% EC	Ekalux+ Econeem polytrin C44	2.0ml + 3.0ml+ 1.5 ml

Special Note: The varietal characters & information given here is based on observation recorded at our trial station. Change in place, cultural practice may result in variation.

Corporate Office :

FF6, Maurya Arcade,
Opp SG Sola Highway, Sola
Science City Road, Sola,
Ahmedabad 380061-GUJ-INDIA
TEL FAX : +91 79 4005 3080

Web : http://www.drishtiagrotech.com
Email : group@drishtiagrotech.com
info@drishtibio.com
GST TIN No : 240 7360 5802
CST TIN No : 245 7360 5802

Onion - D Selection

Cultivation Tips

Soil: well drained sandy loam is suitable.
Sowing time: May-June
Transplanting: 40-45 days after sowing.
Spacing : Row to Row: 15cm, Plant :10cm
Seed Rate: 2.0 kg/acre
Preparation of Main Field: Deep ploughing of main followed by a 1-2 harrowing. Add 7-8 tones per acre well decomposed FYM followed by harrowing to mix in the soil thoroughly. Apply basal dose. At the time of transplanting irrigate the field and transplant the seedlings.

Fertilizer requirement: Fertilizer requirement varies with soil fertility.

- Apply basal dose at the time of planting: 30:30:30 NPK Kg/acre.
- Apply top dressing 25 days after planting: 25:25:25 NPK kg/acre.
- Apply top dressing 50 days after planting: 15:0:0 NPK kg/acre.
- Apply Sulphur (Bensulf) at the rate of : 10kg/acre

Plant protection

Follow the plant protection measures as and when required as recommended below.

Disease/insect	Chemical Name	Trade Name	Dosage per liter of water
Nursery damping off	Carbendazim 50% wp Metalaxy+ Mancozeb 72% WP	Bavistin Ridomyl , Master	1.0g-Drenching, 2.0g Drenching
Main field Anthracnose	Carbendazim 50%wp, Mancozeb 75% wp	Balvistin Dithane M-45,IM-45	1.0g, 2.0g
Stemphyllium	Chlorothalanil 80% wp	Kavach	2.0g
Leaf blight	Mancozeb	DM-45,IM-45	2.0g.
Purple Blotch	Iprodion 50% wp	Rovral	1.0g.
Thirps	Imidochloprid 200SL Azadara, Monocrotophos 36% SL	Confidor, Econeem, Neemark, Nuvacron	0.3ml, 3.0ml, 1.5ml

Harvesting:

Stop irrigation two weeks before harvest. After harvest keep the bulb in the field for curing for 5-6 days along with tops. Cover the bulbs to avoid sun scalding. After proper drying remove roots and neck, don't cut the neck close the bulb.

Special Note: The varietal characters & information given here is based on observations recorded at our trail station. Change in place, season, practice may result in variation.

Corporate Office :

FF6, Maurya Arcade,
Opp SG Sola Highway, Sola
Science City Road, Sola,
Ahmedabad 380061-GUJ-INDIA
TEL FAX : +91 79 4005 3080

Web : http://www.drishtiagrotech.com
Email : group@drishtiagrotech.com
info@drishtibio.com
GST TIN No : 240 7360 5802
CST TIN No : 245 7360 5802

Hybrid Okra

Soil: well drained sandy loams and clay loam soils are ideal for the crop.
Season: As per the regional practices and timings.
Seed Rate: 2.0-2.5kg/acre
Spacing: Row to Row: 60 cm, plant to Plant:30 cm
Optimum temp. for germination:25-30C.

Main field preparation:

Deep ploughing and Harrowing. Apply 10 tones of well decomposed FYM followed by harrowing to mix in the soil. Form ridges and furrows. Apply basal dose of fertilizers in furrows and cover the fertilizer. Irrigate the field one day prior to sowing. Dibble one seed per hill, immediately give one light irrigation for quick and better establishment.

Chemical Fertilizers: Fertilizer application varies with soil fertility.

- Basal dose: 30:40:40 NPK kg/acre.
- First top dressing 20-25 days after sowing: 20:00:40 NPK kg/acre.
- Second top dressing 20-25 days after first top 25:00:00 NPK Kg/acre Dressing.

Plant protection: Follow the plant protection measures as when required as recommended below.

Disease/Insect pest	Chemical Name	Trade Name	Dose per liter of water
Powdery mildew	Sulphur 80% WP, Trademefon 80%EC	Sulfex, Thiovit, Traidemefon 80% Ec	2.0 g, 0.3g
Insect Pest Whitely	Trazophos 40% EC+Deltamethrin 2.8% EC	Spark	1.0ml
Jassids	Imidochloprid 200Sl+ Azaradactin Thiamethoxarm 25% Wg	Cofidor, Econeem, Neemark, Actara	0.3 ml, 3.0ml, 0.5ml
Fruit borer	Profenophos+ Cypermethrin 4%EC, Quinophos 25% EC	Polytrin C44, Ekalux,	1.50ml, 2.0ml
Miltes	Ethlon 50% EC Abamectin 1.9% EC	Mit-505, Dhanu mit, Vertimec	1.0ml,0.5ml

Special note:

The varietal characters & information given here is based on observations recorded at our trail station. Change in place, season, cultural practice may result in variation.

Corporate Office :

FF6, Maurya Arcade,
Opp SG Sola Highway, Sola
Science City Road, Sola,
Ahmedabad 380061-GUJ-INDIA
TEL FAX : +91 79 4005 3080

Web : http://www.drishtiagrotech.com
Email : group@drishtiagrotech.com
info@drishtibio.com
GST TIN No : 240 7360 5802
CST TIN No : 245 7360 5802

Hybrid Bottle Gourd

Soil: well drained sandy loams and alluvial soils are best suited for the crop.
Season: As per the regional practices and things.
Seed Rate: 500-600 gm/acre

Spacing:
Staking system: Row to Row:120 cm, plant to plant:45cm ,
Ground system: Row to Row: 180cm, plant to plant: 45cm.
Optimum temp.for germination:28-32 C

Main field preparation:
Deep ploughing and Harrowing. Add well decomposed FYM 10-12tons/ acre*. Open ridges & furrows at required spacing. Irrigate one day before sowing.

*Chemical Fertilizers**
Fertilizer application varies with soil fertility
Basal dose before sowing: 25:50:50kg/acre.
Top dressing 30 days after sowing: 25:0:50 NPK kg/acre.

Plant protection
Follow the plant protection measures as recommended below as and when required.

Disease/ Insect post	Chemical Name	Trade Name	Dose per liter of water
Powdery mildew	Dinocap 48% EC, Triadimefon 25% WP	Karathane, Bayleton	2.0ml,0.3g
Insect pest: Leafminer	Triazophos 40% EC or Deltamethrin 2.8% EC Profenophos+ Cypermethrin	Spark polytrin C44	1.0ml, 1.5ml
Red Pumpin beetle	Thiamethoxam 25% WG Carbaryl 50% WP	Actara Sevin 50% WP	0.5ml, 4.0G

Special Note: The varietal characters & information given here is based on observations at our trial station. Change in place, season, cultural practice may result in variation.