

24.118: Paradox and Infinity

Ryan Lacey <rlacey@mit.edu>

Collaborator(s): Evan Thomas

March 20, 2014

1. Define P_n to be the n^{th} prime. Eg. $P_0 = 2$, $P_1 = 3$, $P_2 = 5$, etc

For each P_n , if n is an even number, map it to $n/2$; and if n is an odd number, map it to $-((n+1)/2)$.

$$P_0 \rightarrow 0/2 = 0$$

$$P_1 \rightarrow -(1+1)/2 = -1$$

$$P_2 \rightarrow 2/2 = 1$$

$$P_3 \rightarrow -(3+1)/2 = -2$$

2. We build a matrix of pairs $\langle n, m \rangle$ with n specifying the row and m specifying the column of the pair's cell. The setup is similar to the figure below, but with a natural number pair instead of a ratio of natural numbers. We can traverse the matrix in the same increasing diagonal manner as depicted in the diagram. This strategy ensures that we have a counting strategy (appropriate for the natural numbers) and that we are guaranteed to encounter all pairs of the natural numbers.

0 1/1	2 1/2	3 1/3	9 1/4	10 1/5	...
1 2/1	4 2/2	8 2/3	11 2/4	2/5	...
5 3/1	7 3/2	12 3/3	3/4	3/5	...
6 4/1	13 4/2	4/3	4/4	4/5	...
14 5/1	5/2	5/3	5/4	5/5	...
...

Figure 1: Matrix of rational numbers

3. Define a function from the interval $[0,1)$ to the interval $[0,\infty)$ as the following

$$f(x) = \frac{1}{1-x} - 1$$

Let's observe the extremes of the input range

$$f(0) = \frac{1}{1-0} - 1 = 0$$

$$f(1) = \frac{1}{1-1} - 1 = \infty$$

\therefore we can map anything from zero to infinity from the interval of zero to one.

4. skipped

5. Yes, there is a one-to-one correspondence between the natural numbers and nodes of the tree. We can traverse the tree by row and within each row visit the nodes. Each node visited will be assigned a natural number. Eg. Row 0 has one node, which is assigned value 0. Row 1 has two nodes, which are assigned values 1 and 2, etc.

Proof for real numbers applies to tree path, just under a different representation. Proof taken from reals vs ration proof in the book.

No, there is not a one-to-one correspondence between the paths of the trees and the natural numbers. Assume that it is possible to assign a different natural number to each path. If we have such an assignment, we can use it to make a list of all paths. We just let the first member of our list be the path that has been paired with the smallest natural number, the second member of our list be the path that has been paired with the second smallest natural number, and so forth. This allows us to build a table like the following

0,	a_1	a_2	a_3	a_4	a_5	...
0,	b_1	b_2	b_3	b_4	b_5	...
0,	c_1	c_2	c_3	c_4	c_5	...
0,	d_1	d_2	d_3	d_4	d_5	...
0,	e_1	e_2	e_3	e_4	e_5	...
...

$$d = 0, a_1 b_2 c_3 d_4 e_5 \dots$$

Now consider the number $d = 0.a_1 b_2 c_3 d_4 e_5 \dots$. We know that value d must be different from the first member of our list because $f(a_1)$ is not equal to a_1 , and it is different from the second member of our list because $f(b_2)$ is not equal to b_2 , and so forth. But d is a path of 0s and 1s. So we have succeeded in identifying a path not on our list. That contradicts our assumption. So our proof by reductio is complete.