Adaptive (weighted) lasso in GCLMs

0 Setting (DDK §§2–3)

Data and model. $X_1, \dots, X_n \in \mathbb{R}^p$ are i.i.d. $N(0, \Sigma^*)$ with $M^*\Sigma^* + \Sigma^* M^{*\top} + C = 0$, where $M^* \in \mathrm{Stab}_p$ and $C \in PD_p$ is known. $\hat{\Sigma} = \frac{1}{n}\sum_i X_i X_i^\top$.

Design and Gram matrix. For symmetric Σ:

$$A(\Sigma) = (\Sigma \otimes I_p) + (I_p \otimes \Sigma)K^{(p,p)}, \qquad \Gamma(\Sigma) = A(\Sigma)^\top A(\Sigma), \qquad g(\Sigma) = -A(\Sigma)^\top \mathrm{vec}(C),$$

so that

$$L(M) := \tfrac{1}{2}\|M\Sigma + \Sigma M^\top + C\|_F^2 = \tfrac{1}{2}\,\mathrm{vec}(M)^\top \Gamma(\Sigma)\,\mathrm{vec}(M) - g(\Sigma)^\top \mathrm{vec}(M) + \tfrac{1}{2}\|C\|_F^2.$$

Shorthand. $\hat{\Gamma} = \Gamma(\hat{\Sigma})$, $\Gamma^* = \Gamma(\Sigma^*)$, $\hat{g}$, g^*, $\Delta_\Gamma = \hat{\Gamma} - \Gamma^*$, $\Delta_g = \hat{g} - g^*$. Indices are pairs (j,k); $m = \mathrm{vec}(M)$, $m^* = \mathrm{vec}(M^*)$.

Support. $S = \{(j,k) : M^*_{jk} \neq 0\}$, $d = |S|$; Since M^* solves the Lyapunov equation, $\nabla L(m^*) = 0 \iff \Gamma^* m^* = g^*$.

Constants (DDK).

$$c_{\Gamma^*} = \|\!|(\Gamma^*_{SS})^{-1}|\!\|_\infty, \qquad c_{M^*} = \|m^*\|_\infty,$$
$$P := \Gamma^*_{S^cS}(\Gamma^*_{SS})^{-1}, \qquad \rho := \|\!|P|\!\|_\infty, \qquad \beta_{\min} := \min_{(j,k)\in S} |M^*_{jk}|$$

1 Estimator, weights, weighted irrepresentability constant

Pilot and weights

- The random pilot $\hat{M}^0$ is the lasso (diagonals penalized), at the smallest λ of its grid.
- With $\hat{\mu} := \max_{(k,l)} |\hat{M}^0_{kl}|$, the maximum over all entries including the diagonal,

$$\hat{w}_{jk} = \frac{\hat{\mu}}{|\hat{M}^0_{jk}|} \quad \text{if } \hat{M}^0_{jk} \neq 0, \qquad \hat{w}_{jk} = \infty \ \text{ (entry excluded, stays 0)} \quad \text{if } \hat{M}^0_{jk} = 0.$$

- All finite weights are ≥ 1, with equality at the largest pilot entry.

Estimator (weighted direct Lyapunov lasso, all entries penalised):

$$\begin{aligned} \hat{M} &\in \operatorname*{arg\,min}_{M\in\mathbb{R}^{p\times p}} \ \tfrac{1}{2}\|M\hat{\Sigma} + \hat{\Sigma}M^\top + C\|_F^2 + \lambda \sum_{j,k} \hat{w}_{jk}|M_{jk}| \\ &= \operatorname*{arg\,min}_{m} \ \tfrac{1}{2} m^\top \hat{\Gamma} m - \hat{g}^\top m + \lambda \|\hat{W} m\|_1, \qquad \hat{W} = \mathrm{diag}(\hat{w}). \end{aligned} \tag{1.1}$$

An infinite weight means the constraint $m_j = 0$.

Scale invariance. Replacing $(\lambda, \hat{w})$ by $(\lambda/c, c\hat{w})$ leaves the objective unchanged. The products $\lambda\hat{w}_j$ matter.

Quantities. Define

$$\underline{w} := \min_{j \in S^c} \hat{w}_j, \qquad \bar{w}_S := \max_{k \in S} \hat{w}_k < \infty, \qquad r := \frac{\bar{w}_S}{\underline{w}} = \frac{\max_{j \in S^c} |\hat{M}^0_j|}{\min_{k \in S} |\hat{M}^0_k|},$$

- $\bar{w}_S < \infty$ requires the pilot to have no zero on S, a necessary condition for recovery. In particular, all infinite weights lie in S^c.

Weighted irrepresentability constant (Zhou, van de Geer & Bühlmann 2009, Def. 8.1):

$$\rho_w := \max_{j \in S^c} \frac{1}{\hat{w}_j} \sum_{k \in S} |P_{jk}|\, \hat{w}_k \quad \Big(= \| \hat{W}^{-1}_{S^c} P_{S^c \cdot} \hat{W}_{SS} \|_\infty \Big). \tag{1.2}$$

κ**:** Fix any $\kappa \geq 1 + \rho$. In DDK's setting, $\rho \leq 1 - \alpha$ and $\kappa = 2 - \alpha$.

2 W.1 (deterministic; analogue of DDK Theorem D.1)

Theorem W.1. *Let $M^* \in \mathrm{Stab}_p$ have support S, let $\hat{w}$ be weights with $\bar{w}_S < \infty$, and let $\kappa \geq 1 + \rho$. Assume:*

(W1) *Γ^*_{SS} is invertible;*

(W2) *(weighted irrepresentability) $\rho_w \leq 1 - \alpha$ for some $\alpha \in (0, 1]$;*

(W3) *$\| (\Delta_\Gamma)_{\cdot S} \|_\infty \leq \varepsilon_1$ and $\|\Delta_g\|_\infty \leq \varepsilon_2$, with*

$$\varepsilon_1 \leq \frac{\alpha}{3 c_{\Gamma^*}(\kappa r + \alpha)};$$

(W4) *the effective penalty off the support satisfies*

$$\lambda\, \underline{w} > \frac{3\kappa}{\alpha} \max\{c_{M^*}\varepsilon_1,\ \varepsilon_2\}.$$

Then:

(a) *The minimiser $\hat{M}$ of (1.1) is unique, its support is contained in the true support ($\hat{S} \subseteq S$), and*

$$\begin{aligned} \|\hat{M} - M^*\|_\infty \leq b_w &:= \frac{c_{\Gamma^*}}{1 - c_{\Gamma^*}\varepsilon_1}\big(c_{M^*}\varepsilon_1 + \varepsilon_2 + \lambda \bar{w}_S\big) \\ &< c_{\Gamma^*}\, \lambda\Big(\bar{w}_S + \frac{\alpha}{\kappa}\, \underline{w}\Big). \end{aligned} \tag{2.1}$$

(b) *If moreover $\beta_{\min} = \min_{(j,k) \in S} |M^*_{jk}| > b_w$, then $\hat{S} = S$ and $\operatorname{sign}(\hat{M}_{jk}) = \operatorname{sign}(M^*_{jk})$ for all $(j, k) \in S$.*

Reduction to DDK Theorem D.1. Take $\hat{w} = \mathbf{1}$ and $\kappa = 2 - \alpha$ with $\rho \leq 1 - \alpha$. Then:

- (W2) is DDK's (D.1), $\rho \leq 1 - \alpha$;
- (W3) is DDK's $\varepsilon_1 \leq \alpha/(6c_{\Gamma^*})$, since $r = 1$;
- (W4) is DDK's $\lambda > \frac{3(2-\alpha)}{\alpha} \max\{c_{M^*}\varepsilon_1, \varepsilon_2\}$;
- the right-hand bound in (2.1) is $c_{\Gamma^*}\lambda(1 + \frac{\alpha}{2-\alpha}) = \frac{2c_{\Gamma^*}\lambda}{2-\alpha}$, which is DDK's bound;
- DDK's beta-min $\beta_{\min} > 2c_{\Gamma^*}\lambda/(2 - \alpha)$ implies the one in (b).

3 Proof of W.1

KKT conditions Problem (1.1) is convex, since $\hat{\Gamma} \succeq 0$.

$$\begin{aligned} &\big(\hat{\Gamma}\hat{m} - \hat{g}\big)_j + \lambda\, \hat{w}_j \hat{z}_j = 0 \quad \text{(for every } j \text{ with } \hat{w}_j < \infty), \\ &\hat{z}_j = \operatorname{sign}(\hat{m}_j) \text{ if } \hat{m}_j \neq 0, \qquad \hat{z}_j \in [-1,1] \text{ if } \hat{m}_j = 0. \end{aligned} \tag{3.1}$$

Block form. Since $\Gamma^* m^* = g^*$, we have $\hat{\Gamma}\hat{m} - \hat{g} = \Gamma^*(\hat{m} - m^*) + \Delta_\Gamma \hat{m} - \Delta_g$. With $\Delta_M := \hat{m} - m^*$, the KKT conditions (3.1) read

$$\begin{bmatrix} \Gamma^*_{SS} & \Gamma^*_{SS^c} \\ \Gamma^*_{S^cS} & \Gamma^*_{S^cS^c} \end{bmatrix} \begin{bmatrix} (\Delta_M)_S \\ (\Delta_M)_{S^c} \end{bmatrix} + \begin{bmatrix} (\Delta_\Gamma)_{SS} & (\Delta_\Gamma)_{SS^c} \\ (\Delta_\Gamma)_{S^cS} & (\Delta_\Gamma)_{S^cS^c} \end{bmatrix} \begin{bmatrix} \hat{m}_S \\ \hat{m}_{S^c} \end{bmatrix} - \begin{bmatrix} (\Delta_g)_S \\ (\Delta_g)_{S^c} \end{bmatrix} + \lambda \begin{bmatrix} (\hat{w} \circ \hat{z})_S \\ (\hat{w} \circ \hat{z})_{S^c} \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix},$$

where the S^c-rows are taken only for j with $\hat{w}_j < \infty$.

▶ The PDW constructs a pair $(\tilde{m}, \tilde{z})$ satisfying (3.1) with $\tilde{m}_{S^c} = 0$.

Step 1 (restricted problem) Let

$$\tilde{m} = \underset{m:\ m_{S^c}=0}{\arg\min}\ \tfrac{1}{2} m^\top \hat{\Gamma} m - \hat{g}^\top m + \lambda \|\hat{W} m\|_1. \tag{3.2}$$

The restricted objective is strictly convex (as $\hat{\Gamma}$ positive semidefinite and invertible according to DDK) and $\tilde{m}$ is unique.

At $\tilde{m}$, where $\tilde{m}_{S^c} = 0$, the S-rows of 3.1 give

$$(\hat{\Gamma}\tilde{m} - \hat{g})_S + \lambda(\hat{w} \circ \tilde{z})_S = \hat{\Gamma}_{SS}\tilde{m}_S - \hat{g}_S + \lambda(\hat{w} \circ \tilde{z})_S = 0. \tag{3.3}$$

Error bound $\tilde{m}_S - m^*_S$***:***

Since $m^*_{S^c} = 0$ and $\Gamma^*_{SS} m^*_S - g^*_S = 0$, we have

$$\hat{\Gamma}_{SS} m^*_S - \hat{g}_S = \underbrace{\Gamma^*_{SS} m^*_S - g^*_S}_{=0} + (\Delta_\Gamma)_{SS} m^*_S - (\Delta_g)_S = (\Delta_\Gamma)_{SS} m^*_S - (\Delta_g)_S.$$

Subtracting this from (3.3) gives

$$\hat{\Gamma}_{SS} \underbrace{(\tilde{m}_S - m^*_S)}_{=\tilde{\Delta}_S} + \lambda(\hat{w} \circ \tilde{z})_S = -((\Delta_\Gamma)_{SS} m^*_S - (\Delta_g)_S)$$

As $\hat{\Gamma}_{SS}$ invertible (shown in DDK):

$$\tilde{\Delta}_S := \tilde{m}_S - m^*_S = -(\hat{\Gamma}_{SS})^{-1} \underbrace{\Big((\Delta_\Gamma)_{SS} m^*_S - (\Delta_g)_S + \lambda(\hat{w} \circ \tilde{z})_S\Big)}_{=:v}.$$

Then: $\|\tilde{\Delta}_S\|_\infty \leq |||(\hat{\Gamma}_{SS})^{-1}|||_\infty \|v\|_\infty$.

For v,

$$\begin{aligned} \|v\|_\infty &\leq \|(\Delta_\Gamma)_{SS} m^*_S\|_\infty + \|(\Delta_g)_S\|_\infty + \lambda \|(\hat{w} \circ \tilde{z})_S\|_\infty && \text{(triangle inequality)} \\ &\leq |||(\Delta_\Gamma)_{\cdot S}|||_\infty\, c_{M^*} + \|\Delta_g\|_\infty + \lambda \bar{w}_S && (\|m^*_S\|_\infty \leq c_{M^*},\ |\tilde{z}_k| \leq 1) \\ &\leq c_{M^*}\varepsilon_1 + \varepsilon_2 + \lambda \bar{w}_S && \text{(W3)}. \end{aligned}$$

For the inverse, write $\hat{\Gamma}_{SS} = \Gamma^*_{SS} + (\Delta_\Gamma)_{SS} = \Gamma^*_{SS}(I + X)$ with $X := (\Gamma^*_{SS})^{-1}(\Delta_\Gamma)_{SS}$. Then

$$\begin{aligned} |||X|||_\infty &\le |||(\Gamma^*_{SS})^{-1}|||_\infty \, |||(\Delta_\Gamma)_{SS}|||_\infty && \text{(submultiplicativity)} \\ &\le c_{\Gamma^*}\varepsilon_1 && \text{(W3)} \\ &\le \frac{\alpha}{3(\kappa r + \alpha)} \le \frac{1}{3} && \text{(W3)}. \end{aligned}$$

So the Neumann series $(I + X)^{-1} = \sum_{k\ge 0}(-X)^k$ converges, $\hat{\Gamma}_{SS}$ is invertible, and

$$\begin{aligned} |||(\hat{\Gamma}_{SS})^{-1}|||_\infty &= |||(I + X)^{-1}(\Gamma^*_{SS})^{-1}|||_\infty \\ &\le |||(I + X)^{-1}|||_\infty \, c_{\Gamma^*} && \text{(submultiplicativity)} \\ &\le \sum_{k\ge 0} |||X|||_\infty^k \, c_{\Gamma^*} = \frac{c_{\Gamma^*}}{1 - |||X|||_\infty} && \text{(Neumann series)} \\ &\le \frac{c_{\Gamma^*}}{1 - c_{\Gamma^*}\varepsilon_1} && (|||X|||_\infty \le c_{\Gamma^*}\varepsilon_1). \end{aligned} \tag{3.4}$$

Multiplying the two bounds gives

$$\|\tilde{\Delta}_S\|_\infty \le \frac{c_{\Gamma^*}}{1 - c_{\Gamma^*}\varepsilon_1}\big(c_{M^*}\varepsilon_1 + \varepsilon_2 + \lambda \bar{w}_S\big) =: b_w. \tag{3.5}$$

Converting the bound into λ:

By (W4), $c_{M^*}\varepsilon_1 + \varepsilon_2 \le 2\max\{c_{M^*}\varepsilon_1, \varepsilon_2\} < \frac{2\alpha}{3\kappa}\lambda\underline{w}$. By (W3), $\frac{1}{1-c_{\Gamma^*}\varepsilon_1} \le \frac{3(\kappa r+\alpha)}{3\kappa r+2\alpha}$. Hence

$$\begin{aligned} b_w &< c_{\Gamma^*} \cdot \frac{3(\kappa r + \alpha)}{3\kappa r + 2\alpha} \cdot \lambda\underline{w}\Big(r + \frac{2\alpha}{3\kappa}\Big) \\ &= c_{\Gamma^*}\lambda\underline{w}\,\frac{\kappa r + \alpha}{\kappa} = c_{\Gamma^*}\lambda\Big(\bar{w}_S + \frac{\alpha}{\kappa}\underline{w}\Big). \end{aligned} \tag{3.6}$$

▶ This is (D.4). With $\hat{w} \equiv 1$ and $\kappa = 2 - \alpha$ the right-hand side is DDK's $\frac{6c_{\Gamma^*}}{6-\alpha} \cdot \frac{6-\alpha}{3(2-\alpha)}\lambda = \frac{2c_{\Gamma^*}\lambda}{2-\alpha}$.

Step 2 ($\tilde{z}_S$: subgradient on S). Take $\tilde{z}_S$ from (3.3). By construction it is a subgradient of $\|\cdot\|_1$ at $\tilde{m}_S$.

Step 3 ($\tilde{z}_S^c$: dual vector on S^c). Evaluate the block form at $(\tilde{m}, \tilde{z})$. Since $\tilde{m}_{S^c} = m^*_{S^c} = 0$, we have $(\Delta_M)_{S^c} = 0$ and $(\Delta_M)_S = \tilde{\Delta}_S$:

$$\begin{bmatrix} \Gamma^*_{SS} & \Gamma^*_{SS^c} \\ \Gamma^*_{S^cS} & \Gamma^*_{S^cS^c} \end{bmatrix} \begin{bmatrix} \tilde{\Delta}_S \\ 0 \end{bmatrix} + \begin{bmatrix} (\Delta_\Gamma)_{SS} & (\Delta_\Gamma)_{SS^c} \\ (\Delta_\Gamma)_{S^cS} & (\Delta_\Gamma)_{S^cS^c} \end{bmatrix} \begin{bmatrix} \tilde{m}_S \\ 0 \end{bmatrix} - \begin{bmatrix} (\Delta_g)_S \\ (\Delta_g)_{S^c} \end{bmatrix} + \lambda \begin{bmatrix} (\hat{w} \circ \tilde{z})_S \\ (\hat{w} \circ \tilde{z})_{S^c} \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}.$$

The zeros remove the S^c-columns of Γ^* and Δ_Γ. Collect the noise in

$$u := (\Delta_\Gamma)_{\cdot S}\tilde{m}_S - \Delta_g, \qquad \text{i.e.} \quad u_S = (\Delta_\Gamma)_{SS}\tilde{m}_S - (\Delta_g)_S, \quad u_{S^c} = (\Delta_\Gamma)_{S^cS}\tilde{m}_S - (\Delta_g)_{S^c}.$$

The two block rows read

$$\begin{aligned} \Gamma^*_{SS}\tilde{\Delta}_S + u_S + \lambda(\hat{w} \circ \tilde{z})_S = 0 \qquad & (S\text{-rows}), \\ \Gamma^*_{S^cS}\tilde{\Delta}_S + u_{S^c} + \lambda(\hat{w} \circ \tilde{z})_{S^c} = 0 \qquad & (S^c\text{-rows with } \hat{w}_j < \infty). \end{aligned}$$

Solving the S-rows gives $\tilde{\Delta}_S = -(\Gamma^*_{SS})^{-1}\big(u_S + \lambda(\hat{w} \circ \tilde{z})_S\big)$.

Define $\tilde{z}_j$, for $j \in S^c$ with $\hat{w}_j < \infty$, by the S^c-rows, solved for the $\tilde{z}$-term. Then insert $\tilde{\Delta}_S$ and write $P = \Gamma^*_{S^cS}(\Gamma^*_{SS})^{-1}$:

$$\begin{aligned}
\lambda(\hat{w} \circ \tilde{z})_{S^c} &= -\Gamma^*_{S^cS}\tilde{\Delta}_S - u_{S^c} && (S^c\text{-rows}) \\
&= \Gamma^*_{S^cS}(\Gamma^*_{SS})^{-1}(u_S + \lambda(\hat{w} \circ \tilde{z})_S) - u_{S^c} && (\text{insert } \tilde{\Delta}_S) \\
&= P(u_S + \lambda(\hat{w} \circ \tilde{z})_S) - u_{S^c} && (\text{definition of } P) \\
&= Pu_S - u_{S^c} + \lambda P(\hat{w} \circ \tilde{z})_S && (\text{expand}),
\end{aligned}$$

and dividing row j by $\lambda\hat{w}_j$,

$$\tilde{z}_j = \underbrace{\frac{1}{\lambda\hat{w}_j}\left[Pu_S - u_{S^c}\right]_j}_{\text{noise}} + \underbrace{\frac{1}{\hat{w}_j}\left[P(\hat{w} \circ \tilde{z})_S\right]_j}_{\text{deterministic}}. \tag{3.7}$$

▶ This is (D.5) with $\operatorname{sign}(\operatorname{vec}(\tilde{M})_S)$ replaced by $(\hat{w} \circ \tilde{z})_S$ and row j divided by $\hat{w}_j$.

Deterministic term. Since $|\tilde{z}_k| \le 1$,

$$\left|\frac{1}{\hat{w}_j}\left[P(\hat{w} \circ \tilde{z})_S\right]_j\right| \le \frac{1}{\hat{w}_j}\sum_{k \in S}|P_{jk}|\hat{w}_k \le \rho_w \le 1 - \alpha \qquad \text{by (W2).} \tag{3.8}$$

▶ DDK bound this term by $\rho \le 1 - \alpha$. This is the only place where the weighted condition enters.

Noise term.

$$\begin{aligned}
|\text{noise}_j| &\le \frac{1}{\lambda\hat{w}_j}(\rho\|u_S\|_\infty + \|u_{S^c}\|_\infty) && (|||P|||_\infty = \rho) \\
&\le \frac{\kappa}{\lambda\underline{w}}\|u\|_\infty && (1 + \rho \le \kappa,\ \hat{w}_j \ge \underline{w}) \\
&\le \frac{\kappa}{\lambda\underline{w}}(\varepsilon_1\|\tilde{m}_S\|_\infty + \varepsilon_2) && \text{Def. of } u, \text{ (W3)} \\
&\le \underbrace{\frac{\kappa\, c_{M^*}\varepsilon_1}{\lambda\underline{w}}}_{G_1} + \underbrace{\frac{\kappa\,\varepsilon_1\|\tilde{\Delta}_S\|_\infty}{\lambda\underline{w}}}_{G_2} + \underbrace{\frac{\kappa\,\varepsilon_2}{\lambda\underline{w}}}_{G_3} && (\|\tilde{m}_S\|_\infty \le c_{M^*} + \|\tilde{\Delta}_S\|_\infty).
\end{aligned} \tag{3.9}$$

▶ These are DDK's G_1, G_2, G_3 (pp. 532–533), with $(2-\alpha)/\lambda$ replaced by $\kappa/(\lambda\underline{w})$. Each is below $\alpha/3$:

- $G_1, G_3 < \alpha/3$ by (W4);
- by (3.5)–(3.6), $G_2 < \frac{\kappa\varepsilon_1}{\lambda\underline{w}}\, c_{\Gamma^*}\lambda\underline{w}\,\frac{\kappa r+\alpha}{\kappa} = c_{\Gamma^*}\varepsilon_1(\kappa r + \alpha) \le \alpha/3$ by (W3).

With (3.8),

$$\max_{j \in S^c,\ \hat{w}_j < \infty}|\tilde{z}_j| < (1 - \alpha) + \tfrac{\alpha}{3} + \tfrac{\alpha}{3} + \tfrac{\alpha}{3} = 1. \tag{3.10}$$

If $\hat{w}_j = \infty$ for every $j \in S^c$ (the pilot is zero on all of S^c), Step 3 is empty: every S^c-coordinate is fixed at 0 by the constraint, so there is no $\tilde{z}_j$ to bound and $\hat{m}_{S^c} = 0$ holds automatically.

Step 4 (the witness is the unique minimiser; $\hat{S} \subseteq S$). The pair $(\tilde{m}, \tilde{z})$ satisfies (3.1):

- the S-rows by (3.3) and the rows $j \in S^c$ with $\hat{w}_j < \infty$ by construction;
- $\tilde{m}_j = 0$ wherever $\hat{w}_j = \infty$, since all infinite weights lie in S^c;

- $\tilde{z}$ is a valid subgradient, since $|\tilde{z}_j| < 1$ at the zero coordinates $j \in S^c$ with $\hat{w}_j < \infty$ by (3.10), and $\tilde{z}_S$ comes from Step 2.

So $\tilde{m}$ is a minimiser.

Every minimiser has support in S (Wainwright 2009, Lemma 1, cited by DDK on p. 532). Strict dual feasibility implies hat $\hat{S} \subseteq S$.

Uniqueness. Every minimiser lives on $\{m_{S^c} = 0\}$, where the objective is strictly convex (Step 1). Part (a) follows from (3.5)–(3.6). $\square_{(a)}$

Part (b). For $(j,k) \in S$, (a) gives $|\hat{M}_{jk} - M^*_{jk}| \le b_w < \beta_{\min} \le |M^*_{jk}|$. Hence $\hat{M}_{jk} \neq 0$ with the sign of M^*_{jk}, so $S \subseteq \hat{S}$, and with (a), $\hat{S} = S$.

4 Theorem W.2 (probabilistic; the analogue of DDK Theorem 2)

DDK's tools

- Lemma E.6: $|||\Delta_\Sigma|||_2 < \min\{\varepsilon_1/(\sqrt{d}(4 + 8c_{\Sigma^*})),\ \varepsilon_2/(2c_C)\}$ implies $|||(\Delta_\Gamma)_{\cdot S}|||_\infty < \varepsilon_1$ and $\|\Delta_g\|_\infty < \varepsilon_2$, with $c_{\Sigma^*} = |||\Sigma^*|||_2$ and $c_C = \|\operatorname{vec}(C)\|_2$.
- Corollary E.5 (from Wainwright 2019, Thm 6.5) bounds $|||\Delta_\Sigma|||_2$.
- As in DDK,

$$\tilde{c} = \max\Big\{\frac{4\max\{1, c^2_{\Sigma^*}\}(4 + 8c_{\Sigma^*})^2}{c_3},\ 16c_1^2 c^2_{\Sigma^*}(4 + 8c_{\Sigma^*})^2,\ \frac{16\max\{1, c^2_{\Sigma^*}\}c_C^2}{c_3},\ 64c_1^2 c^2_{\Sigma^*} c_C^2\Big\},$$

  with the universal constants c_1, c_2, c_3 of Theorem E.4.

Theorem W.2. *In the setting of DDK Theorem 2, let $\kappa \ge 1 + \rho$, $\alpha \in (0,1]$ and $\bar{r} \ge 0$. Let $\mathcal{E}_w$ be an event on which the weights satisfy $\bar{w}_S < \infty$, $\rho_w \le 1 - \alpha$ and $r \le \bar{r}$. Put*

$$\varepsilon := \sqrt{\tau_1 \tilde{c}\, d\, p/n}, \qquad c_w := \frac{3c_{\Gamma^*}(\kappa\bar{r} + \alpha)}{\alpha} \qquad \Big(\textit{DDK: } c_* = \frac{6c_{\Gamma^*}}{\alpha}\Big).$$

Suppose $n > \tau_1 \tilde{c}\, d\, p \max\{c_w^2, 1/4\}$ for some $\tau_1 > 1$, and that on $\mathcal{E}_w$

$$\lambda\, \underline{w} > \frac{3\kappa}{\alpha}\max\{c_{M^*}, 1\}\,\varepsilon.$$

For the implemented weights ($\hat{w} \ge 1$) it suffices that $\lambda > \frac{3\kappa}{\alpha}\max\{c_{M^*}, 1\}\varepsilon$; this is DDK's condition with $2 - \alpha$ replaced by κ. Then, with probability at least $1 - c_2\exp(-\tau_1 p) - P(\mathcal{E}_w^c)$:

(a) *the minimiser of (1.1) is unique, $\hat{S} \subseteq S$, and*

$$\|\hat{M} - M^*\|_\infty \le \frac{c_{\Gamma^*}}{1 - c_{\Gamma^*}\varepsilon}\big((c_{M^*} + 1)\varepsilon + \lambda\bar{w}_S\big) < c_{\Gamma^*}\lambda\Big(\bar{w}_S + \frac{\alpha}{\kappa}\underline{w}\Big);$$

(b) *if $\beta_{\min}$ exceeds the first bound in (a), then $\hat{S} = S$ with the signs of M^*.*

Proof. DDK Appendix F, steps 1–3, with $\varepsilon_1 = \varepsilon_2 = \varepsilon$.

- **Steps 1–2 are unchanged.** $\varepsilon/(\sqrt{d}(4+8c_{\Sigma^*}))$ and $\varepsilon/(2c_C)$ lie in the admissible range of Corollary E.5. The upper end uses only $\max\{c_w^2, 1/4\} \geq 1/4$, the lower end only $\tilde{c}$. So with probability at least $1 - c_2\exp(-\tau_1 p)$, $\|\!|(\Delta_\Gamma)_{\cdot S}|\!\|_\infty < \varepsilon$ and $\|\Delta_g\|_\infty < \varepsilon$. Both are bounds on the same random variable $\|\!|\Delta_\Sigma|\!\|_2$, so no union bound is needed.
- **Step 3 replaces c_* by c_w.** $\varepsilon < 1/\max\{c_w, 1/2\} \leq 1/c_w = \alpha/(3c_{\Gamma^*}(\kappa\bar{r}+\alpha)) \leq \alpha/(3c_{\Gamma^*}(\kappa r+\alpha))$ on $\mathcal{E}_w$, which is (W3).
- **Conclude.** The λ-condition is (W4), since $\max\{c_{M^*}\varepsilon, \varepsilon\} = \max\{c_{M^*}, 1\}\varepsilon$. Apply Theorem W.1 on $\mathcal{E}_w \cap \{\text{concentration}\}$. □

Reading. With $\hat{w} \equiv 1$, $\kappa = 2-\alpha$ and $\bar{r} = 1$, Theorem W.2 is DDK's Theorem 2. Sample size generally depends on the quality of weights via c_w and $\bar{r}$. With a good pilot, $\bar{r} \to 0$ and one may take $\alpha \to 1$, so $c_w \to 3c_{\Gamma^*}$. The sample size then no longer depends on an irrepresentability gap: it becomes $n \gtrsim \tilde{c}\, dp \cdot 9c_{\Gamma^*}^2$, against DDK's $\tilde{c}\, dp \cdot 36c_{\Gamma^*}^2/\alpha^2$ with α the (often non-existent) lasso gap. The order $\Omega(dp)$ in (d,p) is unchanged.

Corollary W.3 (A pilot close to the truth)**.** *Let $\|\hat{M}^0 - M^*\|_\infty \leq \delta < \beta_{\min}/(1+\rho)$ and put $\bar{r} := \delta/(\beta_{\min}-\delta)$, $\alpha := 1-\rho\bar{r}$. Then $\alpha \in (0,1]$ and*

(i) $\bar{w}_S < \infty$, and all infinite weights lie in S^c;

(ii) $r \leq \bar{r}$;

(iii) $\rho_w \leq \rho r \leq 1-\alpha$, i.e. (W2) holds.

*Hence, **whatever the size of** ρ, Theorem W.1 applies with this α whenever (W1), (W3), (W4) hold, and Theorem W.2 applies with $\mathcal{E}_w = \{\|\hat{M}^0 - M^*\|_\infty \leq \delta\}$ and this $\bar{r}$.*

Proof. Multiplying by $\beta_{\min} - \delta > 0$, $\rho\bar{r} < 1 \iff (1+\rho)\delta < \beta_{\min}$, so $\alpha \in (0,1]$.

(i) For $j \in S^c$, $M_j^* = 0$, so $|\hat{M}_j^0| = |\hat{M}_j^0 - M_j^*| \leq \delta$, i.e. weights can be infinite. For $k \in S$, the reverse triangle inequality gives $|\hat{M}_k^0| \geq |M_k^*| - |\hat{M}_k^0 - M_k^*| \geq \beta_{\min} - \delta > 0$. So the pilot has no zero on S: every $\hat{w}_k = \hat{\mu}/|\hat{M}_k^0|$, $k \in S$, is finite, and infinite weights (pilot zeros) occur only where $M_j^* = 0$.

(ii) Since $\hat{w}_j = \hat{\mu}/|\hat{M}_j^0|$ is decreasing in $|\hat{M}_j^0|$ (with $\hat{\mu}/0 := \infty$), the two bounds in (i) give

$$r = \frac{\bar{w}_S}{\underline{w}} = \frac{\max_{j\in S^c}|\hat{M}_j^0|}{\min_{k\in S}|\hat{M}_k^0|} \leq \frac{\delta}{\beta_{\min}-\delta} = \bar{r}.$$

(iii) Bounding $\hat{w}_k \leq \bar{w}_S$ ($k \in S$) and $\hat{w}_j \geq \underline{w}$ ($j \in S^c$) in (1.2) gives $\rho_w \leq r\,|||P|||_\infty = \rho r \leq \rho\bar{r} = 1-\alpha$.

Finally, (i) and (iii) are the hypotheses of Theorem W.1 that involve the pilot, and (i)–(iii) are exactly what Theorem W.2 requires to hold on $\mathcal{E}_w$. □

On $\beta_{\min}$: Since the diagonal is penalised, $\beta_{\min}$ includes it. Diagonal entries are typically the largest, so in practice $\beta_{\min}$ is the weakest edge. As $\delta \uparrow \beta_{\min}/(1+\rho)$, $\alpha \downarrow 0$ and (W3)–(W4) become more demanding: ρ no longer restricts the model, only the required pilot accuracy. **Getting $P(\mathcal{E}_w), \bar{r}$:** The assumption for this corollary is more harsh than $(W2)$. But it would give us $\bar{r}$ that is necessary for the sample size computation and tell us whether $\mathcal{E}_w$ holds.