


University of Stuttgart
Germany

inm Institute for
Nonlinear Mechanics

Nonsmooth Dynamics

a journey from convex analysis to computer code

Remco Leine



ENOC 2024
TU Delft
22- 26 July 2024




Simulation of granular media



10 hours of simulation
time on the
super computer of the
University of Stuttgart?


Not really...

Application examples of Nonsmooth Dynamics



Simulations by: Giuseppe Capobianco, Simon Eugster, Jonas Breuling, R.L.

A motivating example: The block-on-belt system



equation of motion



$$m\ddot{q} + c\dot{q} + kq = F_T$$

relative velocity

$$\gamma_T = \dot{q} - v_{\text{dr}}$$


friction law

$$(\gamma_T, -F_T) \in \text{Graph } \mathcal{F}$$




Decomposition of the Stribeck friction law


Stribeck



differentiable part




pure Coulomb friction



Coulomb friction is a
set-valued force law expressing
the difference in causality
in the stick and slip phase


constraint force
in stick

impressed force
in slip




Regularization methods

Arctangent smoothing



$$-\lambda_T = \mu \lambda_N \frac{2}{\pi} \arctan(\varepsilon \gamma_T)$$

stiff ODE
in stick phase




LuGre model

$$-\lambda_T = \sigma_0 z + \sigma_1 \dot{z}$$


$$\dot{z} = \gamma_T - \frac{\sigma_0}{\mu \lambda_N} |\gamma_T| z$$

stiff ODE
in stick and slip phase



Coulomb friction as a normal cone inclusion


Primal form



$-\lambda_T \in \mu\lambda_N \text{ Sign}(\gamma_T)$

set-valued sign function

Dual form





$\gamma_T \in N_{C_T}(-\lambda_T)$

normal cone

force reservoir: $C_T = [-\mu\lambda_N, \mu\lambda_N]$

Normal cone from convex analysis




$$N_C(x) = \{y \mid y^T(x^* - x) \leq 0, x \in C, \forall x^* \in C\}$$


$x \in \text{bdry } C \Rightarrow$ normal cone $N_C(x)$ consists of outward normal ray(s)

$x \in \text{int } C \Rightarrow$ normal cone $N_C(x)$ is zero

Coulomb's spatial friction law



Proximal point to a convex set



proximal point function:

$$\mathbf{x} = \text{prox}_C(\mathbf{z})$$

$$\Updownarrow$$

$$\frac{1}{r}(\mathbf{z} - \mathbf{x}) \in N_C(\mathbf{x}), \quad r > 0$$

substitute:

$$\begin{aligned} \mathbf{z} = -\boldsymbol{\lambda} + r\boldsymbol{\gamma} \\ \mathbf{x} = -\boldsymbol{\lambda} \end{aligned} \Rightarrow \begin{cases} \boldsymbol{\gamma} = \frac{1}{r}(\mathbf{z} - \mathbf{x}) \\ \boldsymbol{\lambda} = -\mathbf{x} \end{cases}$$


$$\boldsymbol{\gamma} \in N_C(-\boldsymbol{\lambda}) \iff -\boldsymbol{\lambda} = \text{prox}_C(-\boldsymbol{\lambda} + r\boldsymbol{\gamma}) \quad r > 0$$

normal cone inclusion


implicit equation

Coulomb friction as proximal point equation

Primal form $-\lambda_T \in \mu\lambda_N \operatorname{Sign}(\gamma_T)$




Dual form $\gamma_T \in N_{C_T}(-\lambda_T)$



Proximal point equation

$$-\lambda_T = \operatorname{prox}_{C_T}(-\lambda_T + r\gamma_T)$$

force reservoir: $C_T = [-\mu\lambda_N, \mu\lambda_N]$




The block-on-belt system as DAE

Differential inclusion

equation of motion $m\ddot{q} + c\dot{q} + kq = F_D(\gamma_T) + \lambda_T$

relative velocity $\gamma_T = \dot{q} - v_{\text{dr}}$

friction law $-\lambda_T \in \mu\lambda_N \text{Sign}(\gamma_T)$



$$-\lambda_T \in \mu\lambda_N \text{Sign}(\gamma_T) \iff \gamma_T \in N_{C_T}(-\lambda_T) \iff -\lambda_T = \text{prox}_{C_T}(-\lambda_T + r\gamma_T)$$

Differential algebraic equation

$$\begin{aligned}\dot{q} &= u \\ \dot{u} &= -\frac{c}{m}u - \frac{k}{m}q + \frac{1}{m}F_D(\gamma_T) + \frac{1}{m}\lambda_T\end{aligned}$$

$$0 = \lambda_T + \text{prox}_{C_T}(-\lambda_T + r\gamma_T)$$

with $\gamma_T = u - v_{\text{dr}}$

}

Differential algebraic equation

$$\dot{x} = f(x, \lambda)$$

$$0 = c(x, \lambda)$$

$$x = (q \quad u)^T, \quad \lambda = \lambda_T$$

A semi-implicit index-2 DAE solver

Differential algebraic equation

$$\dot{x} = f(x, \lambda)$$

Hessenberg-2 DAE

$$0 = c(x, \lambda)$$

Semi-implicit scheme

$$x_{i+1} = x_i + \Delta t f(x_i, \lambda_{i+1})$$

$$0 = c(x_{i+1}, \lambda_{i+1})$$

in each step solve $0 = c(x_i + \Delta t f(x_i, \lambda_{i+1}), \lambda_{i+1})$

solvable if $\det(c_x f_\lambda \Delta t + c_\lambda) \neq 0$

index must be 1 or 2

Semi-implicit scheme


$$q_{i+1} = q_i + \Delta t u_i$$

event capturing method!!

$$u_{i+1} = u_i + \frac{\Delta t}{m} (-c u_i - k q_i + F_D(\gamma_{T,i}) + \lambda_{T,i+1})$$

$$0 = \lambda_{T,i+1} + \text{prox}_{C_T}(-\lambda_{T,i+1} + r \gamma_{T,i+1})$$

with $\gamma_{T,i} = u_i - v_{\text{dr}}$



Multibody system with set-valued force law

Multibody system

kinematics $\dot{\mathbf{q}} = \mathbf{u}$

kinetics $\mathbf{M}\dot{\mathbf{u}} - \mathbf{h}(\mathbf{q}, \mathbf{u}) = \mathbf{W}\boldsymbol{\lambda}$

kinematic var. $\boldsymbol{\gamma} = \mathbf{W}^T \mathbf{u} + \boldsymbol{\chi}$

force law $\boldsymbol{\gamma} \in N_C(-\boldsymbol{\lambda})$

Semi-implicit scheme

$$\mathbf{q}_{i+1} = \mathbf{q}_i + \Delta t \mathbf{u}_i \quad h(\mathbf{q}_i, \mathbf{u}_i)$$

$$\mathbf{M}(\mathbf{u}_{i+1} - \mathbf{u}_i) - \Delta t \mathbf{h}_i = \Delta t \mathbf{W} \boldsymbol{\lambda}_{i+1}$$

$$\boldsymbol{\gamma}_{i+1} = \mathbf{W}^T \mathbf{u}_{i+1} + \boldsymbol{\chi}$$

$$\boldsymbol{\gamma}_{i+1} \in N_C(-\boldsymbol{\lambda}_{i+1})$$

$$\boldsymbol{\gamma}_{i+1} = \underbrace{\Delta t \mathbf{W}^T \mathbf{M}^{-1} \mathbf{W} \boldsymbol{\lambda}_{i+1}}_A + \underbrace{\mathbf{W}^T (\mathbf{u}_i + \Delta t \mathbf{M}^{-1} \mathbf{h}_i)}_{-\mathbf{b}} + \boldsymbol{\chi}$$

$$\boldsymbol{\gamma}_{i+1} \in N_C(-\boldsymbol{\lambda}_{i+1})$$

$$\mathbf{y} = -\boldsymbol{\gamma}_{i+1}, \quad \mathbf{x} = -\boldsymbol{\lambda}_{i+1}$$

Linear normal cone inclusion problem

$$\mathbf{y} = \mathbf{A}\mathbf{x} + \mathbf{b}$$

$$-\mathbf{y} \in N_C(\mathbf{x})$$

Overview

Constrained optimization problem

$$\min_{\mathbf{x} \in C} \frac{1}{2} \mathbf{x}^T \mathbf{A} \mathbf{x} + \mathbf{b}^T \mathbf{x}$$

$$\uparrow \mathbf{A} = \mathbf{A}^T > 0$$

Variational
inequality

$$\mathbf{y} = \mathbf{A}\mathbf{x} + \mathbf{b}$$

$$\forall \mathbf{x}^* \in C: \mathbf{y}^T (\mathbf{x}^* - \mathbf{x}) \geq 0$$

 \Leftrightarrow

Normal cone inclusion
problem

$$\mathbf{y} = \mathbf{A}\mathbf{x} + \mathbf{b}$$

$$-\mathbf{y} \in N_C(\mathbf{x})$$

 \Leftrightarrow

Proximal point
equation problem

$$\mathbf{y} = \mathbf{A}\mathbf{x} + \mathbf{b}$$

$$\mathbf{x} = \text{prox}_C(\mathbf{x} - r\mathbf{y})$$

$$\Downarrow C = \mathbb{R}_0^{n+}$$

Linear complementarity problem

$$\mathbf{y} = \mathbf{A}\mathbf{x} + \mathbf{b}$$

$$\mathbf{0} \leq \mathbf{y} \perp \mathbf{x} \geq \mathbf{0}$$

Numerical solution

Normal cone inclusion problem

$$\mathbf{y} = \mathbf{A}\mathbf{x} + \mathbf{b}$$

$$-\mathbf{y} \in N_C(\mathbf{x})$$

\Leftrightarrow

Proximal point equation problem

$$\mathbf{y} = \mathbf{A}\mathbf{x} + \mathbf{b}$$

$$\mathbf{x} = \text{prox}_C(\mathbf{x} - r\mathbf{y})$$

$r > 0$ can be chosen arbitrarily!

Semi-smooth Newton method

$$f(\mathbf{x}) := \mathbf{x} - \text{prox}_C(\mathbf{x} - r(\mathbf{A}\mathbf{x} + \mathbf{b}))$$

for some initial guess \mathbf{x}_0 iterate

$$\mathbf{x}_{k+1} = \mathbf{x}_k - \mathbf{J}^{-1} f(\mathbf{x}_k), \quad \mathbf{J} \in \partial_C f(\mathbf{x}_k)$$

until $\|\mathbf{x}_{k+1} - \mathbf{x}_k\| < tol$



Proximal point iteration

for some initial guess \mathbf{x}_0 iterate

$$\mathbf{y}_k = \mathbf{A}\mathbf{x}_k + \mathbf{b}$$


$$\mathbf{x}_{k+1} = \text{prox}_C(\mathbf{x}_k - r\mathbf{y}_k)$$

until $\|\mathbf{x}_{k+1} - \mathbf{x}_k\| < tol$

r too small: slow convergence

r too large: divergence

Contact law in normal direction



Signorini's law


no penetration $g_N \geq 0$

no adhesion $\lambda_N \geq 0$


no distance effects $g_N > 0 \implies \lambda_N = 0$

$$0 \leq g_N \perp \lambda_N \geq 0$$




Primal form $-\lambda_N \in N_{\mathbb{R}_0^+}(g_N)$



Dual form $g_N \in N_{C_N}(-\lambda_N)$



A Signorini contact problem in statics



kinematics:


$$g_N = b - x$$

equilibrium of forces: $F_{\text{spring}} + \lambda_N = 0$

constitutive equations:

spring law $F_{\text{spring}} = kx$

contact law $0 \leq g_N \perp \lambda_N \geq 0$



Contact problem

$$g_N = \frac{1}{k} \lambda_N + b$$

$$0 \leq g_N \perp \lambda_N \geq 0$$

Linear complementarity problem

$$\mathbf{y} = \mathbf{Ax} + \mathbf{b}$$


$$\mathbf{0} \leq \mathbf{y} \perp \mathbf{x} \geq \mathbf{0}$$

A multi-DOF Signorini contact problem in statics

equilibrium of forces: $\mathbf{K}\mathbf{q} = \mathbf{W}_N\boldsymbol{\lambda}_N + \mathbf{f}$

kinematics: $\mathbf{g}_N = \mathbf{W}_N^T \mathbf{q} + \mathbf{g}_0$

constitutive equation: $\mathbf{0} \leq \mathbf{g}_N \perp \boldsymbol{\lambda}_N \geq \mathbf{0}$



Contact problem

$$\mathbf{g}_N = \underbrace{\mathbf{W}_N^T \mathbf{K}^{-1} \mathbf{W}_N}_{A} \boldsymbol{\lambda}_N + \underbrace{\mathbf{g}_0 + \mathbf{W}_N^T \mathbf{K}^{-1} \mathbf{f}}_{b}$$

$$\mathbf{0} \leq \mathbf{g}_N \perp \boldsymbol{\lambda}_N \geq \mathbf{0}$$

Linear complementarity problem

$$\mathbf{y} = \mathbf{Ax} + \mathbf{b}$$

$$\mathbf{0} \leq \mathbf{y} \perp \mathbf{x} \geq \mathbf{0}$$

Proximal point equation problem

$$\mathbf{y} = \mathbf{Ax} + \mathbf{b}$$

$$\mathbf{x} = \text{prox}_C(\mathbf{x} - r\mathbf{y})$$

\Leftrightarrow

Dynamics: the evolutionary problem

Nonimpulsive dynamics

kinematics: $\dot{\mathbf{q}} = \mathbf{u}$

equation of motion: $\mathbf{M}(\mathbf{q})\ddot{\mathbf{u}} - \mathbf{h}(\mathbf{q}, \mathbf{u}) = \mathbf{W}_N \boldsymbol{\lambda}_N + \mathbf{W}_T \boldsymbol{\lambda}_T$

normal contact law: $g_{Nj} \in N_{C_N}(-\lambda_{Nj})$

friction law: $\gamma_{Tj} \in N_{C_T(\lambda_{Nj})}(-\lambda_{Tj})$






Impulsive dynamics

impact equation: $\mathbf{M}(\mathbf{q})(\mathbf{u}^+ - \mathbf{u}^-) = \mathbf{W}_N \boldsymbol{\Lambda}_N + \mathbf{W}_T \boldsymbol{\Lambda}_T$

normal impact law: $\gamma_{Nj}^+ + e_{Nj} \gamma_{Nj}^- \in N_{C_N}(-\Lambda_{Nj}), \quad j \in \mathcal{I}_N = \{j \mid g_{Nj} = 0\}$

tangential impact law: $\gamma_{Tj}^+ + e_{Tj} \gamma_{Tj}^- \in N_{C_T(\Lambda_{Nj})}(-\Lambda_{Tj}), \quad j \in \mathcal{I}_N$

Accumulation points of collisions



The Woodpecker Toy: a simple code

```

function [t,q,u] = woodpeckertimestepping(t0,te,q0,u0,N)
mM = 3e-4; JM = 5e-9; mS = 4.5e-3; JS = 7e-7; lM = 0.01; lG = 0.015; lS = 0.0201; %system
rM = 0.0031; r0 = 0.0025; hM = 0.0058; hS = 0.02; grav = 9.81; cphi = 0.0056; %parameters
mu = 0.3; eN = [0.5;0;0]; r = 1e-4; tol = 1e-8;
M = [(mS+mM) mS*lM mS*lG; mS*lM (JM+mS*lM^2) mS*lM*lG; mS*lG mS*lM*lG (JS+mS*lG^2)]; %constant
WN = [0 0 0; 0 hM -hM; -hS 0 0]; WT = [1 1 1; lM rM rM; lG-lS 0 0]; %matrices

t = linspace(t0,te,N); dt = (te-t0)/(N-1); %time stamp and time step
q = [q0 zeros(3,N-1)]; u = [u0 zeros(3,N-1)]; PN = zeros(3,1); PT = zeros(3,1); %memory allocation

```

```

for i=1:(N-1)
    tM = t(i)+dt/2; qM = q(:,i) + dt/2*u(:,i); %midpoint
    h = [- (mS+mM)*grav; -cphi*(qM(2)-qM(3))-mS*grav*lM; -cphi*(qM(3)-qM(2))-mS*grav*lG];
    gN = [lM+lG-lS-r0-hS*qM(3);rM-r0+hM*qM(2);rM-r0-hM*qM(2)];
    I = find(gN<=0)'; %index set of closed contacts
    if ~isempty(I)
        gammaNA = WN(:,I)'*u(:,i);
        converged = false;
        while ~converged %fixed point iteration
            u(:,i+1) = u(:,i) + M\ (h*dt + WN(:,I)*PN(I) + WT(:,I)*PT(I));
            gammaNE = WN(:,I)'*u(:,i+1); gammate = WT(:,I)'*u(:,i+1);
            PNnew = proxCN(PN(I)-r*(gammaNE + eN(I).*gammaNA));
            PTnew = proxCT(PT(I)-r*gammate,mu*PN(I));
            converged = norm(PN(I)-PNnew)+norm(PT(I)-PTnew) < tol;
            PN(I) = PNnew; PT(I) = PTnew;
        end
    else %if all contacts are open
        u(:,i+1) = u(:,i) + M\ (h*dt);
    end
    q(:,i+1) = qM + u(:,i+1)*dt/2; %end timestep
end

```


```

function y = proxCN(x), y = max(0,x); %proximal point function for CN = R_0^+
function y = proxCT(x,a), y = max(min(x,a),-a); %proximal point function for CT = [-a,a]

```

only 34 lines
of code...

Moreau-timestepping scheme



A semi-implicit index-2 DAE solver

Differential algebraic equation

$$\dot{x} = f(x, \lambda)$$

Hessenberg-2 DAE

$$0 = c(x, \lambda)$$

Semi-implicit scheme

$$x_{i+1} = x_i + \Delta t f(x_i, \lambda_{i+1})$$

$$0 = c(x_{i+1}, \lambda_{i+1})$$

in each step solve $0 = c(x_i + \Delta t f(x_i, \lambda_{i+1}), \lambda_{i+1})$

solvable if $\det(c_x f_\lambda \Delta t + c_\lambda) \neq 0$

index must be 1 or 2

Semi-implicit scheme


$$q_{i+1} = q_i + \Delta t u_i$$

event capturing method!!

$$u_{i+1} = u_i + \frac{\Delta t}{m} (-c u_i - k q_i + F_D(\gamma_{T,i}) + \lambda_{T,i+1})$$

$$0 = \lambda_{T,i+1} + \text{prox}_{C_T}(-\lambda_{T,i+1} + r \gamma_{T,i+1})$$

with $\gamma_{T,i} = u_i - v_{\text{dr}}$



A semi-implicit index-3 DAE solver

Differential algebraic equation

$$\dot{\mathbf{y}} = \mathbf{g}(\mathbf{y}, \mathbf{z})$$

$$\dot{\mathbf{z}} = \mathbf{h}(\mathbf{y}, \mathbf{z}, \boldsymbol{\lambda}) \quad \text{Hessenberg-3 DAE}$$

$$\mathbf{0} = \mathbf{c}(\mathbf{y}, \mathbf{z}, \boldsymbol{\lambda})$$

Semi-implicit scheme

$$\mathbf{y}_{i+1} = \mathbf{y}_i + \Delta t \mathbf{g}(\mathbf{y}_i, \mathbf{z}_{i+1})$$

$$\mathbf{z}_{i+1} = \mathbf{z}_i + \Delta t \mathbf{h}(\mathbf{y}_i, \mathbf{z}_i, \boldsymbol{\lambda}_{i+1})$$

$$\mathbf{0} = \mathbf{c}(\mathbf{y}_{i+1}, \mathbf{z}_{i+1}, \boldsymbol{\lambda}_{i+1})$$

in each step solve

$$\mathbf{0} = \mathbf{c}(\mathbf{y}_i + \Delta t \mathbf{g}(\mathbf{y}_i, \mathbf{z}_i + \Delta t \mathbf{h}(\mathbf{y}_i, \mathbf{z}_i, \boldsymbol{\lambda}_{i+1}), \boldsymbol{\lambda}_{i+1}), \mathbf{z}_i + \Delta t \mathbf{h}(\mathbf{y}_i, \mathbf{z}_i, \boldsymbol{\lambda}_{i+1}), \boldsymbol{\lambda}_{i+1})$$

solvable if $\det(\mathbf{c}_y \mathbf{g}_z \mathbf{h}_\lambda \Delta t^2 + \mathbf{c}_z \mathbf{h}_\lambda \Delta t + \mathbf{c}_\lambda) \neq 0$ **index must be 3 or less**

Semi-implicit index-3 DAE solver in mechanics

Multibody system (frictionless)

kinematics $\dot{\mathbf{q}} = \mathbf{u}$

kinetics $\mathbf{M}\dot{\mathbf{u}} - \mathbf{h}(\mathbf{q}, \mathbf{u}) = \mathbf{W}_N \boldsymbol{\lambda}_N$ impact law is missing!!!

contact dist. $\mathbf{g}_N = \mathbf{g}_N(\mathbf{q})$

Signorini's law $\mathbf{g}_N \in N_{C_N}(-\boldsymbol{\lambda}_N)$

$$\left. \begin{array}{l} \mathbf{g}_N = \mathbf{g}_N(\mathbf{q}) \\ \mathbf{g}_N \in N_{C_N}(-\boldsymbol{\lambda}_N) \end{array} \right\} \mathbf{0} = \boldsymbol{\lambda}_N + \text{prox}_{C_N}(-\boldsymbol{\lambda}_N + r\mathbf{g}_N(\mathbf{q}))$$

Semi-implicit scheme

$$\begin{aligned} \mathbf{q}_{i+1} &= \mathbf{q}_i + \Delta t \mathbf{u}_{i+1} \\ \mathbf{M}(\mathbf{u}_{i+1} - \mathbf{u}_i) - \Delta t \mathbf{h}_i &= \mathbf{W}_N \mathbf{P}_{N,i+1} \end{aligned}$$

$$\mathbf{g}_{N,i+1} = \mathbf{g}_N(\mathbf{q}_{i+1})$$

$$\mathbf{g}_{N,i+1} \in N_{C_N}(-\mathbf{P}_{N,i+1})$$

agrees for single-contact problems with the inelastic Schatzman-Paoli scheme


SCHATZMAN & PAOLI, 2002

Semi-implicit index-3 DAE solver in mechanics

```
function [t,q,u] = semi_implicit_scheme(sys,t0,te,q0,u0,N,tol)
% Semi-implicit scheme for frictionless inelastic impacts
t = linspace(t0,te,N); dt = (te-t0) / (N-1); %time stamp and time step
q = zeros(sys.dim_q,N); u = zeros(sys.dim_q,N); PN = zeros(length(sys.I),1);
q(:,1) = q0; u(:,1) = u0; %initial condition

for i=1:(N-1)
    h = sys.h(t(i),q(:,i),u(:,i));
    M = sys.M(t(i),q(:,i));
    WN = sys.WChi(t(i),q(:,i),sys.I);
    converged = false;
    while ~converged
        u(:,i+1) = u(:,i) + M\ (h*dt + WN*PN);
        q(:,i+1) = q(:,i) + u(:,i+1)*dt;
        gN = sys.gN(t(i+1),q(:,i+1),sys.I);
        PNnew = proxCN(PN-sys.r/dt*gN);
        converged = norm(PN-PNnew) < tol;
        PN = PNnew;
    end
end

function y = proxCN(x), y = max(0,x); %proximal point function for CN = R_0^+
```



impact law is missing!!!

Schatzman-Paoli-type contact law

completely inelastic



$$\mathbf{g}_{N,i+1} \in N_{C_N}(-\mathbf{P}_{N,i+1})$$

partially elastic

$$\mathbf{g}_{N,i+1} + e_N \mathbf{g}_{N,i-1} \in N_{C_N}(-\mathbf{P}_{N,i+1})$$

$$0 \leq e_N \leq 1$$

Single contact



impact is resolved over two time-steps

Multi contact



$$\Delta t = 0.001005$$



$$\Delta t = 0.001$$

wrong results for $e_N > 0$

if we want frictional and/or elastic impacts...

Constitutive equations

friction laws

anisotropic Coulomb friction

Coulomb-Contensou friction

impact laws

gen. Newton's impact law

Poisson impact law

Glocker, Brogliato, Panagiotopoulos, Curnier, Leine, de Saxé, ...

Measure differential inclusions

Moreau, Monteiro Marques, Schatzman, Ballard, Stewart, ...

Event capturing schemes

Moreau's timestepping scheme

Moreau, Jean

Nonsmooth generalized alpha method

Acary, Brüls, Capobianco

Moreau timestepping

Measure differential inclusion

$$d\mathbf{q} = \mathbf{u} dt$$

$$\mathbf{M}(\mathbf{q})d\mathbf{u} - \mathbf{h}(t, \mathbf{q}, \mathbf{u})dt = \mathbf{W}_N d\mathbf{P}_N$$

$$\xi_{Nj} = \gamma_{Nj}^+ + e_N \gamma_{Nj}^-$$

$$g_{Nj}(\mathbf{q}) = 0 : \quad \xi_{Nj} \in N_{C_N}(-dP_{Nj})$$

$$g_{Nj}(\mathbf{q}) > 0 : \quad dP_{Nj} = 0$$

Time stepping scheme (Moreau)

$$\mathbf{q}_M = \mathbf{q}_i + \mathbf{u}_i \frac{\Delta t}{2}$$

$$\mathbf{M}(\mathbf{q}_M)(\mathbf{u}_{i+1} - \mathbf{u}_i) - \mathbf{h}(\mathbf{q}_M, \mathbf{u}_i) \Delta t = \mathbf{W}_N \mathbf{P}_{N,i+1}$$

$$\xi_{Nj,i+1} = \gamma_{Nj,i+1} + e_N \gamma_{Nj,i}$$


$$g_{Nj}(\mathbf{q}_M) \leq 0 : \quad \xi_{Nj,i+1} \in N_{C_N}(-P_{Nj,i+1})$$

$$g_{Nj}(\mathbf{q}_M) > 0 : \quad P_{Nj,i+1} = 0$$


$$\mathbf{q}_{i+1} = \mathbf{q}_M + \frac{1}{2}\mathbf{u}_{i+1}\Delta t$$









contact law on velocity level!



MOREAU, JEAN, 1988



if we want to do something...
...we can do it



Moreau timestepping

Measure differential inclusion

$$d\mathbf{q} = \mathbf{u} dt$$

$$\mathbf{M}(\mathbf{q})d\mathbf{u} - \mathbf{h}(\mathbf{q}, \mathbf{u})dt = \mathbf{W}_N d\mathbf{P}_N$$

$$\xi_{Nj} = \gamma_{Nj}^+ + e_N \gamma_{Nj}^-$$

$$g_{Nj}(\mathbf{q}) = 0 : \quad \xi_{Nj} \in N_{C_N}(-dP_{Nj})$$

$$g_{Nj}(\mathbf{q}) > 0 : \quad dP_{Nj} = 0$$

Time stepping scheme (Moreau)

$$\mathbf{q}_M = \mathbf{q}_i + \mathbf{u}_i \frac{\Delta t}{2}$$

$$\mathbf{M}(\mathbf{q}_M)(\mathbf{u}_{i+1} - \mathbf{u}_i) - \mathbf{h}(\mathbf{q}_M, \mathbf{u}_i)\Delta t = \mathbf{W}_N \mathbf{P}_{N,i+1}$$


$$\xi_{Nj,i+1} = \gamma_{Nj,i+1} + e_N \gamma_{Nj,i}$$

$$g_{Nj}(\mathbf{q}_M) \leq 0 : \quad \xi_{Nj,i+1} \in N_{C_N}(-P_{Nj,i+1})$$

$$g_{Nj}(\mathbf{q}_M) > 0 : \quad P_{Nj,i+1} = 0$$

$$\mathbf{q}_{i+1} = \mathbf{q}_M + \mathbf{u}_{i+1} \frac{\Delta t}{2}$$

drift!!
contact law on velocity level!



Nonsmooth RATTLE



Stage 1

$$\mathbf{q}_{i+1} = \mathbf{q}_i + \mathbf{u}_{i+\frac{1}{2}} \Delta t$$

$$\mathbf{M}(\mathbf{q}_i)(\mathbf{u}_{i+\frac{1}{2}} - \mathbf{u}_i) - \mathbf{h}(\mathbf{q}_i, \mathbf{u}_{i+\frac{1}{2}}) \frac{\Delta t}{2} = \mathbf{W}_N \mathbf{P}_{N,1}$$

$$\mathbf{g}_N(\mathbf{q}_{i+1}) \in N_{C_N}(-\mathbf{P}_{N,1})$$

contact law on position level

Stage 2


$$\mathbf{P}_N = \mathbf{P}_{N,1} + \mathbf{P}_{N,2}$$

$$\mathbf{M}(\mathbf{q}_{i+1})(\mathbf{u}_{i+1} - \mathbf{u}_{i+\frac{1}{2}}) - \mathbf{h}(\mathbf{q}_{i+1}, \mathbf{u}_{i+\frac{1}{2}}) \frac{\Delta t}{2} = \mathbf{W}_N \mathbf{P}_{N,2}$$

$$\xi_{Nj,i+1} = \gamma_{Nj,i+1} + e_N \gamma_{Nj,i}$$

$$P_{Nj,1} - r g_{Nj,i+1} \geq 0 : \quad \xi_{Nj,i+1} \in N_{C_N}(-P_{Nj}) \quad \text{contact law on velocity level}$$

$$P_{Nj,1} - r g_{Nj,i+1} < 0 : \quad P_{Nj} = 0$$





BREULING, CAPOBIANCO, EUGSTER, LEINE: A NONSMOOTH RATTLE ALGORITHM FOR MECHANICAL SYSTEMS WITH FRICITONAL UNILATERAL CONSTRAINTS, NONLINEAR ANALYSIS: HYBRID SYSTEMS, 2024



Simulation of rockfall protection nets



Rock model



Net model



Conclusions

- Numerical methods from nonsmooth dynamics are mature and easy to implement
- The analysis of nonsmooth systems requires the extension of Nonlinear Dynamics to nonsmooth dynamical systems:
 - stability theory
 - invariant manifolds (normal form theory, nonlinear modes)
 - bifurcation theory
 - chaos theory

With many thanks to some of my PhD students and co-workers:



Jonas
Breuling



Fabia
Bayer




Lisa
Eberhardt



Yassine
Karoui



Balkis
Youssef



Giuseppe
Capobianco
(Erlangen)



Simon
Eugster
(TU Eindhoven)



MATLAB codes