

Categorical predictors, interactions, logistic regression, and hierarchical models

Roger Levy

Massachusetts Institute of Technology

April 25, 2024

Categorical predictors: contrast matrices

- ▶ Often we want to bring effects of categorical predictors into our studies

Categorical predictors: contrast matrices

- ▶ Often we want to bring effects of categorical predictors into our studies
 - ▶ Is the relative clause being processed subject-extracted or object-extracted (Grodner and Gibson, 2005; Traxler et al., 2002)?
 - The reporter who attacked the senator (subject-extracted)*
 - The reporter who the senator attacked (object-extracted)*

Categorical predictors: contrast matrices

- ▶ Often we want to bring effects of categorical predictors into our studies
 - ▶ Is the relative clause being processed subject-extracted or object-extracted (Grodner and Gibson, 2005; Traxler et al., 2002)?

The reporter who attacked the senator (subject-extracted)
The reporter who the senator attacked (object-extracted)
 - ▶ Is the vowel lax or tense?

Categorical predictors: contrast matrices


- ▶ Often we want to bring effects of categorical predictors into our studies
 - ▶ Is the relative clause being processed subject-extracted or object-extracted (Grodner and Gibson, 2005; Traxler et al., 2002)?

The reporter who attacked the senator (subject-extracted)
The reporter who the senator attacked (object-extracted)
 - ▶ Is the vowel lax or tense?
- ▶ Example in word naming: the category of the initial sound of the word

burst	<i>bait</i>
frication	<i>chess</i>
other consonant	<i>wrist</i>
vowel	<i>inch</i>


Categorical predictors

- Initial evidence suggests that there may well be some real effects of initial phone class on word naming RT:



Categorical predictors

- ▶ Initial evidence suggests that there may well be some real effects of initial phone class on word naming RT:



- ▶ **Note:** within-class variance seems to differ across classes—this is sometimes called **heteroskedasticity**. We won't worry about this for now, but it's something to keep in mind and we can eventually incorporate ways to deal with it

Categorical predictors

- ▶ Initial phone class is a four-predictor categorical variable

Categorical predictors

- ▶ Initial phone class is a four-predictor categorical variable
- ▶ Hence, a model which encodes a different predicted mean for each initial phone class must have four parameters (plus a residual noise parameter σ^2)

Categorical predictors

- ▶ Initial phone class is a four-predictor categorical variable
- ▶ Hence, a model which encodes a different predicted mean for each initial phone class must have four parameters (plus a residual noise parameter σ^2)
- ▶ But a model in which phone class doesn't have an effect already has one parameter (plus σ^2):

$$RT = \alpha + \epsilon$$

Categorical predictors

- ▶ Initial phone class is a four-predictor categorical variable
- ▶ Hence, a model which encodes a different predicted mean for each initial phone class must have four parameters (plus a residual noise parameter σ^2)
- ▶ But a model in which phone class doesn't have an effect already has one parameter (plus σ^2):

$$RT = \alpha + \epsilon$$

- ▶ To fit the categorical predictors into the model, we use a **coding scheme** to represent the categorical values as numeric predictors

Categorical predictors

- ▶ Initial phone class is a four-predictor categorical variable
- ▶ Hence, a model which encodes a different predicted mean for each initial phone class must have four parameters (plus a residual noise parameter σ^2)
- ▶ But a model in which phone class doesn't have an effect already has one parameter (plus σ^2):

$$RT = \alpha + \epsilon$$

- ▶ To fit the categorical predictors into the model, we use a **coding scheme** to represent the categorical values as numeric predictors
- ▶ Many different coding schemes available (I'll cover three); critically, these models are **equivalent** in the following key sense: they can express the same set of probability distributions $P(Y|X)$!

Categorical predictors

- ▶ Initial phone class is a four-predictor categorical variable
- ▶ Hence, a model which encodes a different predicted mean for each initial phone class must have four parameters (plus a residual noise parameter σ^2)
- ▶ But a model in which phone class doesn't have an effect already has one parameter (plus σ^2):

$$RT = \alpha + \epsilon$$

- ▶ To fit the categorical predictors into the model, we use a **coding scheme** to represent the categorical values as numeric predictors
- ▶ Many different coding schemes available (I'll cover three); critically, these models are **equivalent** in the following key sense: they can express the same set of probability distributions $P(Y|X)$!
- ▶ Hence, your choice of coding scheme should generally be guided by how you want to interpret model parameters

Dummy coding

With dummy coding, you drop the intercept term and create one 0/1 “dummy” predictor per value of your categorical predictor:

Level of Init	X_1	X_2	X_3	X_4
burst	1	0	0	0
frication	0	1	0	0
other	0	0	1	0
vowel	0	0	0	1

Dummy coding

With dummy coding, you drop the intercept term and create one 0/1 “dummy” predictor per value of your categorical predictor:

Level of Init	X_1	X_2	X_3	X_4
burst	1	0	0	0
frication	0	1	0	0
other	0	0	1	0
vowel	0	0	0	1

This leads to the model

$$Y = \beta_1 X_1 + \beta_2 X_2 + \beta_3 X_3 + \beta_4 X_4 + \epsilon$$

Dummy coding

With dummy coding, you drop the intercept term and create one 0/1 “dummy” predictor per value of your categorical predictor:


Level of Init	X_1	X_2	X_3	X_4
burst	1	0	0	0
frication	0	1	0	0
other	0	0	1	0
vowel	0	0	0	1

This leads to the model

$$Y = \beta_1 X_1 + \beta_2 X_2 + \beta_3 X_3 + \beta_4 X_4 + \epsilon$$

(Note: you cannot use dummy coding for more than one additively combined categorical predictor, since you can only sacrifice the intercept once)


Dummy coding



Predictor	$\hat{\beta}$	$SE(\hat{\beta})$	$p(> t)$
burst	469.65	1.15	$\ll 0.001$
fricative	475.77	1.28	$\ll 0.001$
other	450.76	1.96	$\ll 0.001$
vowel	474.00	5.70	$\ll 0.001$

- Here, each parameter is just the mean RT for the corresponding initial phone type

Dummy coding



Predictor	$\hat{\beta}$	$SE(\hat{\beta})$	$p(> t)$
burst	469.65	1.15	$\ll 0.001$
fricative	475.77	1.28	$\ll 0.001$
other	450.76	1.96	$\ll 0.001$
vowel	474.00	5.70	$\ll 0.001$

- ▶ Here, each parameter is just the mean RT for the corresponding initial phone type
- ▶ The t statistics reflect the null hypothesis that mean RT for that initial phone type is 0 (not a very useful null hypothesis!)

Treatment coding

- ▶ With treatment coding, you leave the intercept in and arbitrarily label one value of the categorical predictor as the “baseline” level

Treatment coding

- ▶ With treatment coding, you leave the intercept in and arbitrarily label one value of the categorical predictor as the “baseline” level
- ▶ You then add a 0/1 predictor variable for each non-baseline level of the predictor

Treatment coding

- ▶ With treatment coding, you leave the intercept in and arbitrarily label one value of the categorical predictor as the “baseline” level
- ▶ You then add a 0/1 predictor variable for each non-baseline level of the predictor
- ▶ E.g., if we make **burst** the baseline level for initial phone:

Treatment coding

- ▶ With treatment coding, you leave the intercept in and arbitrarily label one value of the categorical predictor as the “baseline” level
- ▶ You then add a 0/1 predictor variable for each non-baseline level of the predictor
- ▶ E.g., if we make **burst** the baseline level for initial phone:

Treatment coding

- ▶ With treatment coding, you leave the intercept in and arbitrarily label one value of the categorical predictor as the “baseline” level
- ▶ You then add a 0/1 predictor variable for each non-baseline level of the predictor
- ▶ E.g., if we make **burst** the baseline level for initial phone:

Level of Init	X_1	X_2	X_3
burst	0	0	0
frication	1	0	0
other	0	1	0
vowel	0	0	1

Treatment coding


- ▶ With treatment coding, you leave the intercept in and arbitrarily label one value of the categorical predictor as the “baseline” level
- ▶ You then add a 0/1 predictor variable for each non-baseline level of the predictor
- ▶ E.g., if we make **burst** the baseline level for initial phone:

Level of Init	X_1	X_2	X_3
burst	0	0	0
frication	1	0	0
other	0	1	0
vowel	0	0	1

This leads to the model

$$Y = \alpha + \beta_1 X_1 + \beta_2 X_2 + \beta_3 X_3 + \epsilon$$


Treatment coding



Predictor	$\hat{\beta}$	$SE(\hat{\beta})$	$p(> t)$
Intercept	469.65	1.15	$\ll 0.001$
frication	6.12	1.72	0.000411
other	-18.90	2.27	$\ll 0.001$
vowel	4.35	5.82	0.455

- ▶ Here, weights 2–4 indicate the *difference* between the mean RT for words of the given phone type and for burst-initial words

Treatment coding



Predictor	$\hat{\beta}$	$SE(\hat{\beta})$	$p(> t)$
Intercept	469.65	1.15	$\ll 0.001$
fricative	6.12	1.72	0.000411
other	-18.90	2.27	$\ll 0.001$
vowel	4.35	5.82	0.455

- ▶ Here, weights 2–4 indicate the *difference* between the mean RT for words of the given phone type and for burst-initial words
- ▶ The t -statistic-based p -value indicates that fricative-initial words are slower than burst-initial words, and that words starting with other consonants are faster, but that we have insufficient information to conclude that vowel-initial words are either faster or slower

Sum (or deviation) coding

- ▶ Sum (or deviation) coding is designed to fit weights for all values of the categorical predictor, subject to the constraint that the weights sum to zero

Sum (or deviation) coding

- ▶ Sum (or deviation) coding is designed to fit weights for all values of the categorical predictor, subject to the constraint that the weights sum to zero
- ▶ The intercept remains in the model, and represents the hypothetical “average” response across all conditions

Sum (or deviation) coding

- ▶ Sum (or deviation) coding is designed to fit weights for all values of the categorical predictor, subject to the constraint that the weights sum to zero
- ▶ The intercept remains in the model, and represents the hypothetical “average” response across all conditions

Sum (or deviation) coding

- ▶ Sum (or deviation) coding is designed to fit weights for all values of the categorical predictor, subject to the constraint that the weights sum to zero
- ▶ The intercept remains in the model, and represents the hypothetical “average” response across all conditions

Level of Init	X_1	X_2	X_3
burst	1	0	0
frication	0	1	0
other	0	0	1
vowel	-1	-1	-1

- ▶ This leads to the model

$$Y = \alpha + \beta_1 X_1 + \beta_2 X_2 + \beta_3 X_3 + \epsilon$$

Sum (or deviation) coding

- ▶ Equivalently we can describe the sum-coding model as

Sum (or deviation) coding

- ▶ Equivalently we can describe the sum-coding model as

Sum (or deviation) coding

- ▶ Equivalently we can describe the sum-coding model as

Level of Init	X_1	X_2	X_3	X_4
burst	1	0	0	0
frication	0	1	0	0
other	0	0	1	0
vowel	0	0	0	1

- ▶ with the model

$$Y = \alpha + \beta_1 X_1 + \beta_2 X_2 + \beta_3 X_3 + \beta_4 X_4 + \epsilon$$


subject to the constraint that

$$\beta_1 + \beta_2 + \beta_3 + \beta_4 = 0$$

Or equivalently

$$\beta_4 = -\beta_1 - \beta_2 - \beta_3$$


Sum (or deviation) coding



Predictor	$\hat{\beta}$	$SE(\hat{\beta})$	$p(> t)$
Intercept	467.55	1.57	$\ll 0.001$
X_1	2.11	1.77	0.233
X_2	8.23	1.81	$\ll 0.001$
X_3	-16.79	2.09	$\ll 0.001$

- Parameters 2–4 are the *offsets* of the initial-phone-specific means from the hypothetical “grand average” (which is not the mean RT, because we have different # observations for different initial phones!)

Sum (or deviation) coding




Predictor	$\hat{\beta}$	$SE(\hat{\beta})$	$p(> t)$
Intercept	467.55	1.57	$\ll 0.001$
X_1	2.11	1.77	0.233
X_2	8.23	1.81	$\ll 0.001$
X_3	-16.79	2.09	$\ll 0.001$

- Parameters 2–4 are the *offsets* of the initial-phone-specific means from the hypothetical “grand average” (which is not the mean RT, because we have different # observations for different initial phones!)
- To recover the condition-specific offset $\hat{\beta}_4$ for and its standard error $SE(\hat{\beta}_4)$, recall that

$$\beta_4 = -\beta_1 - \beta_2 - \beta_3 \quad \Rightarrow \quad \hat{\beta}_4 = -\hat{\beta}_1 - \hat{\beta}_2 - \hat{\beta}_3$$

Sum (or deviation) coding




Predictor	$\hat{\beta}$	$SE(\hat{\beta})$	$p(> t)$
Intercept	467.55	1.57	$\ll 0.001$
X_1	2.11	1.77	0.233
X_2	8.23	1.81	$\ll 0.001$
X_3	-16.79	2.09	$\ll 0.001$

- Parameters 2–4 are the *offsets* of the initial-phone-specific means from the hypothetical “grand average” (which is not the mean RT, because we have different # observations for different initial phones!)
- To recover the condition-specific offset $\hat{\beta}_4$ for and its standard error $SE(\hat{\beta}_4)$, recall that

$$\beta_4 = -\beta_1 - \beta_2 - \beta_3 \quad \Rightarrow \quad \hat{\beta}_4 = -\hat{\beta}_1 - \hat{\beta}_2 - \hat{\beta}_3$$

Sum (or deviation) coding



Predictor	$\hat{\beta}$	$SE(\hat{\beta})$	$p(> t)$
Intercept	467.55	1.57	$\ll 0.001$
X_1	2.11	1.77	0.233
X_2	8.23	1.81	$\ll 0.001$
X_3	-16.79	2.09	$\ll 0.001$

- ▶ Parameters 2–4 are the *offsets* of the initial-phone-specific means from the hypothetical “grand average” (which is not the mean RT, because we have different # observations for different initial phones!)
- ▶ To recover the condition-specific offset $\hat{\beta}_4$ for and its standard error $SE(\hat{\beta}_4)$, recall that

$$\beta_4 = -\beta_1 - \beta_2 - \beta_3 \quad \Rightarrow \quad \hat{\beta}_4 = -\hat{\beta}_1 - \hat{\beta}_2 - \hat{\beta}_3$$

This gives us $\hat{\beta}_4 = -6.45$; using the formula for variance of the sum of random variables we can recover that $SE(\hat{\beta}_4) = 4.33$ and hence the p -value for that offset is 0.136

When to care about coding schema?

- ▶ To reiterate: the set of probability distributions $P(Y|X)$ that can be fitted does not depend on choice of coding schema (at least in this current type of case!)

When to care about coding schema?

- ▶ To reiterate: the set of probability distributions $P(Y|X)$ that can be fitted does not depend on choice of coding schema (at least in this current type of case!)
- ▶ So frequentist model comparisons involving the model (e.g., the F test, the likelihood ratio test) will give the same results regardless of the coding schema you choose

When to care about coding schema?

- ▶ To reiterate: the set of probability distributions $P(Y|X)$ that can be fitted does not depend on choice of coding schema (at least in this current type of case!)
- ▶ So frequentist model comparisons involving the model (e.g., the F test, the likelihood ratio test) will give the same results regardless of the coding schema you choose
 - ▶ E.g., in our case, comparing our model M_A against the “null” model $M_0 : RT = \alpha + \epsilon$ gives the result

$$\frac{(RSS_0 - RSS_A)/(m_A - m_0)}{RSS_A/(n - m_A - 1)} = \frac{(211685.57 - 173916.95)/3}{173916.95/535} = 38.73$$

When to care about coding schema?

- ▶ To reiterate: the set of probability distributions $P(Y|X)$ that can be fitted does not depend on choice of coding schema (at least in this current type of case!)
- ▶ So frequentist model comparisons involving the model (e.g., the F test, the likelihood ratio test) will give the same results regardless of the coding schema you choose
 - ▶ E.g., in our case, comparing our model M_A against the “null” model $M_0 : RT = \alpha + \epsilon$ gives the result

$$\frac{(RSS_0 - RSS_A)/(m_A - m_0)}{RSS_A/(n - m_A - 1)} = \frac{(211685.57 - 173916.95)/3}{173916.95/535} = 38.73$$

When to care about coding schema?

- ▶ To reiterate: the set of probability distributions $P(Y|X)$ that can be fitted does not depend on choice of coding schema (at least in this current type of case!)
- ▶ So frequentist model comparisons involving the model (e.g., the F test, the likelihood ratio test) will give the same results regardless of the coding schema you choose
 - ▶ E.g., in our case, comparing our model M_A against the “null” model $M_0 : RT = \alpha + \epsilon$ gives the result

$$\frac{(RSS_0 - RSS_A)/(m_A - m_0)}{RSS_A/(n - m_A - 1)} = \frac{(211685.57 - 173916.95)/3}{173916.95/535} = 38.73$$

- ▶ Consulting the cumulative distribution function for $F_{3,535}$, we find that this is highly significant— $p \ll 0.001$

When to care about coding schema?

- ▶ To reiterate: the set of probability distributions $P(Y|X)$ that can be fitted does not depend on choice of coding schema (at least in this current type of case!)
- ▶ So frequentist model comparisons involving the model (e.g., the F test, the likelihood ratio test) will give the same results regardless of the coding schema you choose
 - ▶ E.g., in our case, comparing our model M_A against the “null” model $M_0 : RT = \alpha + \epsilon$ gives the result

$$\frac{(RSS_0 - RSS_A)/(m_A - m_0)}{RSS_A/(n - m_A - 1)} = \frac{(211685.57 - 173916.95)/3}{173916.95/535} = 38.73$$

- ▶ Consulting the cumulative distribution function for $F_{3,535}$, we find that this is highly significant— $p \ll 0.001$
- ▶ Rather, your choice of contrasts should reflect how you want to *interpret* the parameters in your model!

Interactions between predictors

- ▶ Sometimes X_1 's effect on your response Y will differ depending on the value of another predictor X_2

Interactions between predictors

- ▶ Sometimes X_1 's effect on your response Y will differ depending on the value of another predictor X_2
- ▶ Example: back to age of acquisition and word frequency: let's make a **median split** on AoA and examine frequency effects on lexical decision time:



Interactions between predictors

- ▶ However, we want to generalize this so that we don't rely on the median split, but rather on a continuous dependence between the effects of frequency and AoA

Interactions between predictors

- ▶ However, we want to generalize this so that we don't rely on the median split, but rather on a continuous dependence between the effects of frequency and AoA
- ▶ This is known as an *interaction*

Interactions between predictors

- ▶ However, we want to generalize this so that we don't rely on the median split, but rather on a continuous dependence between the effects of frequency and AoA
- ▶ This is known as an *interaction*
- ▶ Formally, introducing an interaction between predictors X_i and X_j simply means adding a predictor whose value is the *product* of X_i and X_j

$$RT \sim \alpha + \beta_1 X_1 + \beta_2 X_2 + \beta_3 \overbrace{X_3}^{=X_1 X_2}$$

Interactions between predictors

- ▶ However, we want to generalize this so that we don't rely on the median split, but rather on a continuous dependence between the effects of frequency and AoA
- ▶ This is known as an *interaction*
- ▶ Formally, introducing an interaction between predictors X_i and X_j simply means adding a predictor whose value is the *product* of X_i and X_j

$$RT \sim \alpha + \beta_1 X_1 + \beta_2 X_2 + \beta_3 \overbrace{X_3}^{=X_1 X_2}$$

- ▶ Effect on the model matrix for our problem:

$$X = \begin{bmatrix} 1 & \text{Freq}_1 & \text{AoA}_1 & \text{Freq}_1 \text{AoA}_1 \\ 1 & \text{Freq}_2 & \text{AoA}_2 & \text{Freq}_2 \text{AoA}_2 \\ \vdots & \vdots & \vdots & \vdots \\ 1 & \text{Freq}_n & \text{AoA}_n & \text{Freq}_n \text{AoA}_n \end{bmatrix}$$

Interpreting interactions

Resulting model fit:

Predictor	$\hat{\beta}$	$SE(\hat{\beta})$	$p(> t)$
$\hat{\alpha}$	488.55	47.92	$\ll 0.001$
$\hat{\beta}_{\text{Freq}}$	6.07	5.73	0.29
$\hat{\beta}_{\text{AoA}}$	33.47	7.35	$\ll 0.001$
$\hat{\beta}_{\text{Freq} \times \text{AoA}}$	-2.84	0.912	0.00193

- ▶ The t statistic for $\hat{\beta}_{\text{Freq} \times \text{AoA}}$ indicates that the interaction is significant

Interpreting interactions

Resulting model fit:

Predictor	$\hat{\beta}$	$SE(\hat{\beta})$	$p(> t)$
$\hat{\alpha}$	488.55	47.92	$\ll 0.001$
$\hat{\beta}_{\text{Freq}}$	6.07	5.73	0.29
$\hat{\beta}_{\text{AoA}}$	33.47	7.35	$\ll 0.001$
$\hat{\beta}_{\text{Freq} \times \text{AoA}}$	-2.84	0.912	0.00193

- ▶ The t statistic for $\hat{\beta}_{\text{Freq} \times \text{AoA}}$ indicates that the interaction is significant
- ▶ Interpretation of the interaction parameter: as AoA increases, the effect of frequency becomes “less positive”

Interpreting interactions

Resulting model fit:

Predictor	$\hat{\beta}$	$SE(\hat{\beta})$	$p(> t)$
$\hat{\alpha}$	488.55	47.92	$\ll 0.001$
$\hat{\beta}_{\text{Freq}}$	6.07	5.73	0.29
$\hat{\beta}_{\text{AoA}}$	33.47	7.35	$\ll 0.001$
$\hat{\beta}_{\text{Freq} \times \text{AoA}}$	-2.84	0.912	0.00193

- ▶ The t statistic for $\hat{\beta}_{\text{Freq} \times \text{AoA}}$ indicates that the interaction is significant
- ▶ Interpretation of the interaction parameter: as AoA increases, the effect of frequency becomes “less positive”
- ▶ **Critically important issue:** when there is a higher-order interaction in the model, care must be taken with the interpretation of lower-order terms involved in the interaction

Interpreting interactions

Resulting model fit:

Predictor	$\hat{\beta}$	$SE(\hat{\beta})$	$p(> t)$
$\hat{\alpha}$	488.55	47.92	$\ll 0.001$
$\hat{\beta}_{\text{Freq}}$	6.07	5.73	0.29
$\hat{\beta}_{\text{AoA}}$	33.47	7.35	$\ll 0.001$
$\hat{\beta}_{\text{Freq} \times \text{AoA}}$	-2.84	0.912	0.00193

- ▶ The t statistic for $\hat{\beta}_{\text{Freq} \times \text{AoA}}$ indicates that the interaction is significant
- ▶ Interpretation of the interaction parameter: as AoA increases, the effect of frequency becomes “less positive”
- ▶ **Critically important issue:** when there is a higher-order interaction in the model, care must be taken with the interpretation of lower-order terms involved in the interaction
- ▶ Here, the “main” effect of frequency ($\hat{\beta}_{\text{Freq}}$) indicates the effect on RT of increasing word frequency **for a hypothetical word with age of acquisition at 0 years!**

Interpreting interactions

Resulting model fit:

Predictor	$\hat{\beta}$	$SE(\hat{\beta})$	$p(> t)$
$\hat{\alpha}$	488.55	47.92	$\ll 0.001$
$\hat{\beta}_{\text{Freq}}$	6.07	5.73	0.29
$\hat{\beta}_{\text{AoA}}$	33.47	7.35	$\ll 0.001$
$\hat{\beta}_{\text{Freq} \times \text{AoA}}$	-2.84	0.912	0.00193

- ▶ The t statistic for $\hat{\beta}_{\text{Freq} \times \text{AoA}}$ indicates that the interaction is significant
- ▶ Interpretation of the interaction parameter: as AoA increases, the effect of frequency becomes “less positive”
- ▶ **Critically important issue:** when there is a higher-order interaction in the model, care must be taken with the interpretation of lower-order terms involved in the interaction
- ▶ Here, the “main” effect of frequency ($\hat{\beta}_{\text{Freq}}$) indicates the effect on RT of increasing word frequency **for a hypothetical word with age of acquisition at 0 years!**
- ▶ But such words can't exist!

Interpreting interactions

- ▶ How do we interpret lower-order terms—e.g., “main effects” of X_1, X_2 —in the presence of an interaction term $X_{1 \times 2}$?


Interpreting interactions

- ▶ How do we interpret lower-order terms—e.g., “main effects” of X_1, X_2 —in the presence of an interaction term $X_{1 \times 2}$?
- ▶ One possible answer: we want a “main effect” of X_2 to indicate the effect of X_2 as X_1 remains constant at its average value for the dataset

Interpreting interactions


- ▶ How do we interpret lower-order terms—e.g., “main effects” of X_1, X_2 —in the presence of an interaction term $X_{1 \times 2}$?
- ▶ One possible answer: we want a “main effect” of X_2 to indicate the effect of X_2 as X_1 remains constant at its average value for the dataset
- ▶ We can achieve this by centering the predictor(s)

Interpreting interactions—centering predictors




Interpreting interactions—centering predictors

density(x = dat.english\$AoA)




density(x = dat.english\$cAoA)




Interpreting interactions—centering predictors


density(x = dat.english\$AoA)



density(x = dat.english\$cAoA)




density(x = dat.english\$WrittenFrequencyL)



Written frequency count (log base 2)

Interpreting interactions—centering predictors


density(x = dat.english\$AoA)



Age of acquisition --- years




density(x = dat.english\$cAoA)



Age of acquisition --- years (centered mean 0)


density(x = dat.english\$WrittenFrequencyL)



Written frequency count (log base 2)



density(x = dat.english\$cWrittenFrequencyL)



Written frequency count (log base 2, centered to mea)

Interpreting interactions

With the centered predictors, our model becomes:

Predictor	$\hat{\beta}$	$SE(\hat{\beta})$	$p(> t)$
$\hat{\alpha}$	602.04	1.94	$\ll 0.001$
$\hat{\beta}_{\text{cFreq}}$	-11.79	1.53	$\ll 0.001$
$\hat{\beta}_{\text{cAoA}}$	10.14	1.30	$\ll 0.001$
$\hat{\beta}_{\text{cFreq} \times \text{cAoA}}$	-2.84	0.912	0.00193

Interpreting interactions

With the centered predictors, our model becomes:

Predictor	$\hat{\beta}$	$SE(\hat{\beta})$	$p(> t)$
$\hat{\alpha}$	602.04	1.94	$\ll 0.001$
$\hat{\beta}_{\text{cFreq}}$	-11.79	1.53	$\ll 0.001$
$\hat{\beta}_{\text{cAoA}}$	10.14	1.30	$\ll 0.001$
$\hat{\beta}_{\text{cFreq} \times \text{cAoA}}$	-2.84	0.912	0.00193

- ▶ This shows that the “average”-AoA word in our dataset *does* have a substantial frequency effect

Interpreting interactions

With the centered predictors, our model becomes:

Predictor	$\hat{\beta}$	$SE(\hat{\beta})$	$p(> t)$
$\hat{\alpha}$	602.04	1.94	$\ll 0.001$
$\hat{\beta}_{\text{cFreq}}$	-11.79	1.53	$\ll 0.001$
$\hat{\beta}_{\text{cAoA}}$	10.14	1.30	$\ll 0.001$
$\hat{\beta}_{\text{cFreq} \times \text{cAoA}}$	-2.84	0.912	0.00193

- ▶ This shows that the “average”-AoA word in our dataset *does* have a substantial frequency effect
- ▶ Likewise, the “average”-frequency word in our dataset *does* have an AoA frequency effect

Interpreting interactions

- ▶ Even better, however, is *visualizing* the interactions

Interpreting interactions

- ▶ Even better, however, is *visualizing* the interactions
- ▶ The interaction means that as one predictor changes value, the other predictor's **effect** changes (and vice versa)

Interpreting interactions

- ▶ Even better, however, is *visualizing* the interactions
- ▶ The interaction means that as one predictor changes value, the other predictor's **effect** changes (and vice versa)
- ▶ Going back to the regression equation, we can write it as

$$\begin{aligned} RT &= \alpha + \beta_1 X_1 + \beta_2 X_2 + \beta_3 \overbrace{X_3}^{=X_1 X_2} + \epsilon \\ &= \alpha + \beta_1 X_1 + (\beta_2 + \beta_3 X_1) X_2 + \epsilon \end{aligned}$$

So

$$\hat{y} = \hat{\alpha} + \hat{\beta}_1 X_1 + (\hat{\beta}_2 + \hat{\beta}_3 X_1) X_2 + \epsilon$$

Interpreting interactions

- ▶ Even better, however, is *visualizing* the interactions
- ▶ The interaction means that as one predictor changes value, the other predictor's **effect** changes (and vice versa)
- ▶ Going back to the regression equation, we can write it as

$$\begin{aligned} RT &= \alpha + \beta_1 X_1 + \beta_2 X_2 + \beta_3 \overbrace{X_3}^{=X_1 X_2} + \epsilon \\ &= \alpha + \beta_1 X_1 + (\beta_2 + \beta_3 X_1) X_2 + \epsilon \end{aligned}$$

So

$$\hat{y} = \hat{\alpha} + \hat{\beta}_1 X_1 + (\hat{\beta}_2 + \hat{\beta}_3 X_1) X_2 + \epsilon$$

- ▶ So the estimated effect size of X_2 is $(\hat{\beta}_2 + \hat{\beta}_3 X_1)$

Interpreting interactions

- ▶ Even better, however, is *visualizing* the interactions
- ▶ The interaction means that as one predictor changes value, the other predictor's **effect** changes (and vice versa)
- ▶ Going back to the regression equation, we can write it as

$$\begin{aligned} RT &= \alpha + \beta_1 X_1 + \beta_2 X_2 + \beta_3 \overbrace{X_1 X_2}^{=X_1 X_2} + \epsilon \\ &= \alpha + \beta_1 X_1 + (\beta_2 + \beta_3 X_1) X_2 + \epsilon \end{aligned}$$

So

$$\hat{y} = \hat{\alpha} + \hat{\beta}_1 X_1 + (\hat{\beta}_2 + \hat{\beta}_3 X_1) X_2 + \epsilon$$

- ▶ So the estimated effect size of X_2 is $(\hat{\beta}_2 + \hat{\beta}_3 X_1)$
- ▶ Because the estimates $\{\hat{\beta}_i\}$ are multivariate-normal distributed, we can also use this to compute a standard error (and thus confidence intervals) of the effect size of X_2 **for a given value of X_1**

Visualizing interactions

- Let us explore the effect of a word's frequency as a function of its AoA in this model



Visualizing interactions

- Let us explore the effect of a word's frequency as a function of its AoA in this model



- What are some possible theoretical interpretations of this

Interactions with categorical variables

- ▶ Case of an interaction between a k -level categorical variable X_1 and a continuous variable X_2

Interactions with categorical variables

- ▶ Case of an interaction between a k -level categorical variable X_1 and a continuous variable X_2
- ▶ Review: how many parameters does an additive model of $Y \sim X_1 + X_2$ have?

Interactions with categorical variables

- ▶ Case of an interaction between a k -level categorical variable X_1 and a continuous variable X_2
- ▶ Review: how many parameters does an additive model of $Y \sim X_1 + X_2$ have?
 - ▶ 1 for the intercept

Interactions with categorical variables

- ▶ Case of an interaction between a k -level categorical variable X_1 and a continuous variable X_2
- ▶ Review: how many parameters does an additive model of $Y \sim X_1 + X_2$ have?
 - ▶ 1 for the intercept
 - ▶ $k - 1$ more for X_1

Interactions with categorical variables

- ▶ Case of an interaction between a k -level categorical variable X_1 and a continuous variable X_2
- ▶ Review: how many parameters does an additive model of $Y \sim X_1 + X_2$ have?
 - ▶ 1 for the intercept
 - ▶ $k - 1$ more for X_1
 - ▶ 1 more for X_2

Interactions with categorical variables

- ▶ Case of an interaction between a k -level categorical variable X_1 and a continuous variable X_2
- ▶ Review: how many parameters does an additive model of $Y \sim X_1 + X_2$ have?
 - ▶ 1 for the intercept
 - ▶ $k - 1$ more for X_1
 - ▶ 1 more for X_2
- ▶ So, a total of $k + 1$

Interactions with categorical variables

- ▶ Case of an interaction between a k -level categorical variable X_1 and a continuous variable X_2
- ▶ Review: how many parameters does an additive model of $Y \sim X_1 + X_2$ have?
 - ▶ 1 for the intercept
 - ▶ $k - 1$ more for X_1
 - ▶ 1 more for X_2
- ▶ So, a total of $k + 1$
- ▶ Adding the interaction between X_1 and X_2 allows for a separate X_2 slope for each value of X_1 (instead of the same slope for all values of X_1)

Interactions with categorical variables

- ▶ Case of an interaction between a k -level categorical variable X_1 and a continuous variable X_2
- ▶ Review: how many parameters does an additive model of $Y \sim X_1 + X_2$ have?
 - ▶ 1 for the intercept
 - ▶ $k - 1$ more for X_1
 - ▶ 1 more for X_2
- ▶ So, a total of $k + 1$
- ▶ Adding the interaction between X_1 and X_2 allows for a separate X_2 slope for each value of X_1 (instead of the same slope for all values of X_1)
- ▶ Hence this adds a total **a total of $k - 1$ new parameters** beyond the additive (“main effects”) model

Interactions with categorical variables


Example: different effects of AoA on naming time for old versus young native English speakers:

Predictor	$\hat{\beta}$	$SE(\hat{\beta})$	$p(> t)$
$\hat{\beta}_{\text{Age=old}}$	623.81	4.81	$\ll 0.001$
$\hat{\beta}_{\text{Age=young}}$	452.88	4.81	$\ll 0.001$
$\hat{\beta}_{\text{AoA:Age=old}}$	5.52	0.744	$\ll 0.001$
$\hat{\beta}_{\text{AoA:Age=young}}$	2.57	0.744	0.000581

Interactions with categorical variables

Example: different effects of AoA on naming time for old versus young native English speakers:


Predictor	$\hat{\beta}$	$SE(\hat{\beta})$	$p(> t)$
$\hat{\beta}_{\text{Age=old}}$	623.81	4.81	$\ll 0.001$
$\hat{\beta}_{\text{Age=young}}$	452.88	4.81	$\ll 0.001$
$\hat{\beta}_{\text{AoA:Age=old}}$	5.52	0.744	$\ll 0.001$
$\hat{\beta}_{\text{AoA:Age=young}}$	2.57	0.744	0.000581



Interactions with categorical variables

Example: different effects of AoA on naming time for old versus young native English speakers:

Predictor	$\hat{\beta}$	$SE(\hat{\beta})$	$p(> t)$
$\hat{\beta}_{\text{Age=old}}$	623.81	4.81	$\ll 0.001$
$\hat{\beta}_{\text{Age=young}}$	452.88	4.81	$\ll 0.001$
$\hat{\beta}_{\text{AoA:Age=old}}$	5.52	0.744	$\ll 0.001$
$\hat{\beta}_{\text{AoA:Age=young}}$	2.57	0.744	0.000581




- ▶ With this parameterization, the t statistics aren't useful in assessing the strength of evidence for the interaction

Interactions with categorical variables

Example: different effects of AoA on naming time for old versus young native English speakers:

Predictor	$\hat{\beta}$	$SE(\hat{\beta})$	$p(> t)$
$\hat{\beta}_{\text{Age=old}}$	623.81	4.81	$\ll 0.001$
$\hat{\beta}_{\text{Age=young}}$	452.88	4.81	$\ll 0.001$
$\hat{\beta}_{\text{AoA:Age=old}}$	5.52	0.744	$\ll 0.001$
$\hat{\beta}_{\text{AoA:Age=young}}$	2.57	0.744	0.000581



- ▶ With this parameterization, the t statistics aren't useful in assessing the strength of evidence for the interaction
- ▶ We can use the F -test to do this instead—comparing with a 3-parameter model (sans interaction) gives us $F_{1,1074} = 7.84$, for a p -value of 0.00519

Interactions between categorical variables

- ▶ Case of an interaction between a k_1 -level categorical variable X_1 and another k_2 -level categorical variable X_2 , then you have a total of $k_1 k_2$ parameters

Interactions between categorical variables

- ▶ Case of an interaction between a k_1 -level categorical variable X_1 and another k_2 -level categorical variable X_2 , then you have a total of $k_1 k_2$ parameters
- ▶ Review: how many parameters does an additive model of $Y \sim X_1 + X_2$ have?

Interactions between categorical variables

- ▶ Case of an interaction between a k_1 -level categorical variable X_1 and another k_2 -level categorical variable X_2 , then you have a total of $k_1 k_2$ parameters
- ▶ Review: how many parameters does an additive model of $Y \sim X_1 + X_2$ have?
 - ▶ 1 for the intercept

Interactions between categorical variables

- ▶ Case of an interaction between a k_1 -level categorical variable X_1 and another k_2 -level categorical variable X_2 , then you have a total of $k_1 k_2$ parameters
- ▶ Review: how many parameters does an additive model of $Y \sim X_1 + X_2$ have?
 - ▶ 1 for the intercept
 - ▶ $k_1 - 1$ more for X_1

Interactions between categorical variables

- ▶ Case of an interaction between a k_1 -level categorical variable X_1 and another k_2 -level categorical variable X_2 , then you have a total of $k_1 k_2$ parameters
- ▶ Review: how many parameters does an additive model of $Y \sim X_1 + X_2$ have?
 - ▶ 1 for the intercept
 - ▶ $k_1 - 1$ more for X_1
 - ▶ $k_2 - 1$ more for X_2

Interactions between categorical variables

- ▶ Case of an interaction between a k_1 -level categorical variable X_1 and another k_2 -level categorical variable X_2 , then you have a total of $k_1 k_2$ parameters
- ▶ Review: how many parameters does an additive model of $Y \sim X_1 + X_2$ have?
 - ▶ 1 for the intercept
 - ▶ $k_1 - 1$ more for X_1
 - ▶ $k_2 - 1$ more for X_2
- ▶ So, a total of $k_1 + k_2 - 1$

Interactions between categorical variables

- ▶ Case of an interaction between a k_1 -level categorical variable X_1 and another k_2 -level categorical variable X_2 , then you have a total of $k_1 k_2$ parameters
- ▶ Review: how many parameters does an additive model of $Y \sim X_1 + X_2$ have?
 - ▶ 1 for the intercept
 - ▶ $k_1 - 1$ more for X_1
 - ▶ $k_2 - 1$ more for X_2
- ▶ So, a total of $k_1 + k_2 - 1$
- ▶ Adding the interaction between X_1 and X_2 means that there is a separate effect of each logically possible combination of the two predictors

Interactions between categorical variables

- ▶ Case of an interaction between a k_1 -level categorical variable X_1 and another k_2 -level categorical variable X_2 , then you have a total of $k_1 k_2$ parameters
- ▶ Review: how many parameters does an additive model of $Y \sim X_1 + X_2$ have?
 - ▶ 1 for the intercept
 - ▶ $k_1 - 1$ more for X_1
 - ▶ $k_2 - 1$ more for X_2
- ▶ So, a total of $k_1 + k_2 - 1$
- ▶ Adding the interaction between X_1 and X_2 means that there is a separate effect of each logically possible combination of the two predictors
- ▶ So there is a total of $k_1 k_2$ parameters (one for the intercept and $k_1 k_2 - 1$ for the overall interaction)

Interactions between categorical variables

- ▶ Case of an interaction between a k_1 -level categorical variable X_1 and another k_2 -level categorical variable X_2 , then you have a total of $k_1 k_2$ parameters
- ▶ Review: how many parameters does an additive model of $Y \sim X_1 + X_2$ have?
 - ▶ 1 for the intercept
 - ▶ $k_1 - 1$ more for X_1
 - ▶ $k_2 - 1$ more for X_2
- ▶ So, a total of $k_1 + k_2 - 1$
- ▶ Adding the interaction between X_1 and X_2 means that there is a separate effect of each logically possible combination of the two predictors
- ▶ So there is a total of $k_1 k_2$ parameters (one for the intercept and $k_1 k_2 - 1$ for the overall interaction)
- ▶ Hence this adds a total a total of $k_1 k_2 - k_1 - k_2 + 1$ new parameters beyond the additive (“main effects”) model

Interactions between categorical variables

Example: the relationship between initial phone type and native speaker age on naming times

Predictor	$\hat{\beta}$	$SE(\hat{\beta})$	$p(> t)$
$\hat{\beta}_{\text{Age=old,Init=burst}}$	654.83	1.66	$\ll 0.001$
$\hat{\beta}_{\text{Age=young,Init=burst}}$	469.65	1.66	$\ll 0.001$
$\hat{\beta}_{\text{Age=old,Init=frication}}$	669.72	1.84	$\ll 0.001$
$\hat{\beta}_{\text{Age=young,Init=frication}}$	475.77	1.84	$\ll 0.001$
$\hat{\beta}_{\text{Age=old,Init=other}}$	643.18	2.82	$\ll 0.001$
$\hat{\beta}_{\text{Age=young,Init=other}}$	450.76	2.82	$\ll 0.001$
$\hat{\beta}_{\text{Age=old,Init=vowel}}$	653.65	8.21	$\ll 0.001$
$\hat{\beta}_{\text{Age=young,Init=vowel}}$	474.00	8.21	$\ll 0.001$

- ▶ With this parameterization the model is just encoding eight different means

Interactions between categorical variables

Example: the relationship between initial phone type and native speaker age on naming times

Predictor	$\hat{\beta}$	$SE(\hat{\beta})$	$p(> t)$
$\hat{\beta}_{\text{Age=old,Init=burst}}$	654.83	1.66	$\ll 0.001$
$\hat{\beta}_{\text{Age=young,Init=burst}}$	469.65	1.66	$\ll 0.001$
$\hat{\beta}_{\text{Age=old,Init=frication}}$	669.72	1.84	$\ll 0.001$
$\hat{\beta}_{\text{Age=young,Init=frication}}$	475.77	1.84	$\ll 0.001$
$\hat{\beta}_{\text{Age=old,Init=other}}$	643.18	2.82	$\ll 0.001$
$\hat{\beta}_{\text{Age=young,Init=other}}$	450.76	2.82	$\ll 0.001$
$\hat{\beta}_{\text{Age=old,Init=vowel}}$	653.65	8.21	$\ll 0.001$
$\hat{\beta}_{\text{Age=young,Init=vowel}}$	474.00	8.21	$\ll 0.001$

- ▶ With this parameterization the model is just encoding eight different means
- ▶ Once again, we can use the F test to compare this model with an additive model of parameters, giving us

Interactions between categorical variables

Example: the relationship between initial phone type and native speaker age on naming times

Predictor	$\hat{\beta}$	$SE(\hat{\beta})$	$p(> t)$
$\hat{\beta}_{\text{Age=old,Init=burst}}$	654.83	1.66	$\ll 0.001$
$\hat{\beta}_{\text{Age=young,Init=burst}}$	469.65	1.66	$\ll 0.001$
$\hat{\beta}_{\text{Age=old,Init=frication}}$	669.72	1.84	$\ll 0.001$
$\hat{\beta}_{\text{Age=young,Init=frication}}$	475.77	1.84	$\ll 0.001$
$\hat{\beta}_{\text{Age=old,Init=other}}$	643.18	2.82	$\ll 0.001$
$\hat{\beta}_{\text{Age=young,Init=other}}$	450.76	2.82	$\ll 0.001$
$\hat{\beta}_{\text{Age=old,Init=vowel}}$	653.65	8.21	$\ll 0.001$
$\hat{\beta}_{\text{Age=young,Init=vowel}}$	474.00	8.21	$\ll 0.001$

- ▶ With this parameterization the model is just encoding eight different means
- ▶ Once again, we can use the F test to compare this model with an additive model of 5 parameters, giving us

Interactions between categorical variables

Example: the relationship between initial phone type and native speaker age on naming times

Predictor	$\hat{\beta}$	$SE(\hat{\beta})$	$p(> t)$
$\hat{\beta}_{\text{Age=old,Init=burst}}$	654.83	1.66	$\ll 0.001$
$\hat{\beta}_{\text{Age=young,Init=burst}}$	469.65	1.66	$\ll 0.001$
$\hat{\beta}_{\text{Age=old,Init=frication}}$	669.72	1.84	$\ll 0.001$
$\hat{\beta}_{\text{Age=young,Init=frication}}$	475.77	1.84	$\ll 0.001$
$\hat{\beta}_{\text{Age=old,Init=other}}$	643.18	2.82	$\ll 0.001$
$\hat{\beta}_{\text{Age=young,Init=other}}$	450.76	2.82	$\ll 0.001$
$\hat{\beta}_{\text{Age=old,Init=vowel}}$	653.65	8.21	$\ll 0.001$
$\hat{\beta}_{\text{Age=young,Init=vowel}}$	474.00	8.21	$\ll 0.001$

- ▶ With this parameterization the model is just encoding eight different means
- ▶ Once again, we can use the F test to compare this model with an additive model of 5 parameters, giving us

Interactions between categorical variables

Example: the relationship between initial phone type and native speaker age on naming times

Predictor	$\hat{\beta}$	$SE(\hat{\beta})$	$p(> t)$
$\hat{\beta}_{\text{Age=old,Init=burst}}$	654.83	1.66	$\ll 0.001$
$\hat{\beta}_{\text{Age=young,Init=burst}}$	469.65	1.66	$\ll 0.001$
$\hat{\beta}_{\text{Age=old,Init=frication}}$	669.72	1.84	$\ll 0.001$
$\hat{\beta}_{\text{Age=young,Init=frication}}$	475.77	1.84	$\ll 0.001$
$\hat{\beta}_{\text{Age=old,Init=other}}$	643.18	2.82	$\ll 0.001$
$\hat{\beta}_{\text{Age=young,Init=other}}$	450.76	2.82	$\ll 0.001$
$\hat{\beta}_{\text{Age=old,Init=vowel}}$	653.65	8.21	$\ll 0.001$
$\hat{\beta}_{\text{Age=young,Init=vowel}}$	474.00	8.21	$\ll 0.001$

- ▶ With this parameterization the model is just encoding eight different means
- ▶ Once again, we can use the F test to compare this model with an additive model of 5 parameters, giving us $F_{3,1070} = 2.52$, for a p -value of 0.0563

General words of wisdom for categorical factors & interactions

- ▶ There are many different ways to parameterize any model

General words of wisdom for categorical factors & interactions

- ▶ There are many different ways to parameterize any model
- ▶ It's easy to get lost in the apparent cleverness of setting model contrasts for interesting parameterizations

General words of wisdom for categorical factors & interactions

- ▶ There are many different ways to parameterize any model
- ▶ It's easy to get lost in the apparent cleverness of setting model contrasts for interesting parameterizations
- ▶ Don't lose sight of the more important issue: **model class**

General words of wisdom for categorical factors & interactions

- ▶ There are many different ways to parameterize any model
- ▶ It's easy to get lost in the apparent cleverness of setting model contrasts for interesting parameterizations
- ▶ Don't lose sight of the more important issue: **model class**
- ▶ Because we're operating under maximum likelihood, **if one model is a reparameterization of another, they make the same predictions and yield the same inferences for model comparison!**

General words of wisdom for categorical factors & interactions

- ▶ There are many different ways to parameterize any model
- ▶ It's easy to get lost in the apparent cleverness of setting model contrasts for interesting parameterizations
- ▶ Don't lose sight of the more important issue: **model class**
- ▶ Because we're operating under maximum likelihood, **if one model is a reparameterization of another, they make the same predictions and yield the same inferences for model comparison!**
- ▶ Model comparisons (for simple linear regression, the F test; more generally, the likelihood ratio test) are your friend!

Dichotomous categorical response variables

- ▶ We have generalized linear regression to categorical *predictor* variables

Dichotomous categorical response variables

- ▶ We have generalized linear regression to categorical *predictor* variables
- ▶ However, we have not yet addressed the case when the *response* variable is categorical

Dichotomous categorical response variables

- ▶ We have generalized linear regression to categorical *predictor* variables
- ▶ However, we have not yet addressed the case when the *response* variable is categorical
- ▶ Let's consider the case of a dichotomous response variable

Dichotomous categorical response variables

- ▶ We have generalized linear regression to categorical *predictor* variables
- ▶ However, we have not yet addressed the case when the *response* variable is categorical
- ▶ Let's consider the case of a dichotomous response variable
- ▶ Example: the dative alternation (Bresnan et al., 2007)

Sally sent [the children]_{Recip} [toys]_{Theme}

Sally sent [toys]_{Theme} to [the children]_{Recip}

Double Object

Prepositional Object

Dichotomous categorical response variables

- ▶ We have generalized linear regression to categorical *predictor* variables
- ▶ However, we have not yet addressed the case when the *response* variable is categorical
- ▶ Let's consider the case of a dichotomous response variable
- ▶ Example: the dative alternation (Bresnan et al., 2007)

Sally sent [the children]_{Recip} [toys]_{Theme}


Sally sent [toys]_{Theme} to [the children]_{Recip}

Double Object

Prepositional Object

- ▶ We looked briefly before at the effects of definiteness of the **theme** (*toys/the toys*) and **recipient** (*children/the children*)


Dichotomous categorical responses



- ▶ We could learn these four separate means, but we would fail to capture the systematicity of the effects seen here

Dichotomous categorical responses

Another way of representing the four means:



- ▶ This looks like what we called an *additive pattern* for linear regression

$$Y = \alpha + \beta_1 X_1 + \beta_2 X_2 + \epsilon$$

where X_1 is 1 iff the theme is indefinite, and X_2 is 1 iff the recipient is indefinite (both 0 otherwise)

Problems for linear models with categorical response

$$Y = \alpha + \beta_1 X_1 + \beta_2 X_2 + \epsilon \quad X_1 = 1 \text{ iff theme indefinite, } X_2 = 1 \text{ iff recipient indefinite (both 0 otherwise)}$$

1. **Bad predictions for individual observations:** in linear regression, the noise term ϵ is *Gaussian* (normally distributed)—it predicts that any continuous value is possible

Problems for linear models with categorical response

$$Y = \alpha + \beta_1 X_1 + \beta_2 X_2 + \epsilon \quad X_1 = 1 \text{ iff theme indefinite, } X_2 = 1 \text{ iff recipient indefinite (both 0 otherwise)}$$

1. **Bad predictions for individual observations:** in linear regression, the noise term ϵ is *Gaussian* (normally distributed)—it predicts that any continuous value is possible
 - ▶ The only really possible outcomes for individual observations are 0 (PP recipient) and 1 (NP recipient)

Problems for linear models with categorical response

$$Y = \alpha + \beta_1 X_1 + \beta_2 X_2 + \epsilon \quad X_1 = 1 \text{ iff theme indefinite, } X_2 = 1 \text{ iff recipient indefinite (both 0 otherwise)}$$

1. **Bad predictions for individual observations:** in linear regression, the noise term ϵ is *Gaussian* (normally distributed)—it predicts that any continuous value is possible
 - ▶ The only really possible outcomes for individual observations are 0 (PP recipient) and 1 (NP recipient)
 - ▶ Remember that our observed “means” are averages of many 0 and 1 observations!

## Definiteness of recipient and theme			
## Realization	Rec=def, Theme=def	Rec=def, Theme=indef	
## Double obj.	19	78	
## Prep. obj.	34	23	
## Definiteness of recipient and theme			
## Realization	Rec=indef, Theme=def	Rec=indef, Theme=indef	
## Double obj.	0	4	
## Prep. obj.	5	9	

Problems for linear models with categorical response

$$Y = \alpha + \beta_1 X_1 + \beta_2 X_2 + \epsilon \quad X_1 = 1 \text{ iff theme indefinite, } X_2 = 1 \text{ iff recipient indefinite (both 0 otherwise)}$$

1. **Bad predictions for individual observations:** in linear regression, the noise term ϵ is *Gaussian* (normally distributed)—it predicts that any continuous value is possible
 - ▶ The only really possible outcomes for individual observations are 0 (PP recipient) and 1 (NP recipient)
 - ▶ Remember that our observed “means” are averages of many 0 and 1 observations!

##	Definiteness of recipient and theme		
## Realization	Rec=def, Theme=def	Rec=def, Theme=indef	
## Double obj.	19	78	
## Prep. obj.	34	23	
##	Definiteness of recipient and theme		
## Realization	Rec=indef, Theme=def	Rec=indef, Theme=indef	
## Double obj.	0	4	
## Prep. obj.	5	9	

2. **Bad predicted means:** in linear regression, there is no guarantee that the predicted mean response \hat{y} will fall between 0 and 1, even if all individual observations fall within this range

Bad predicted means with linear regression for categorical response

- Consider a case where a predictor is continuous and the response is categorical:

Recipient is NP

Mary sent **John** a shiny toy

Mary sent **her friend** a shiny toy

Mary sent **every kid in the room** a shiny toy

Recipient is PP

Mary sent a shiny toy to **John**

Mary sent a shiny toy to **her friend**

Mary sent a shiny toy to **every kid in the room**

Bad predicted means with linear regression for categorical response

- ▶ Consider a case where a predictor is continuous and the response is categorical:

Recipient is NP

Mary sent **John** a shiny toy

Mary sent **her friend** a shiny toy

Mary sent **every kid in the room** a shiny toy

Recipient is PP

Mary sent a shiny toy to **John**


Mary sent a shiny toy to **her friend**

Mary sent a shiny toy to **every kid in the room**

- ▶ We could quantify the *size* of the recipient in any number of ways (here we'll use length in # of words)


Dichotomous categorical response variables

Here's what happens when we learn a linear regression model on recipient length:



Bad predicted means with linear regression for categorical response


Same problem if we use *log* of recipient length as a predictor:



Bad predicted means with linear regression for categorical response

Even spline-based methods (which we haven't covered yet) give us the same problem, too, at the far end of the range of lengths:

Realization of Recipient (NP=1, PP=0)



Problems with linear regression for categorical response

- ▶ So linear regression is bad for categorical response variables in:

Problems with linear regression for categorical response

- ▶ So linear regression is bad for categorical response variables in:
 1. The **noise distribution** it assumes around the predicted mean

Problems with linear regression for categorical response

- ▶ So linear regression is bad for categorical response variables in:
 1. The **noise distribution** it assumes around the predicted mean
 2. The **range of the predicted mean** allowed

Problems with linear regression for categorical response

- ▶ So linear regression is bad for categorical response variables in:
 1. The **noise distribution** it assumes around the predicted mean
 2. The **range of the predicted mean** allowed
- ▶ Fortunately, the framework of generalized linear models (GLMs) gives us the flexibility to deal with these problems!

Reviewing GLMs

Assumptions of the generalized linear model (GLM):

1. Predictors $\{X_i\}$ influence Y through the mediation of a **linear predictor** η ;

Reviewing GLMs

Assumptions of the generalized linear model (GLM):

1. Predictors $\{X_i\}$ influence Y through the mediation of a **linear predictor** η ;
2. η is a linear combination of the $\{X_i\}$:

Reviewing GLMs

Assumptions of the generalized linear model (GLM):

1. Predictors $\{X_i\}$ influence Y through the mediation of a **linear predictor** η ;
2. η is a linear combination of the $\{X_i\}$:

$$\eta = \alpha + \beta_1 X_1 + \cdots + \beta_m X_m \quad (\text{linear predictor})$$

Reviewing GLMs

Assumptions of the generalized linear model (GLM):

1. Predictors $\{X_i\}$ influence Y through the mediation of a **linear predictor** η ;
2. η is a linear combination of the $\{X_i\}$:

$$\eta = \alpha + \beta_1 X_1 + \cdots + \beta_m X_m \quad (\text{linear predictor})$$

3. η determines the predicted mean μ of Y

$$\eta = l(\mu) \quad (\text{link function})$$

Reviewing GLMs

Assumptions of the generalized linear model (GLM):

1. Predictors $\{X_i\}$ influence Y through the mediation of a **linear predictor** η ;
2. η is a linear combination of the $\{X_i\}$:

$$\eta = \alpha + \beta_1 X_1 + \cdots + \beta_m X_m \quad (\text{linear predictor})$$

3. η determines the predicted mean μ of Y

$$\eta = l(\mu) \quad (\text{link function})$$

4. There is some **noise distribution** of Y around the predicted mean μ of Y :

$$P(Y = y; \mu)$$

Logit GLMs for dichotomous responses

- ▶ Choosing a different link function and noise distribution gives us the **logit model**

Logit GLMs for dichotomous responses

- ▶ Choosing a different link function and noise distribution gives us the **logit model**
- ▶ Logit link function:

$$\eta = \log \frac{\mu}{1 - \mu}$$

Logit GLMs for dichotomous responses

- ▶ Choosing a different link function and noise distribution gives us the **logit model**
- ▶ Logit link function:

$$\eta = \log \frac{\mu}{1 - \mu}$$

- ▶ Bernoulli noise distribution around predicted mean μ :

$$P(Y = y|\mu) = \begin{cases} \mu & y = 1 \\ 1 - \mu & y = 0 \\ 0 & \text{otherwise} \end{cases}$$

Logit GLMs for dichotomous responses

- ▶ Choosing a different link function and noise distribution gives us the **logit model**
- ▶ Logit link function:

$$\eta = \log \frac{\mu}{1 - \mu}$$

- ▶ Bernoulli noise distribution around predicted mean μ :

$$P(Y = y|\mu) = \begin{cases} \mu & y = 1 \\ 1 - \mu & y = 0 \\ 0 & \text{otherwise} \end{cases}$$

- ▶ The linear predictor remains as it was before:

$$\eta = \alpha + \beta_1 X_1 + \cdots + \beta_m X_m$$

Logit GLMs for dichotomous responses

- ▶ Choosing a different link function and noise distribution gives us the **logit model**
- ▶ Logit link function:

$$\eta = \log \frac{\mu}{1 - \mu}$$

- ▶ Bernoulli noise distribution around predicted mean μ :

$$P(Y = y|\mu) = \begin{cases} \mu & y = 1 \\ 1 - \mu & y = 0 \\ 0 & \text{otherwise} \end{cases}$$

- ▶ The linear predictor remains as it was before:

$$\eta = \alpha + \beta_1 X_1 + \cdots + \beta_m X_m$$

- ▶ Using logit GLMs to fit data with dichotomous response variables is called **logistic regression**

The logit link function

- ▶ A lot of the action in logistic regression is in the logit link function:

$$\eta = \log \frac{\mu}{1 - \mu}$$

The logit link function

- ▶ A lot of the action in logistic regression is in the logit link function:

$$\eta = \log \frac{\mu}{1 - \mu}$$

- ▶ Looking at its *inverse* is equally useful:

$$\mu = \frac{e^{\eta}}{1 + e^{\eta}}$$

The logit link function

- ▶ A lot of the action in logistic regression is in the logit link function:

$$\eta = \log \frac{\mu}{1 - \mu}$$

- ▶ Looking at its *inverse* is equally useful:

$$\mu = \frac{e^{\eta}}{1 + e^{\eta}}$$


The logit link function

- ▶ A lot of the action in logistic regression is in the logit link function:

$$\eta = \log \frac{\mu}{1 - \mu}$$

- ▶ Looking at its *inverse* is equally useful:

$$\mu = \frac{e^{\eta}}{1 + e^{\eta}}$$



Estimating parameters in logistic regression

$$\eta = \alpha + \beta_1 X_1 + \cdots + \beta_m X_m$$

$$\mu = \frac{e^\eta}{1 + e^\eta}$$

$$P(Y = y|\mu) = \begin{cases} \mu & y = 1 \\ 1 - \mu & y = 0 \\ 0 & \text{otherwise} \end{cases}$$

- ▶ As with linear regression, the regression weights $\langle \alpha, \beta_1, \dots, \beta_m \rangle$ must be learned

Estimating parameters in logistic regression

$$\eta = \alpha + \beta_1 X_1 + \cdots + \beta_m X_m$$

$$\mu = \frac{e^\eta}{1 + e^\eta}$$

$$P(Y = y|\mu) = \begin{cases} \mu & y = 1 \\ 1 - \mu & y = 0 \\ 0 & \text{otherwise} \end{cases}$$

- ▶ As with linear regression, the regression weights $\langle \alpha, \beta_1, \dots, \beta_m \rangle$ must be learned
- ▶ Unlike linear regression, there is no additional noise parameter to be learned (σ^2 in linear regression)

Estimating parameters in logistic regression

$$\eta = \alpha + \beta_1 X_1 + \cdots + \beta_m X_m$$

$$\mu = \frac{e^\eta}{1 + e^\eta}$$

$$P(Y = y|\mu) = \begin{cases} \mu & y = 1 \\ 1 - \mu & y = 0 \\ 0 & \text{otherwise} \end{cases}$$

- ▶ As with linear regression, the regression weights $\langle \alpha, \beta_1, \dots, \beta_m \rangle$ must be learned
- ▶ Unlike linear regression, there is no additional noise parameter to be learned (σ^2 in linear regression)
- ▶ Once again, we can use the **method of maximum likelihood** (which we use in the example here) or **Bayesian inference** to estimate parameters

Interpreting an additive logistic regression model

- ▶ Here's a logistic regression model for additive effects of theme and recipient definiteness:

$$\eta = \alpha + \beta_{\text{ThemeDef}} X_{\text{ThemeDef}} + \beta_{\text{RecDef}} X_{\text{RecDef}}$$

$$\mu = \frac{e^{\eta}}{1 + e^{\eta}}$$

$$P(Y = y | \mu) = \begin{cases} \mu & y = 1 \\ 1 - \mu & y = 0 \\ 0 & \text{otherwise} \end{cases}$$

Interpreting an additive logistic regression model

- ▶ Here's a logistic regression model for additive effects of theme and recipient definiteness:

$$\eta = \alpha + \beta_{\text{ThemeDef}} X_{\text{ThemeDef}} + \beta_{\text{RecDef}} X_{\text{RecDef}}$$

$$\mu = \frac{e^{\eta}}{1 + e^{\eta}}$$


$$P(Y = y|\mu) = \begin{cases} \mu & y = 1 \\ 1 - \mu & y = 0 \\ 0 & \text{otherwise} \end{cases}$$

- ▶ The maximum likelihood estimate for the three regression parameters is

$$\begin{array}{ll} \hat{\alpha} & -0.61 \\ \hat{\beta}_{\text{ThemeDef}} & 1.8 \\ \hat{\beta}_{\text{RecDef}} & -2.2 \end{array}$$


Interpreting an additive logistic regression model

$\hat{\alpha}$	-0.61
$\hat{\beta}_{\text{ThemeDef}}$	1.8
$\hat{\beta}_{\text{RecDef}}$	-2.2



Interpreting an additive logistic regression model

This additive model does a decent job of modeling the true means!



Confidence regions for logistic regression

- ▶ Let us write the linear-predictor part of our GLM in matrix form:

$$\eta = X\beta$$

Confidence regions for logistic regression

- ▶ Let us write the linear-predictor part of our GLM in matrix form:

$$\boldsymbol{\eta} = \mathbf{X}\boldsymbol{\beta}$$

- ▶ In linear regression, we built confidence regions for parameter estimates on the basis that the covariance matrix of the MLE $\hat{\boldsymbol{\beta}}$ can be written *exactly* as

$$\text{Cov}(\hat{\boldsymbol{\beta}}) = \sigma^2(\mathbf{X}^T\mathbf{X})^{-1}$$

Confidence regions for logistic regression

- ▶ Let us write the linear-predictor part of our GLM in matrix form:

$$\eta = X\beta$$

- ▶ In linear regression, we built confidence regions for parameter estimates on the basis that the covariance matrix of the MLE $\hat{\beta}$ can be written *exactly* as

$$\text{Cov}(\hat{\beta}) = \sigma^2(X^T X)^{-1}$$

- ▶ In logistic regression and other GLMs, we make confidence regions and model comparisons based on constructs whose *asymptotic* (=approximately true, and increasingly accurate as sample sizes increase) form we can state

Confidence regions for logistic regression

- ▶ Let us write the linear-predictor part of our GLM in matrix form:

$$\boldsymbol{\eta} = \mathbf{X}\boldsymbol{\beta}$$

- ▶ In linear regression, we built confidence regions for parameter estimates on the basis that the covariance matrix of the MLE $\hat{\boldsymbol{\beta}}$ can be written *exactly* as

$$\text{Cov}(\hat{\boldsymbol{\beta}}) = \sigma^2(\mathbf{X}^\top \mathbf{X})^{-1}$$

- ▶ In logistic regression and other GLMs, we make confidence regions and model comparisons based on constructs whose *asymptotic* (=approximately true, and increasingly accurate as sample sizes increase) form we can state
- ▶ For confidence regions: asymptotically, the covariance matrix of $\hat{\boldsymbol{\beta}}$ is

$$\text{Cov}(\hat{\boldsymbol{\beta}}) = \begin{bmatrix} \frac{\partial^2 L(\boldsymbol{\beta}_1)}{\partial \beta_1^2} & \frac{\partial^2 L(\boldsymbol{\beta}_2)}{\partial \beta_1 \beta_2} & \cdots & \frac{\partial^2 L(\boldsymbol{\beta}_m)}{\partial \beta_1 \beta_m} \\ \frac{\partial^2 L(\boldsymbol{\beta}_1)}{\partial \beta_1 \beta_2} & \frac{\partial^2 L(\boldsymbol{\beta}_2)}{\partial \beta_2^2} & \cdots & \frac{\partial^2 L(\boldsymbol{\beta}_m)}{\partial \beta_2 \beta_m} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial^2 L(\boldsymbol{\beta}_1)}{\partial \beta_1 \beta_m} & \frac{\partial^2 L(\boldsymbol{\beta}_2)}{\partial \beta_2 \beta_m} & \cdots & \frac{\partial^2 L(\boldsymbol{\beta}_m)}{\partial \beta_m^2} \end{bmatrix}$$

(when certain regularity conditions hold). This is known as the **Fisher information matrix**.

Confidence regions for logistic regression

- ▶ A confidence region for predictors of a model estimated under maximum likelihood can be constructed similarly to the case in linear regression: for any size- k subset of predictors β' , the quantity

$$(\hat{\beta}' - \beta)^T \left(\text{Cov}(\hat{\beta}') \right)^{-1} (\hat{\beta}' - \beta)^T$$

(a multivariate **Wald statistic**) is *asymptotically* χ_k^2 distributed.

Confidence regions for logistic regression

- ▶ A confidence region for predictors of a model estimated under maximum likelihood can be constructed similarly to the case in linear regression: for any size- k subset of predictors β' , the quantity

$$(\hat{\beta}' - \beta)' \left(\text{Cov}(\hat{\beta}') \right)^{-1} (\hat{\beta}' - \beta)$$

(a multivariate **Wald statistic**) is *asymptotically* χ_k^2 distributed.

- ▶ For a single model parameter β , we can equivalently say that

$$\frac{(\hat{\beta} - \beta)}{SE(\hat{\beta})}$$

is asymptotically normally distributed, where

$SE(\hat{\beta}) = \sqrt{\text{Var}(\hat{\beta})}$. This quantity for $\beta = 0$ is often called the **Wald z-statistic**.

Confidence regions for logistic regression

- ▶ A confidence region for predictors of a model estimated under maximum likelihood can be constructed similarly to the case in linear regression: for any size- k subset of predictors β' , the quantity

$$(\hat{\beta}' - \beta)^T \left(\text{Cov}(\hat{\beta}') \right)^{-1} (\hat{\beta}' - \beta)^T$$

(a multivariate **Wald statistic**) is *asymptotically* χ_k^2 distributed.

- ▶ For a single model parameter β , we can equivalently say that

$$\frac{(\hat{\beta} - \beta)}{SE(\hat{\beta})}$$


is asymptotically normally distributed, where

$SE(\hat{\beta}) = \sqrt{\text{Var}(\hat{\beta})}$. This quantity for $\beta = 0$ is often called the **Wald z-statistic**.

- ▶ **Caution!** These approximations break down when the estimates $\hat{\beta}$ are large—most notably, when a single predictor allows *perfect* prediction of an outcome (always 0, or always 1)

Confidence regions in logistic regression

For example, a confidence region for the effects of recipient and theme definiteness:



Interactions in logistic regression

- ▶ Interactions work exactly the same for all GLMs, including logistic regression, as for linear regression

Interactions in logistic regression

- ▶ Interactions work exactly the same for all GLMs, including logistic regression, as for linear regression
- ▶ Critically, the interaction terms go into the equation for the linear predictor

$$\begin{aligned}\eta = & \alpha + \beta_{\text{RecDef}} X_{\text{RecDef}} \\ & + \beta_{\text{ThemeDef}} X_{\text{ThemeDef}} \\ & + \beta_{\text{RecDef:ThemeDef}} X_{\text{RecDef}} X_{\text{ThemeDef}}\end{aligned}$$

Interactions in logistic regression

- ▶ Interactions work exactly the same for all GLMs, including logistic regression, as for linear regression
- ▶ Critically, the interaction terms go into the equation for the linear predictor

$$\begin{aligned}\eta = & \alpha + \beta_{\text{RecDef}} X_{\text{RecDef}} \\ & + \beta_{\text{ThemeDef}} X_{\text{ThemeDef}} \\ & + \beta_{\text{RecDef:ThemeDef}} X_{\text{RecDef}} X_{\text{ThemeDef}}\end{aligned}$$

- ▶ Crucial to remember the coding scheme for these categorical predictors—here we'll stay with $X_{\text{ThemeDef}} = 1$ iff theme indefinite, $X_{\text{RecDef}} = 1$ iff recipient indefinite (both 0 otherwise)

Interactions in logistic regression

- MLE fit of the with-interactions model for the *send* data:

$\hat{\alpha}$	-0.5819215
$\hat{\beta}_{\text{RecDef}}$	-16
$\hat{\beta}_{\text{ThemeDef}}$	1.803136
$\hat{\beta}_{\text{RecDef:ThemeDef}}$	13.952

- However, the standard error of $\hat{\beta}_{\text{RecDef:ThemeDef}}$ is huge:
1151563

Interactions in logistic regression

- MLE fit of the with-interactions model for the *send* data:

$\hat{\alpha}$	-0.5819215
$\hat{\beta}_{\text{RecDef}}$	-16
$\hat{\beta}_{\text{ThemeDef}}$	1.803136
$\hat{\beta}_{\text{RecDef:ThemeDef}}$	13.952

- However, the standard error of $\hat{\beta}_{\text{RecDef:ThemeDef}}$ is huge:
1151563
- This ultimately arose because there was a *perfect prediction* possible:

```
##               Definiteness of recipient and theme
## Realization   Rec=indef, Theme=def
##   Double obj.                0
##   Prep. obj.                 5
```

Interactions in logistic regression

- ▶ MLE fit of the with-interactions model for the *send* data:

$\hat{\alpha}$	-0.5819215
$\hat{\beta}_{\text{RecDef}}$	-16
$\hat{\beta}_{\text{ThemeDef}}$	1.803136
$\hat{\beta}_{\text{RecDef:ThemeDef}}$	13.952

- ▶ However, the standard error of $\hat{\beta}_{\text{RecDef:ThemeDef}}$ is huge: 1151563
- ▶ This ultimately arose because there was a *perfect prediction* possible:

```
##                               Definiteness of recipient and theme
## Realization    Rec=indef, Theme=def
##   Double obj.                                0
##   Prep. obj.                                  5
```

- ▶ Remember, in these situations you cannot trust the Wald z-statistic $(\frac{\hat{\beta}}{SE(\hat{\beta})})!$

The likelihood ratio test

- ▶ For linear regression, hypothesis testing for a single model parameter using the t -statistic yielded *exactly* the same result as explicit model comparison with the F -statistic

The likelihood ratio test

- ▶ For linear regression, hypothesis testing for a single model parameter using the t -statistic yielded *exactly* the same result as explicit model comparison with the F -statistic
- ▶ This is a special property of *linear* regression, and does *not* generalize to other GLMs (or to the more complex models we'll see further down the line)

The likelihood ratio test

- ▶ For linear regression, hypothesis testing for a single model parameter using the t -statistic yielded *exactly* the same result as explicit model comparison with the F -statistic
- ▶ This is a special property of *linear* regression, and does *not* generalize to other GLMs (or to the more complex models we'll see further down the line)
- ▶ Instead, the more general method for hypothesis testing is the **likelihood ratio test**

The likelihood ratio test

- ▶ For linear regression, hypothesis testing for a single model parameter using the t -statistic yielded *exactly* the same result as explicit model comparison with the F -statistic
- ▶ This is a special property of *linear* regression, and does *not* generalize to other GLMs (or to the more complex models we'll see further down the line)
- ▶ Instead, the more general method for hypothesis testing is the **likelihood ratio test**
- ▶ We saw this before, in the end of the chapter on frequentist hypothesis testing

The likelihood ratio test

- ▶ For *nested* models $M_0 \subset M_A$ with k_0 and k_A free parameters respectively, the statistic

$$-2 \log \frac{\max \text{Lik}_{M_0}(y)}{\max \text{Lik}_{M_A}(y)}$$

is (asymptotically, in the limit of large data) distributed as $\chi^2_{k_A - k_0}$ if M_0 is true

The likelihood ratio test

- ▶ For *nested* models $M_0 \subset M_A$ with k_0 and k_A free parameters respectively, the statistic

$$-2 \log \frac{\max \text{Lik}_{M_0}(y)}{\max \text{Lik}_{M_A}(y)}$$

is (asymptotically, in the limit of large data) distributed as $\chi^2_{k_A - k_0}$ if M_0 is true

- ▶ This statistic doesn't have the same problems that the Wald z statistic has, so it can be used very generally to compare nested models

The likelihood ratio test

- ▶ For *nested* models $M_0 \subset M_A$ with k_0 and k_A free parameters respectively, the statistic

$$-2 \log \frac{\max \text{Lik}_{M_0}(y)}{\max \text{Lik}_{M_A}(y)}$$

is (asymptotically, in the limit of large data) distributed as $\chi^2_{k_A - k_0}$ if M_0 is true

- ▶ This statistic doesn't have the same problems that the Wald z statistic has, so it can be used very generally to compare nested models
- ▶ In our case, the additive model for recipient and theme animacy had log-likelihood of -97.1 , whereas the interactive model had log-likelihood of -96.8

The likelihood ratio test

- ▶ For *nested* models $M_0 \subset M_A$ with k_0 and k_A free parameters respectively, the statistic

$$-2 \log \frac{\max \text{Lik}_{M_0}(y)}{\max \text{Lik}_{M_A}(y)}$$

is (asymptotically, in the limit of large data) distributed as $\chi^2_{k_A - k_0}$ if M_0 is true

- ▶ This statistic doesn't have the same problems that the Wald z statistic has, so it can be used very generally to compare nested models
- ▶ In our case, the additive model for recipient and theme animacy had log-likelihood of -97.1 , whereas the interactive model had log-likelihood of -96.8
- ▶ They differed in 1 parameter, and the cumulative distribution function of χ^2_1 for 0.66 is 0.582, so we conclude that the interaction is statistically

The likelihood ratio test

- ▶ For *nested* models $M_0 \subset M_A$ with k_0 and k_A free parameters respectively, the statistic

$$-2 \log \frac{\max \text{Lik}_{M_0}(y)}{\max \text{Lik}_{M_A}(y)}$$

is (asymptotically, in the limit of large data) distributed as $\chi^2_{k_A - k_0}$ if M_0 is true

- ▶ This statistic doesn't have the same problems that the Wald z statistic has, so it can be used very generally to compare nested models
- ▶ In our case, the additive model for recipient and theme animacy had log-likelihood of -97.1 , whereas the interactive model had log-likelihood of -96.8
- ▶ They differed in 1 parameter, and the cumulative distribution function of χ^2_1 for 0.66 is 0.582, so we conclude that the interaction is statistically

The likelihood ratio test

- ▶ For *nested* models $M_0 \subset M_A$ with k_0 and k_A free parameters respectively, the statistic

$$-2 \log \frac{\max \text{Lik}_{M_0}(y)}{\max \text{Lik}_{M_A}(y)}$$

is (asymptotically, in the limit of large data) distributed as $\chi^2_{k_A - k_0}$ if M_0 is true

- ▶ This statistic doesn't have the same problems that the Wald z statistic has, so it can be used very generally to compare nested models
- ▶ In our case, the additive model for recipient and theme animacy had log-likelihood of -97.1 , whereas the interactive model had log-likelihood of -96.8
- ▶ They differed in 1 parameter, and the cumulative distribution function of χ^2_1 for 0.66 is 0.582, so we conclude that the interaction is statistically **insignificant**

References I

- Bresnan, J., Cueni, A., Nikitina, T., and Baayen, H. (2007). Predicting the dative alternation. In Boume, G., Kraemer, I., and Zwarts, J., editors, *Cognitive Foundations of Interpretation*, pages 69–95. Amsterdam: Royal Netherlands Academy of Science.
- Grodner, D. and Gibson, E. (2005). Some consequences of the serial nature of linguistic input. *Cognitive Science*, 29(2):261–290.
- Traxler, M. J., Morris, R. K., and Seely, R. E. (2002). Processing subject and object relative clauses: Evidence from eye movements. *Journal of Memory and Language*, 47:69–90.