

Directional Data Slides

Brett Presnell

September 5, 2015

Introduction

Examples


Wish to analyze data in which response is a “direction”:

- ▶ 2d directional data are called *circular* data
- ▶ 3d directional data are called *spherical* data
- ▶ not all “directional” data are directions in the usual sense
- ▶ “directional” data may also arise in higher dimensions

Wind Directions

- ▶ Recorded at Col de la Roa, Italian Alps
- ▶ $n = 310$ (first 40 listed below)
- ▶ Radians, clockwise from north
- ▶ Source: Agostinelli (CSDA 2007); also R package circular


6.23	1.03	0.15	0.72	2.20
0.46	0.63	1.45	0.37	1.95
0.08	0.15	0.33	0.09	0.09
6.23	0.05	6.14	6.28	6.17
6.24	6.02	6.14	6.25	0.01
5.38	5.30	5.63	0.77	1.34
6.14	0.22	6.23	2.33	3.61
0.49	6.12	0.01	0.00	0.46



Arrival Times at an ICU

- ▶ 24-hour clock times (format `hrs.mins`)
- ▶ $n = 254$ (first 32 listed below)
- ▶ Source: Cox & Lewis (1966); also Fisher (1993) and R package `circular`

11.00	17.00	23.15	10.00	
12.00	8.45	16.00	10.00	
15.30	20.20	4.00	12.00	
2.20	12.00	5.30	7.30	
12.00	16.00	16.00	1.30	
11.05	16.00	19.00	17.45	
20.20	21.00	12.00	12.00	
18.00	22.00	22.00	22.05	



Primate Vertebrae

- ▶ Orientation of left superior facet of last lumbar vertebra in humans, gorillas, and chimpanzees
- ▶ Source: Keifer (2005 UF Anthropology MA Thesis)

! [img] (<https://en.wikipedia.org/wiki/Vertebra#/media/File:0>)

Plot of Human Data


! [img](vertebraeOnSphere.png) "Orientation of left superior

Butterfly Migrations

- ▶ Direction of travel observed for 2649 migrating butterflies in Florida
- ▶ Source: Thomas J Walker, University of Florida, Dept of Entomology and Nematology
- ▶ Other variables:
 - ▶ site: 23 locations in Florida
 - ▶ observer: Thomas Walker (tw) or James J. Whitesell (jw)
 - ▶ species: cloudless sulphur (cs), gulf fritillary (gf), long-tailed skipper (lt)
 - ▶ distance to coast (km)
 - ▶ date and time of observation
 - ▶ percentage of sky free of clouds
 - ▶ quality of sunlight: (b)right, (h)aze, (o)bstructed, (p)artly obstructed
 - ▶ presence/absence and direction (N, NE, E, SE, S, SW, W, NW) of wind
 - ▶ temperature

Why is the Analysis of Directional Data Different?

- ▶ First three observations from the wind directions data: 6.23, 1.03, 0.15
- ▶ The mean of these three numbers is 2.47
- ▶ What do you think?



Graphical Display of Directional Data

Graphical Display of Circular Data (in R)

- ▶ Have already seen simple dot plots for circular data, e.g., for the wind data:

```
windc <- circular(wind, type = "angles",
  units = "radians", template = "geographics")
require("circular")
par(mar = c(0, 0, 0, 0) + 0.1, oma = c(0,
  0, 0, 0) + 0.1)
plot(windc, cex = 1.5, axes = FALSE, bin = 360,
  stack = TRUE, sep = 0.035, shrink = 1.3)
axis.circular(at = circular(seq(0, (7/4) *
  pi, pi/4), template = "geographics"),
  labels = c("N", "NE", "E", "SE", "S",
    "SW", "W", "NW"), cex = 1.4)
ticks.circular(circular(seq(0, (15/8) * pi,
  pi/8)), zero = pi/2, rotation = "clock",
  tcl = 0.075)
```

Graphical Display of Circular Data (in R) (ctd)

- ▶ and for the ICU data:

```
## Note that pch=17 does not work properly here.  
par(mar=c(0,0,0,0)+0.1, oma=c(0,0,0,0)+0.1)  
plot(fisherB1c, cex=1.5, axes=TRUE,  
      bin=360, stack=TRUE, sep=0.035, shrink=1.3)
```

- ▶ and one more ...

Graphical Display of Circular Data (in R) (ctd)




Figure 1: "Walking directions of long-legged desert ants under three different experimental conditions:"

Graphical Display of Circular Data (in R) (ctd)

```
par(mar = c(0, 0, 0, 0) + 0.1, oma = c(0,
      0, 0, 0) + 0.1)
plot(fisherB10c$set1, units = "degrees",
      zero = pi/2, rotation = "clock", pch = 16,
      cex = 1.5)
ticks.circular(circular(seq(0, (11/6) * pi,
      pi/6)), zero = pi/2, rotation = "clock",
      tcl = 0.075)
points(fisherB10c$set2, zero = pi/2, rotation = "clock",
      pch = 16, col = "darkgrey", next.points = -0.1,
      cex = 1.5)
points(fisherB10c$set3, zero = pi/2, rotation = "clock",
      pch = 1, next.points = 0.1, cex = 1.5)
```

Circular Histograms

- ▶ Circular histograms exist (see Fisher and Mardia and Jupp) but is there a ready-made function in R?

Rose Diagrams

- ▶ Invented by Florence Nightingale (elected first female member of the Royal Statistical Society in 1859; honorary member of ASA)
- ▶ Nightingale's rose in R (see also this post and the R graph catalog)
- ▶ Note that radii of segments are proportional to *square root* of the frequencies (counts), so that areas are proportional to frequencies. Is this the right thing to do?
- ▶ Rose diagrams suffer from the same problems as histograms. The impression conveyed may depend strongly on:
 - ▶ the binwidth of the cells
 - ▶ the choice of starting point for the bins

Adding a Rose Diagram to the Plot of Wind Directions

```
rose.diag(windc, bins=16, col="darkgrey",  
          cex=1.5, prop=1.35, add=TRUE)
```

Adding a Rose Diagram to the Plot of Wind Directions






Figure 2: "Wind direction data with rose diagram with segment areas are proportional to counts (segment radii are proportional to square roots of counts)."

Changing the Binwidth



Changing the Radii

- ▶ I think that the default “radii proportional to counts” is generally best, but this is not always obvious. The scale certainly makes a big difference however.

```
rose.diag(windc, bins = 16, col = "darkgrey",  
          radii.scale = "linear", cex = 1.5, prop = 2.4,  
          add = TRUE)
```

Changing the Radii




Figure 3: "Wind direction data with rose diagram (segment radii proportional to counts)."

Kernel Density Estimates

```
lines(density.circular(windc, bw=40), lwd=2, lty=1)
```

Kernel Density Estimates




Figure 4: "Wind direction data with rose diagram and kernel density estimate."

Spherical Data

- ▶ Are there any canned routines for plotting spherical data in R?

Basic Summary Statistics

Mean Direction and Mean Resultant Length

- First three observations from the wind directions data:

theta	x	y
6.23	-0.06	1.00
1.03	0.86	0.51
0.15	0.15	0.99

- resultant (sum of direction vectors): (0.952, 2.5)
- mean vector: $(\bar{x}, \bar{y}) = (0.317, 0.833)$
- resultant length (Euclidean norm of resultant): $R = 2.675$
- mean resultant length: $\bar{R} = \text{src_R}\{\text{round}(\text{meanResultantLength}, 3)\}$
- mean direction: $(\bar{x}, \bar{y})/\bar{R} = (0.356, 0.934)$
- $\tilde{\theta} = 0.364$

Plot




Figure 5: "First three observations from the wind directions data and their sample mean direction."

Generating Random Points on the Sphere

- ▶ Wish to generate a random “direction” in d -dimensions; i.e., an observation from the uniform distribution in the $d - 1$ sphere.
- ▶ Usual way: let $X \sim N_d(0, I)$ and return $U = X/\|X\|$.
- ▶ An alternative rejection sampler:
 - ▶ Repeat until $\|X\| \leq 1$
 - ▶ Let X be uniformly distributed on the cube $[-1, 1]^d$
 - ▶ Return $U = X/\|X\|$
- ▶ What is the acceptance rate for the rejection sampler:
 - ▶ Volume of the $d - 1$ sphere is $\pi^{d/2}/\Gamma(d/2 + 1)$
 - ▶ Volume of $[-1, 1]^d$ is 2^d
 - ▶ Acceptance rate is $(\pi^{1/2}/2)^d/\Gamma(d/2 + 1)$
 - ▶ Curse of dimensionality

dimension	2	3	4	5	6	7	8	9	10
accept rate (%)	79	52	31	16	8	4	2	1	0

```

runifSphere <- function(n, dimension, method = c("norm",
"cube", "slownorm")) {
  method <- match.arg(method)
  if (method == "norm") {
    u <- matrix(rnorm(n * dimension),
      ncol = dimension)
    u <- sweep(u, 1, sqrt(apply(u * u,
      1, sum)), "/")
  } else if (method == "slownorm") {
    u <- matrix(nrow = n, ncol = dimension)
    for (i in 1:n) {
      x <- rnorm(dimension)
      xnorm <- sqrt(sum(x^2))
      u[i, ] <- x/xnorm
    }
  } else {
    u <- matrix(nrow = n, ncol = dimension)
    for (i in 1:n) {
      x <- runif(dimension, -1, 1)
      xnorm <- sqrt(sum(x^2))
      while (xnorm > 1) {
        x <- runif(dimension, -1,
          1)
        xnorm <- sqrt(sum(x^2))
      }
      u[i, ] <- x/xnorm
    }
  }
  u
}

```

Easy fix for Borel's paradox in 3-d

Take longitude $\phi \sim U(0, 2\pi)$ independent of latitude $\theta = \arcsin(2U - 1)$, $U \sim U(0, 1)$.


Rotationally Symmetric Distributions

Comparison of Projected Normal and Langevin Distributions


One way that we might compare the $L(\mu, \kappa)$ and $PN(\gamma\mu, I)$ distributions by choosing κ and γ to give the same mean resultant lengths and comparing the densities of the cosine of the angle θ between U and μ .

Of course matching mean resultant lengths is not necessarily the best way to compare these families of distributions.

$d = 2$



$d = 3$



$$d = 4$$

