

Asteroids Discovered Before Impact: Confidence Ranking of the 13 Cases

From 2008 TC3 to 2026 RW1 (CERNQ52)

2026-09-07

Summary

Thirteen asteroids have been observed in space before entering Earth’s atmosphere. The newest, 2026 RW1, was posted to the NEO Confirmation Page as CERNQ52 and discovered by the Catalina Sky Survey at Mount Lemmon (G96) on 2026 September 6 at 08:45 UTC. It entered the atmosphere at about 16:07 UTC over the Indian Ocean near 13.8 S, 116.4 E, roughly 750 km northwest of the Western Australian coast, at local midnight. It is the first Aten-class object found before impact, and the first for which a prior-night precovery (Pan-STARRS 2, F52, on September 5) was linked while the object was still in flight.

Two of the 13, 2026 JN4 and 2026 RW1, have no independent confirmation that the impact occurred. Starting with 2026 JN4 in May 2026, the Minor Planet Center adopted new wording in its discovery MPECs: it does “not add the object to the list of past impactors” until it receives a report of a ground or air impact. The same wording appears in MPEC 2026-R64 for 2026 RW1.

Two kinds of confidence

Confidence in a predicted impact combines two separate probabilities.

- **In-flight confidence, P(impact | astrometry).** Did the tracked object hit? This depends on the number of observations, the number of independent telescopes, the length of the arc and the advance notice, and the orbit uncertainty parameter U. Arcs inside Earth’s gravity well stiffen quickly, so even short arcs can yield near-certain impact solutions.
- **Post-impact confidence, P(same object | impact evidence).** Is the fireball or meteorite the tracked object? Meteorites with a camera-network trajectory that matches the pre-impact orbit are the gold standard. A calibrated sensor detection (US government satellite sensors, GOES lightning mapper, multi-station infrasound) at the predicted time and place is next. Eyewitness video is strong when it fixes a trajectory and weak when it is a single report.

For 2019 MO and 2024 UQ the second term did all the work: neither was recognized as an impactor until after the fireball was in the satellite record. They are linked impacts rather than predicted ones.

Ranking table

The table is sorted by in-flight confidence only. The in-flight rank is the sum of four rank positions: number of observations (descending), number of contributing telescopes (descending), advance notice from discovery to impact (descending), and U (ascending), with ties broken by observation count. The “Evid. rank” column preserves the earlier ordering by total evidence, which weights post-impact confirmation heavily.

Column key: **Disc.** is the MPC observatory code of the discovery station. **Tel.** is the number of distinct MPC station codes contributing astrometry. **Obs** is the total observation count in the MPC database, verified through the MPC Explorer observations API on 2026-09-07. **U** is the MPC orbit uncertainty parameter (JPL condition code). **Warning** is discovery observation to impact; an asterisk marks retrospective cases where the impact solution was computed after the event.

Rank	Object	Disc.	Tel.	Obs	U	Warning	Confirmation evidence	Evid. rank	Tier
1	2008 TC3	G96	29	883	4	20.1 h	Meteosat 8 flash, US sensors (1 kt), infrasound, KLM pilots; pre-impact spectrum and lightcurve 600 Almahata Sitta stones, 10.5 kg (ureilite)	1	Meteorites
2	2023 CX1	K88	22	368	4	6.7 h	FRIPON camera trajectory, thousands of witnesses across France and England Saint-Pierre-le-Viger stones, about 1 kg (L chondrite)	2	Meteorites
3	2024 RW1	G96	19	187	5	10.9 h	US sensors 0.2 kt at 18.0N 122.9E, matching ESA's 0.19 kt model Videos from Luzon through typhoon cloud; no stones	5	Sensor + video
4	2024 XA1	V00	15	79	5	10.3 h	Many videos across Yakutia near Olekminsk No US sensor entry; no stones found	8	Video only
5	2024 BX1	K88	16	328	5	2.7 h	Widely filmed over Berlin and Leipzig; no US sensor entry 200+ Ribbeck stones, 1.8 kg (aubrite)	3	Meteorites
6	2026 RW1	G96	7	30	4	7.4 h	None reported as of 2026-09-07; absent from CNEOS fireball database, which lags about three weeks First Aten; first prior-night precovery (F52) linked while in flight	12	Unconfirmed
7	2022 WJ1	G96	10	51	6	3.6 h	All-sky camera trajectory, widespread video from Ontario and four US states Weather radar tracked falling stones into Lake Ontario; none recovered	6	Trajectory + radar
8	2022 EB5	K88	7	177	6	2.0 h	US sensors 3.8 kt at 70.0N 9.1W; infrasound in Greenland and Norway Possible flash seen from northern Iceland; ESA model only 0.4 kt	7	Sensor
9	2018 LA	G96	4	18	6	8.5 h	US sensors (0.9 kt), many dashcam and security videos in Botswana Motopi Pan stones, Central Kalahari (howardite)	4	Meteorites
10	2019 MO	T08	2	7	9	11.6 h*	GOES-16 lightning mapper, US sensors (6 kt), infrasound, NEXRAD radar 30-minute ATLAS arc linked after the fact; Pan-STARRS precovery extended it to 2.3 h	9	Retrospective link
11	2014 AA	G96	1	7	9	20.8 h	Weak infrasound at three CTBTO stations; no optical or sensor detection Predicted corridor spanned Central America to East Africa	11	Infrasound only
12	2024 UQ	T05	2	9	7	1.8 h*	GOES flash, US sensors 0.15 kt at 30N 136W 14-minute ATLAS arc; recognized as an impactor only after the flash	10	Retrospective link
13	2026 JN4	U68	3	6	9	3.2 h	No sensor or camera detection; entered Earth's shadow 20 min before impact One eyewitness in Darwin with matching time and direction; MPC declined to list	13	Unconfirmed

Observation counts are MPC totals. JPL's Small-Body Database uses fewer in its fits for several objects (2008 TC3: 575 used; 2024 RW1: 76 used, because 92 observations from station D29 are excluded; 2023 CX1: 331; 2024 XA1: 60). The ranking uses the MPC totals.

Reading the table

The tiers by evidence. Four cases have recovered meteorites and are certain: 2008 TC3, 2023 CX1, 2024 BX1, and 2018 LA. Four more have trajectories or calibrated sensor detections at the predicted time and place: 2024 RW1, 2022 WJ1, 2022 EB5, and 2024 XA1. Three rest on a single confirming channel or a retrospective linkage: 2019 MO, 2024 UQ, and 2014 AA. Two have nothing yet: 2026 RW1 and 2026 JN4.

The in-flight ordering tells a different story. By astrometry alone 2026 RW1 rises from 12th to 6th. Thirty observations from seven stations on three continents, a 27-hour arc including the Pan-STARRS 2 precovery pair, and $U = 4$ tie it with 2008 TC3 and 2023 CX1 for the best-determined orbit in the set. Its $P(\text{impact} | \text{astrometry})$ is effectively unity. Conversely 2018 LA drops from 4th to 9th: its 18 observations from four stations were thin, and the meteorites in the Kalahari are what make it certain. 2024 BX1 drops from 3rd to 5th because its 2.7-hour warning was short even though its 328 observations were plentiful.

Where 2026 RW1 stands. What it lacks is the second term. The entry was over open ocean at local midnight, at a slow 12 km/s, with an energy near 0.01 to 0.02 kt. Objects that small usually escape the US sensor record: 2022 WJ1, 2023 CX1, 2024 BX1, and 2024 XA1 all have blank sensor entries. The Desert Fireball Network cameras in Western Australia are at the edge of their range, and there is no geostationary lightning mapper over the Indian Ocean. It may stay unconfirmed, and by the MPC's new standard it may stay off the MPC list, like 2026 JN4.

2026 RW1's two firsts. Every earlier pre-impact discovery was an Apollo. 2026 RW1 is an Aten with $a = 0.94$ AU and aphelion at 1.12 AU, so Earth overtook it near aphelion from behind at only 5 km/s. It is also the first case where observations from the previous night, by a different survey, were identified and linked while the object was still inbound. 2019 MO's Pan-STARRS precovery was found after the impact, and the earlier same-night observations of 2024 XA1 and 2026 JN4 by ZTF preceded discovery by only one to two hours.

What the class tells us

- **The population is a selection effect, not a sample.** Every object was found within roughly a lunar distance, at magnitude 16 to 20, from the night side. Warning time scales with size and aperture: the 1.5 m Mount Lemmon telescope and Pan-STARRS gave 7 to 21 hours; the 0.6 m Piskésető Schmidt and the 0.28 m SynTrack gave 2 to 3 hours. Sizes have shrunk from 3 to 4 m in 2008 to about 1 m now because the surveys improved, not because the flux changed.
- **The first Aten is what the selection effect predicts.** Night-side discovery of an Aten requires catching it near aphelion, where Earth overtakes it slowly. That is why 2026 RW1 arrived with the lowest relative velocity in the set, the gentlest entry, and the least energy, which in turn is why it is hard to confirm. Slow, low-inclination, near-1 AU orbits dominate the whole list for the same reason: they linger in the detectable volume longest.
- **Orbits are ordinary NEO orbits.** All inclinations are below 13 degrees. Perihelia run 0.59 to 0.94 AU. Ten of 13 have aphelia inside 3 AU. Three (2019 MO, 2022 EB5, 2024 RW1) reach past 4 AU.

- **Compositions are diverse.** The four recovered falls are a ureilite, a howardite, an L chondrite, and an aubrite. Four falls, four classes, none of them the H chondrites that dominate the overall fall record. Small numbers, but they hint that meter-scale impactors sample the full NEO source mix.
- **Confirmation depends on geography, not on the object.** Every meteorite case fell over populated land at night. Every unconfirmed or sensor-only case fell over ocean or remote terrain. The two unconfirmed events both fell into the same sensor gap northwest of Australia, four months apart.
- **We catch a few percent.** The CNEOS fireball database logs 30 to 40 events per year above 0.1 kt. Pre-impact discoveries ran about one per year through 2019 and now run two to four. Nine of the 13 were discovered between September and March, matching long Northern Hemisphere nights and the latitudes of the discovering telescopes.
- **Modeled energies scatter against measured ones.** Where both exist the ratio ranges from 0.2 to 10 (2022 EB5: 4 kt measured versus 0.4 kt modeled), showing that H alone cannot size a meter-class object to better than a factor of two in diameter.

One inconsistency in the sources: the ESA past-impactors table gives 2026 JN4 an entry time of 15:39 UTC, while the MPC, the International Meteor Organization, and independent computations give 13:36 to 13:44 UTC. It looks like a table error rather than a real disagreement.

Sources

- MPEC 2026-R64 (2026 RW1): <https://www.minorplanetcenter.net/mpec/K26/K26R64.html>
- MPEC 2026-J143 (2026 JN4): <https://minorplanetcenter.net/mpec/K26/K26JE3.html>
- MPC Explorer observations (verification of counts and station codes): <https://data.minorplanetcenter.net/explorer/>
- JPL Small-Body Database API (orbits, U): <https://ssd-api.jpl.nasa.gov/sbdb.api>
- CNEOS fireball database API (US government sensor detections): <https://ssd-api.jpl.nasa.gov/fireball.api>
- ESA NEOCC past impactors: <https://neo.ssa.esa.int/past-impactors>
- ESA NEOCC 2026 JN4: <https://neo.ssa.esa.int/past-impactors/2026jn4>
- Wikipedia, list of predicted asteroid impacts: https://en.wikipedia.org/wiki/List_of_predicted_asteroid_impacts_on_Earth
- Wikipedia object pages: 2014 AA, 2019 MO, 2022 EB5, 2022 WJ1, 2024 RW1, 2024 UQ, 2024 XA1, 2026 JN4, 2026 RW1 (fr)
- The Watchers on 2026 RW1: <https://watchers.news/2026/09/07/asteroid-2026-rw1-impacts-earth-over-indian-ocean-september-6-2026-13th-predicted-impactor/>
- The Watchers on 2026 JN4: <https://watchers.news/2026/05/16/asteroid-2026-jn4-impacts-earth-arafura-sea-may-15-2026/>
- EarthSky on 2026 RW1: <https://earthsky.org/space/small-asteroid-impact-australia-sep-6-2026/>