

Beyond the Ranking: Circumstances, Science, and Comparative Context

What the 13 pre-impact discoveries tell us about small impactors, and how they compare across the Solar System

2026-09-08

Purpose

report/impactor-confidence-report.md ranks the 13 pre-impact discoveries by how sure we can be that the tracked object hit and that the recovered evidence is really it. This document goes a layer deeper: the circumstances and published science behind each of the 13, the patterns that fall out of comparing them, what those patterns say about the objects versus about our detection and confirmation infrastructure, and how the whole 13-object record compares to impacts on other worlds and to Earth impacts we never saw coming. Per-object detail lives in data/impactors/*.json; comparison-event detail lives in data/comparisons/*.json (schemas documented at the end of this file).

The 13, beyond the ranking

Object	Discovered by	Recovered fall	S
2008 TC3	R. Kowalski, Catalina (G96)	Almahata Sitta — polymict ureilite, 600 stones, 10.5 kg	First pre-impact telescope spectrum matched to a recovered meteorite (Jenniskens et al. 2009, <i>Nature</i>); fall proved to be an extraordinarily heterogeneous rubble-pile breccia
2014 AA	R. Kowalski, Catalina (G96)	none (deep Atlantic)	Only one of the 13 confirmed solely by CTBTO infrasound, with zero optical or satellite corroboration
2018 LA	R. Kowalski, Catalina (G96)	Motopi Pan — howardite, 24 stones, 215 g	Orbit + petrology jointly traced the source to 4 Vesta via the v6 resonance (Jenniskens et al. 2021)
2019 MO	ATLAS, Mauna Loa (T08)	none (Caribbean)	Not recognized as a real impactor until <i>after</i> the GOES fireball; Pan-STARRS2 precovery frames found retrospectively on a partially dead detector region
2022 EB5	K. Sárneczky, Konkoly (K88)	none (Greenland Sea)	First European-discovered imminent impactor; 3.8 kt measured vs. 0.4 kt modeled — the largest measured/modeled mismatch in the set
2022 WJ1	D. Rankin, Catalina (G96)	none — likely in Lake Ontario	Weather radar tracked debris to ~850 m altitude; refined to 40–60 cm, the smallest object ever confirmed with a pre-impact orbit at the time
2023 CX1	K. Sárneczky, Konkoly (K88)	Saint-Pierre-le-Viger — L5-6 chondrite, 60 stones, 1.1 kg	First “targeted” meteor observation in history — FRIPON alerted in advance; fell 220 years after the 1803 L’Aigle fall, about 110 km away in the same region of Normandy (Egal et al. 2025, <i>Nature</i>)
2024 BX1	K. Sárneczky, Konkoly (K88)	Ribbeck — aubrite, ~200 stones, 1.8 kg	First aubrite ever recovered with a known pre-impact orbit; a candidate spectral analog for Mercury’s surface
2024 RW1	J. Fazekas, Catalina (G96)	none (Philippine Sea, Typhoon Yagi)	Measured (0.2 kt) and ESA-modeled (0.19 kt) energy agreed almost exactly; predicted time/place verified to ~0.1 s and ~100 m
2024 UQ	ATLAS, Haleakala (T05)	none (NE Pacific)	Same “linked, not predicted” pattern as 2019 MO: recognized only after the GOES flash, Catalina precovery found afterward
2024 XA1	V. F. Carvajal, Catalina Sky Survey, Bok (V00)	none yet (remote Siberian taiga)	No camera network or sensor hit; strewn field computed <i>ab initio</i> from orbit + wind alone (Gianotto et al. 2025, <i>Icarus</i>) — untested against 2008 TC3/2023 CX1’s camera-based method
2026 JN4	N. Saini et al., SynTrack (U68)	none; MPC declined to list	First case where MPC formally withheld “past impactor” status pending independent confirmation (MPEC 2026-J143)
2026 RW1	Catalina (G96)	none reported	First Aten in the set; Earth overtook it from behind near aphelion at ~5 km/s encounter velocity, entering at ~12 km/s, the slowest and lowest-energy entry in the set

What the 13 tell us about the objects

- Composition is diverse and skews rare.** All four recovered falls are different meteorite classes — ureilite, howardite, L-chondrite, aubrite — and one, the L chondrite, belongs to the ordinary-chondrite group that makes up about four fifths of witnessed falls. Two of the four are achondrites (howardite, aubrite), a class that makes up under a tenth of falls generally. With $n = 4$ this could be selection bias (differentiated, more fragile material may be more likely to survive detection and produce a well-characterized strewn field) as easily as a real signal about the meter-scale population — see the small-numbers caution below.
- Orbits are unremarkable except for one.** Twelve of the 13 are ordinary Apollo-class, Earth-crossing, low-inclination ($\leq 13^\circ$) orbits with perihelia 0.59–0.94 AU. The exception, 2026 RW1, is the first Aten (aphelion just past 1 AU) ever caught before impact — and its Aten geometry is exactly why it produced the slowest, lowest-energy, hardest-to-confirm entry in the set (Earth overtook it from behind near aphelion at ~5 km/s before gravitational acceleration; its atmospheric entry speed of ~12 km/s is the lowest in the set, just below 2008 TC3’s 12.8 km/s).
- Sizes have shrunk as detection has improved, not because the flux changed.** 2008 TC3 was ~4 m; by 2022–2026, confirmed objects routinely run under 1 m (2022 WJ1: 40–60 cm; 2026 RW1: ~0.8 m). That is a detection-system trend, not an asteroid-population trend.
- Entry geometry spans extremes.** 2024 UQ (21 km/s) and 2024 RW1 (19.7 km/s) are the fastest; 2026 RW1, entering at ~12 km/s after a ~5 km/s encounter, is the slowest, marginally below 2008 TC3 at 12.8 km/s. Warning time varies with aperture, discovery geometry, and pipeline latency rather than object hazard (0.6 m Konkoly 2.0–6.7 h; 0.5 m ATLAS 1.8 and 11.6 h, both retrospective; 0.28 m SynTrack 3.2 h; 1.5 m Mount Lemmon 3.6–21 h; 2.3 m Bok 10.3 h).

What the 13 tell us about us

- Discovery is concentrated in a small footprint.** Seven of 13 came from the Catalina Sky Survey: six from its Mount Lemmon telescope (G96) and one from its Bok telescope (V00), across 18 years and at least three named staff observers. Within that, one person — Richard Kowalski — personally discovered three (2008 TC3, 2014 AA, 2018 LA), and one more individual — Krisztián Sárneczky, working alone at Konkoly’s 0.6 m Schmidt — discovered another three (2022 EB5, 2023 CX1, 2024 BX1). Together, two people account for nearly half of all pre-impact discoveries ever made. This is as much a statement about who runs a NEOCP-monitoring pipeline nightly as about telescope power.

- **The confirmation chain has become institutional, not improvised.** 2008 TC3 was discovered and its impact confirmed through ad hoc, largely manual coordination — the first time this had ever been attempted. By 2024–2026, ESA's Aegis/Meerkat and NASA's Scout systems routinely cross-validate a NEOCP object from a 1% probability alert to a 100% impact call within hours (2024 XA1: 07:31 to 07:50 UTC) and publish predicted time/location to sub-second, sub-kilometer precision before impact (2024 RW1). The 18-year span of this dataset is also, in effect, a build log for automated imminent-impactor infrastructure.
- **“Confirmed” is a moving institutional target, not a fixed physical fact.** MPEC 2026-J143 (2026 JN4, May 2026) is the first MPC circular to explicitly withhold “past impactor” status pending independent ground/air confirmation; the same wording was reused four months later for 2026 RW1 (MPEC 2026-R64). Two objects with effectively unity P(impact | astrometry) sit outside the MPC list because both were predicted over ocean north and northwest of Australia, beyond sensor and camera coverage — a policy and geography effect, not evidence the impacts didn't happen. ESA's NEOCC, by contrast, lists 2026 JN4 as its twelfth past impactor.
- **Retrospective linkage is a real, recurring category.** 2014 AA (infrasound only), 2019 MO, and 2024 UQ (both recognized only after a GOES/sensor flash, with precovery frames found afterward) show that for roughly a quarter of this small sample, the tracked orbit alone was not what proved the impact — an independent detection channel (treaty-monitoring infrasound, geostationary lightning mappers, retrospective image search) did the confirming work instead.
- **The pace accelerated after, and largely because of, external events.** Pre-impact discoveries ran at roughly one every few years through 2013 (2008 TC3, then a 6-year gap to 2014 AA); from 2018 onward the rate is two to four per year. That acceleration follows the February 2013 Chelyabinsk airburst, which — as detailed below — triggered the creation of IAWN, SMPAG, and NASA's Planetary Defense Coordination Office, alongside the independent arrival of new surveys (ATLAS from 2017) and automated triage pipelines (Scout, Aegis, Meerkat). Technology and institution-building are intertwined here; this dataset cannot cleanly separate them, but the timing lines up.

A small-numbers caution

Thirteen objects, and only four with recovered material, is not enough to characterize a population rigorously. Every pattern above — the achondrite-heavy meteorite mix, the single-Aten outlier, the two-person discovery concentration — is consistent with a real signal but equally consistent with the accidents of who happened to be looking, with what equipment, on which nights, across less than two decades. Treat the patterns as hypotheses the next 13 detections will test, not as established statistics.

Comparison: impacts on other worlds

Body / event	Date(s)	Advance warning	Energy scale
Comet Shoemaker-Levy 9, Jupiter	1994-07-16 to 07-22	~16 months (discovered Mar 1993, already tidally captured in Jovian orbit)	Individual fragments up to ~millions of Mt TNT; largest fragments ~1–2 km
2009 Jupiter impact scar (“Wesley impact”)	2009-07-19	None	Impactor ~200–500 m; scar comparable in scale to some SL9 scars
Jupiter amateur-detected fireballs	2010-06, 2010-08, 2012-09, and others through the 2010s	None	Impactors ~5–20 m — Chelyabinsk-scale
NASA Lunar Impact Monitoring Program	2006–present, continuous	None (no atmosphere, no fireball-warning phase possible even in principle)	>300 flashes logged; brightest (2013-03-17) ~0.3–0.4 m, ~5 t TNT
Mars HiRISE/CTX new-crater survey	2006–present, continuous	None	Mostly meter-to-tens-of-meters impactors; one notable crater only ~4 m across
Mars InSight seismic detections	2020-05, 2021-02, 2021-08, 2021-12	None	Dec 2021 event: seismic magnitude ~4, ~150 m crater, impactor 5–12 m, small enough to have burned up in Earth's atmosphere

Apart from Shoemaker-Levy 9, predicted 16 months ahead, and the artificial WE0913A lunar impact, predicted weeks ahead, every impact in the table above was reconstructed after the fact. Earth's 13-object series is, as far as these sources show, the only sustained record of natural meter-scale impacts predicted *before* they happened. The InSight December 2021 event is the sharpest single comparison available: a 5–12 m body that would have burned up in Earth's atmosphere instead reached the surface and left a clean 150 m crater, because Mars's atmosphere is roughly 1% of Earth's surface density. Energy alone does not determine consequence — atmosphere and terrain do at least as much.

Comparison: Earth impacts we did not see coming

Event	Date	Energy	
Chelyabinsk superbolide	2013-02-15	~500 kt (Popova et al. 2013, <i>Science</i>)	Approached from the Sun's direction — the one blind spot ground-based night-sky surveys cannot cover — and at ~19 m was below the size of most surveys
Tunguska event	1908-06-30	~3–15 Mt (Boslough & Crawford 2008 favor 3–5 Mt; earlier blast-scaling estimates 10–15 Mt; older estimates up to 30–40 Mt)	No detection capability of any kind existed
Sikhote-Alin meteorite fall	1947-02-12	Not modeled in kt (pre-dates airburst energy modeling)	Daylight entry over remote taiga, decades before any survey existed
Carancas impact event	2007-09-15	~0.002–0.24 kt (estimates vary with target-strength assumptions)	Meter-class body, far below any survey's reach; the event class itself was thought impossible
Benešov bolide	1991-05-07	Sub-kiloton	Optically recorded in real time by the European Fireball Network, but no asteroid-survey network existed to have caught it beforehand

The energy contrast is stark: the largest *measured* energy among all 13 tracked impactors is 2019 MO at ~6 kt. Chelyabinsk was roughly 80 times larger; Tunguska hundreds to thousands of times larger. In 18 years of pre-impact detection, nothing close to Chelyabinsk scale has ever been caught in advance — the sunward blind spot that hid Chelyabinsk is a gap in kind, not just a gap in luck, since it affects every ground-based night-sky survey equally. Benešov and 2024 XA1 also make a matched pair worth noting directly: a “perfect” trajectory record — whether from cameras in 1991 or from *ab initio* orbital modeling in 2024 — is no guarantee that a meteorite will actually be found; Benešov took 20 years.

Comparison: our own hardware, coming back from deep space

A third comparison class sits alongside natural impactors: artificial objects — spent rocket stages from national space programs — that were sent toward the Moon or beyond decades ago, lost track of, and later rediscovered on Earth-approaching or Earth-bound orbits. This is a narrow category by design: it excludes the routine, growing problem of end-of-life and aborted low-Earth-orbit reentries, and it excludes controlled reentries from crewed or robotic missions returning home on purpose. What is left is a handful of cases where a translunar-injection stage, launched once and never touched again, turned up years or decades later in cislunar or heliocentric space.

Object	Origin (program/mission)	What happened	Confirmed reentry?
WT1190F	Candidate: NASA Lunar Prospector (1998) TLI stage	Tracked on-and-off from 2009; reentered Indian Ocean off Sri Lanka, 2015-11-13	Yes — the only confirmed case
2020 SO	Confirmed: NASA Surveyor 2 (1966) Centaur upper stage	Temporarily captured Nov 2020–Mar 2021, then escaped to heliocentric orbit again	No
J002E3	Confirmed: NASA Apollo 12 (1969) S-IVB third stage	Cycles between high-Earth and heliocentric orbit on a ~40-year period since 1969; last departed 2003, due back in the mid-2040s	No
WE0913A	Confirmed: CNSA Chang'e 5-T1 (2014) Long March 3C third stage	Tracked 7+ years; struck the Moon , not Earth, 2022-03-04	N/A (lunar impact)
1991 VG	Resolved: natural, not artificial	Early candidate for Apollo-era debris; close approach 1991, later shown to have no compelling artificial signature	N/A (false candidate)

WT1190F is the only confirmed case of deep-space-origin artificial debris actually reentering Earth's atmosphere. Wikipedia's own list of derelict hardware in heliocentric orbit names Saturn V S-IVB stages from most of the Apollo missions, the Apollo 10 lunar-module ascent stage *Snoopy*, and multiple lost Soviet planetary-probe stages — a real population of candidates — yet only WT1190F has been observed to come home, and even its identity was never proven, only strongly inferred from its density. 2020 SO and J002E3 show why: both made genuinely close approaches (2020 SO passed about 50,000 km from Earth's center, outside the geostationary belt but well inside the Moon's distance), and both were pushed back out to heliocentric orbit by solar radiation pressure rather than decaying further, because these lightweight hollow stages have area-to-mass ratios far higher than any natural asteroid of similar size. That same high area-to-mass-ratio signature — first used to flag WT1190F and 2020 SO as artificial — was later run in reverse on 2024 PT5 and 1991 VG to help confirm those objects were natural after all, showing the diagnostic cuts both ways.

The identification methods are almost a direct match to the confirmation methods used for the 13 natural impactors, just aimed at a different question. Where a recovered meteorite closes the loop for 2008 TC3 or 2023 CX1, spectroscopy of paint pigment (J002E3), a stainless-steel reflectance signature (2020 SO), or bulk density and radiation-pressure sensitivity alone (WT1190F, never spectroscopically confirmed) does the equivalent work for artificial objects — identifying not just that something is coming, but *what it specifically is*, sometimes down to a named 1960s mission. WE0913A adds a cautionary note from the other direction: its identity was disputed for months, briefly and wrongly pinned on a 2015 SpaceX stage, showing that even careful tracking can misattribute deep-space debris across national programs when no one deliberately claims it.

Humanity's reaction: two events, two institutions

Neither of the two step-changes visible in planetary-defense capability was triggered by anything in the 13-object series itself — both predate or run alongside it, and both are documented here from primary NASA/ESA/UN sources (see `data/comparisons/notes.md` for full sourcing and explicit caveats about what could and could not be verified).

Shoemaker-Levy 9 (1994) created “planetary defense” as a funded program. A 1992 NASA-commissioned Spaceguard Survey report had already proposed a systematic search, but with no dedicated funding. SL9's fragments hitting Jupiter over six days, watched live by Hubble with plumes thousands of kilometers high and scars larger than Earth, is what NASA's own retrospectives credit with turning that proposal into a funded mandate: by 1998 Congress had directed NASA to find 90% of near-Earth objects ≥1 km within ten years, a goal met by the end of 2010.

Chelyabinsk (2013) internationalized the response and shifted the target size down. By 2013 the 1-km Spaceguard goal was essentially met — and Chelyabinsk mattered precisely because that survey generation was never built to catch a 19 m object, and didn't. By coincidence, the UN COPUOS Working Group on Near-Earth Objects was meeting in Vienna the very day of the airburst; the timing gave its recommendations unusual force, leading to the UN's endorsement in December 2013 of the International Asteroid Warning Network (IAWN) and the Space Mission Planning Advisory Group (SMPAG). NASA formalized its Planetary Defense Coordination Office in January 2016; known-NEO counts grew 84% from 2013 to 2018; the 2017 close approach of 2012 TC4 became the first live test of the resulting international detect-track-communicate pipeline. That pipeline — Scout, Aegis, Meerkat, cross-agency real-time triage — is the same infrastructure that confirms and publishes impact solutions for objects like 2024 XA1 and 2026 RW1 within hours today.

The general pattern: continuous technical progress (better telescopes, detectors, automated pipelines) proceeds on its own funding cycle regardless of any single event, but the *scale, deadlines, and international scope* of what gets built tracks closely with vivid, undeniable events — each one closing a specific, previously underweighted gap (kilometer-scale objects after SL9; decameter-scale, populated-area risk after Chelyabinsk) rather than uniformly raising resources at every scale at once.

Synthesis

The 13-object record is simultaneously a (thin) sample of a real small-asteroid population and a much richer, better-attested record of how fast — and how unevenly — the capability to find these objects has grown. Reading it as pure asteroid science overclaims (n = 4 recovered falls cannot characterize a population); reading it as pure institutional history undersells the genuine, hard-won achievements inside it (a telescope spectrum matched to a meteorite in hand; an impact time called to a tenth of a second). The comparisons make the boundary of current capability legible in a way the 13 objects alone cannot: every other body in the Solar System, including a planet more than 300 times Earth's mass, has its impacts reconstructed only after they happen; and even on Earth, the one body where advance detection now routinely works, it has never yet reached the size class that produced Chelyabinsk, let alone Tunguska.

Data store

- `data/impactors/<slug>.json` — one file per pre-impact discovery (13 files: `2008-tc3.json` ... `2026-rw1.json`). Fields: `discovery`, `impact`, `warning_time_hours`, `orbit`, `physical`, `confirmation` (meteorites/sensors/cameras/eyewitness), `scientific_results`, `notable_papers`, `narrative`, `sources`.
- `data/comparisons/planetary-impacts.json` — six other-world events (Shoemaker-Levy 9, the 2009 Jupiter scar, 2010s Jupiter amateur fireballs, NASA's Lunar Impact Monitoring Program, Mars HiRISE/CTX crater survey, Mars InSight seismic detections). Fields: `body`, `event_name`, `date`, `description`, `discoverer_or_mission`, `scientific_significance`, `energy_or_scale`, `key_papers`, `sources`.
- `data/comparisons/earth-undetected.json` — five Earth events with no advance detection (Chelyabinsk, Tunguska, Sikhote-Alin, Carancas, Benešov). Fields: `name`, `date_utc`, `location`, `energy_estimate_kt`, `size_estimate_m`, `why_undetected`, `damage_casualties`, `detection_method`, `scientific_significance`, `key_papers`, `sources`.

- `data/comparisons/artificial-deep-space-objects.json` — five deep-space-origin artificial objects (WT1190F, 2020 SO, J002E3, WE0913A, 1991 VG): national-program rocket stages rediscovered in cislunar/heliocentric space, one false-candidate natural object, and one lunar (not Earth) impact kept for contrast. Excludes routine LEO end-of-life/aborted reentries and controlled crewed/robotic mission reentries by design. Fields: `name`, `category`, `date_utc`, `suspected_or_confirmed_origin`, `origin_space_program`, `discovery`, `orbital_history`, `identification_evidence`, `physical_characteristics`, `outcome`, `scientific_significance`, `comparison_to_neo_impactor_set`, `key_papers`, `sources`.
- `data/comparisons/notes.md` — sourced working notes on the SL9→Spaceguard and Chelyabinsk→IAWN/SMPAG/PDCO policy history, with explicit unverified-claim flags.

All files were built from primary sources where available (MPC, JPL SBDB/CNEOS, ESA NEOCC, Meteoritical Bulletin, peer-reviewed papers) with press coverage used only for circumstantial detail; each file carries its own `sources` list. Numeric figures in this document are drawn directly from those files rather than re-derived.

Sources

In addition to the sources already listed in `report/impactor-confidence-report.md` and in each `data/` JSON file: Jenniskens et al. 2009 (*Nature* 458); Jenniskens et al. 2021 (*Meteoritics & Planetary Science* 56); Egal et al. 2025 (*Nature Astronomy*); Spurný et al. 2024 (*Astronomy & Astrophysics* 686); Gianotto et al. 2025 (*Icarus* 433); Farnocchia et al. 2016 (*Icarus* 274); Popova et al. 2013 and Posiolova et al. 2022 (*Science*); Borovička et al. 2013 and Garcia et al. 2022 (*Nature/Nature Geoscience*); Hueso et al. 2010, 2013 (*ApJL, A&A*); Suggs et al. 2014 (*Icarus*); Daubar et al. 2013, 2020 (*Icarus, JGR Planets*); NASA Goddard and NASA Science retrospectives on Shoemaker-Levy 9 and Chelyabinsk; SpaceNews; UN COPUOS/IAWN/SMPAG institutional record.