Finding landslide scars from satellite data in Pacitan

A screening model for Cyclone Cempaka (Nov 2017): what it finds, what it misses, and what it hands to an expert

Model report, November 2026

Radikal Lukafiardi

© 2026 rlukafiardi

The model finds 86.8% of scars but only 9.0% of its flags are real

Answer. On 106 scars of 500 m2 or more in held-out blocks, the model detects 86.8% (target 80%: met). Of the objects it flags, 9.0% touch a scar (target 30%: not met). The criterion stays as written.

Why it still helps. An expert checks 10 false objects per true scar instead of 31 with the baselines, over 1.0% of the area.

What the false alarms are. In a blinded review of 30 large false alarms, 1 was judged a scar. Most look like terraces, river banks or roads, not unmapped landslides.

Use. A screening aid, not a map of every landslide. 5,865 candidates are delivered for expert verdicts, the 1,678 large ones first.

© 2026 rlukafiardi

The goal: flag candidate scars for expert review, with recall of at least 80% and precision of at least 30%

Recall counts inventory polygons found; precision counts flagged objects that touch a polygon.

Aim. A screening aid that flags candidate landslide scars from satellite data, so an expert reviews flags instead of digitizing the whole area by hand.

Event and place. Cyclone Cempaka, November 2017, Pacitan regency, East Java: a 1,370 km2 study domain.

Success, fixed before any model. Object recall of 80% or more on polygons of 500 m2 or more (5 pixels), at object precision of 30% or more, on held-out 3 km blocks.

Why precision is a floor. The inventory is not wall-to-wall: a flag with no polygon under it may still be a real scar that nobody mapped.

© 2026 rlukafiardi

Labels are 743 polygons from a citizen-report inventory, so a missing polygon is not proof of no landslide

Observed: labels and imagery. Assumed: the shift, and that unmapped ground is mostly no landslide.

Labels. Samodra et al. 2024 inventory (CC BY 4.0), built from 328 citizen reports extended with 50 cm imagery and field checks. It states no footprint.

Inputs. Sentinel-2 vegetation index (NDVI) before and after the storm, the 2018 Satellite Embedding, terrain and site factors on one 10 m grid: 97 features.


Rare and small. Scars cover 0.15% of the area; the median polygon is 1,096 m2 and 26% are under 500 m2, below what 10 m pixels show reliably.

Registration. Polygons sit about 12.6 m off the Sentinel-2 grid (found from the data, cause unverified) and were shifted onto it; predictions shift back to compare with the inventory.

Samodra et al. 2024, Mendeley Data 10.17632/vrtx3w6mjd.1; Sentinel-2: Copernicus; embedding: Google DeepMind (CC BY 4.0)
© 2026 rlukafiardi

Scars sit on steeper ground: the median slope at a scar is 22 degrees, against 15 across the study area

The map shows where scars were mapped, not where they happened: the inventory started from citizen reports.



- inventory polygon, centroid (n = 743)

What is drawn. Centroids of the 743 inventory polygons on a hillshade of the release's 10 m elevation model. UTM 49S, north up.

Steeper ground. 37% of scars sit on slopes of 25 degrees or more; 16% of the study area does.


Gaps. Few scars are mapped on the flat western plain. A gap is not proof of no landslide.

Samodra et al. 2024 (CC BY 4.0): inventory and elevation model; slope read from the release's slope layer

© 2026 rlukafiardi

Scores come from whole 3 km blocks held out of training, so spatial neighbours cannot leak into the test

34 of 170 blocks (seed 42) hold 157 of 743 polygons and are scored once. The score on page 9 is one run of a frozen model and threshold.



- held-out test block (34)
- training block (white edges)
- excluded as likely unmapped (5)
- inventory polygon, centroid

Model choice. Repeated 5-fold block cross-validation, 5 seeds; models compared at 80% object recall.

Why not AUC. Random folds flatter average precision by up to 0.06 and AUC hides it.

Excluded. Five blocks look unmapped and stay out of the domain.

Intervals. Block bootstrap 95%; 106 test polygons of 500 m2 or more, so intervals are wide.

© 2026 rlukafiardi

A plain vegetation–loss threshold needs 1.08% of the area to reach 50% recall, and only 7% of those pixels are scars

A threshold alone is a weak detector, so the other features have to remove false alarms.

Pixel recall	Threshold on −dB	Pixel precision	Flagged pixels	Area flagged
20%	0.292	19.4%	16,675	0.16%
50%	**0.111**	**7.0%**	**115,502**	**1.08%**
80%	-0.051	0.6%	2,034,645	19.01%

Note: −dB is NDVI loss (pre-event minus Apr–May post-event); pixel-level; recall of 80% flags a fifth of the area

Source: reports/tables/eda_naive_detector.csv

© 2026 rlukafiardi

Vegetation change and the satellite embedding complement each other: together 7.6% precision at 80% recall, against 5.0% and 3.7% alone

Cross-validated on training blocks, 5 repeats. The intervals overlap; the paired comparison is in the report.

Model	Precision at 80% recall	95% interval	Area flagged
All features (boosting), chosen	**7.6%**	**3.9% to 13.7%**	**1.0%**
dNDVI + site factors (boosting)	5.0%	2.7% to 9.4%	1.8%
Embedding + site factors (boosting)	3.7%	2.8% to 5.3%	1.2%
All features (logistic)	4.1%	2.8% to 7.8%	2.0%
Simple logistic (dNDVI + factors)	2.8%	1.9% to 5.7%	4.0%
Naive threshold on -dB min5	2.7%	2.2% to 3.8%	12%

Source: reports/tables/modeling_results.csv

© 2026 rlukafiardi

Recall target met at 86.8%, precision target not met at 9.0%; the criterion stays unchanged

Frozen model scored once on 34 held-out blocks: 1,170 objects flagged, 105 touching a scar.

86.8%

object recall, 95% interval 79.0% to 93.1% (target 80%, met)

9.0%

object precision, 95% interval 6.7% to 11.5% (target 30%, not met)

10

false objects to check per true scar (baselines: 31)

1.0%

of the test area flagged for review

© 2026 rlukafiardi

At similar recall the model flags about a third as many objects as the baselines and recovers scar shapes better

Same 34 test blocks, 157 polygons (106 of 500 m2 or more). Intervals overlap on recall; they do not on precision.

Model	Object recall	Strict recall	Object precision	Objects flagged
Chosen model (boosting)	**86.8% (79.0–93.1)**	**69.8%**	**9.0% (6.7–11.5)**	**1,170**
Simple logistic (dNDVI + factors)	89.6% (82.9–95.3)	50.9%	3.1% (2.2–4.1)	3,569
Naive threshold on -dB min5	89.6% (81.8–95.9)	34.0%	3.2% (2.1–4.4)	3,821


Note: strict recall needs overlap with a polygon (IoU 0.3); the loose 'touches' rule favours long objects

© 2026 rlukafiardi

About 10 false objects to check per true scar, against 31 for the baselines

The review workload is what an expert pays for each scar found.

False objects per true scar found, test blocks, 95% interval



Source: reports/tables/test_results.csv

© 2026 rlukafiardi

Reaching 30% precision would cost recall: on the test blocks it appears only at about 68%

The curve is descriptive (106 polygons); the operating point was frozen from training-block repeats, not read off this curve.

Object precision against object recall, test blocks, model sweep over thresholds



Source: reports/tables/test_curve.csv, test_results.csv

© 2026 rlukafiardi

Misses are the small scars: 98% of polygons over 20 pixels are found, against 35% of those under 5

Share of polygons touched by a flagged object. The baselines find more small ones, at about three times the review workload.


Polygon size	Chosen model	Simple logistic	Naive threshold
Under 5 px (outside the target)	35%	47%	45%
5 to 20 px	79%	85%	85%
Over 20 px	**98%**	**96%**	**96%**

Source: reports/tables/test_results.csv

© 2026 rlukafiardi

In the densest test block, flagged objects touch 10 of 10 scars of 500 m2 or more, and 44 other objects touch no scar

Block (6, 13) has the most inventory polygons (19) of the 34 test blocks. Its long blue object follows a river bed and counts as true because it touches a scar.



- inventory polygon
- inventory polygon, not touched
- flagged object touching a polygon
- flagged object touching none

Counts. 19 inventory polygons, 10 of them 500 m2 or more; 57 flagged objects, 13 touching a polygon.

Untouched polygons. 3 polygons are not touched by any object, all under 500 m2.

Reading it. False alarms here are small and scattered. The loose 'touches' rule also credits long objects, which is why strict recall is reported next to it.


Note: inventory frame, UTM 49S; dashed line = block edge; objects counted by their representative point

© 2026 rlukafiardi

In a blinded review, 1 of 30 large false alarms was judged a scar, 17 were not and 12 were unsure

Reviewers recognised 8 of 10 known scars shuffled in as controls; the most common note on false alarms was terrace farming (7).

Blinded visual review on Sentinel-2 true colour, share of objects by verdict



Note: only objects of 20 px or more (24% of false alarms), n = 30; the reviewer saw the same imagery as the model

Source: blinded review answers (aggregated counts)

© 2026 rlukafiardi

Four follow-up ideas were tried after the blinded review; none gave a precision gain that held up, so none was kept

Precision at 80% recall in block cross-validation, first repeat; the size prior is the mean of 5 repeats.

Object-level second stage. 12.4% against 7.8% for the single stage, but only because objects merged into large blobs and shape recovery fell. A geometry guard was added to the validation.

Prior-year NDVI. 6.9% against 7.8% without it. Terrace false alarms have the same seasonal swing as scars.

Post-event persistence (2019, 2020). 6.6% against 7.8%. Scars here are back to pre-event NDVI within a year.


Object-size prior. Best rule 8.4% against 7.6% without it, a gain of 0.8 points whose interval includes zero. Not adopted.

Source: reports/tables/stage2_results.csv, prev_raw.csv, post_raw.csv, size_prior_results.csv

© 2026 rlukafiardi

The deliverable ranks 5,865 candidates across 1,370 km2; experts review the 1,678 large ones first

Each candidate carries a score and a blank verdict column, so expert answers can become new labels.



- large candidate (1,678)
- small candidate, 3 to 19 px (4,187)
- held-out test block
- excluded as likely unmapped

Order. Candidates are ranked by score: the 1,678 large ones (20 pixels or more) come first, then 4,187 small ones.

Fields. Score, size, slope, land use, distance to river and to the nearest mapped polygon, and empty verdict and note columns.

Layers. Two layers: on the Sentinel-2 grid, and shifted back to the inventory's frame.

Note: test objects: 1,154 in the candidate file (built over the whole area), 1,170 in the scored run (built over the test blocks only)

Labels: Samodra et al. 2024 (CC BY 4.0); Sentinel-2: Copernicus; embedding: Google DeepMind (CC BY 4.0)

© 2026 rlukafiardi

Read it as a screening aid: one event, 34 test blocks and an incomplete inventory limit what the numbers say

Observed: test scores. Assumed: label registration shift and land-use class names (inferred, unconfirmed by the inventory authors).

Inventory not wall-to-wall. It started from 328 citizen reports, so precision is a floor; the blinded review suggests this explains little of the gap for large objects.

Small test set. 34 blocks and 106 polygons of 500 m2 or more: intervals are wide and the point estimates would move with another split.

Calibration. The threshold was set for 80% recall on training blocks; test recall (86.8%) is above that, inside its interval.

Ten-metre imagery. Scars under about 500 m2 are not reliably visible, and the review used the same imagery as the model.

One event, one place. Nothing here says how the model transfers to another storm or region.

© 2026 rlukafiardi

Next: expert verdicts on the candidates become new labels; two data questions stay open

Nothing in this list is scheduled; it is what the evidence points to if the work continues.

Expert verdicts. Reviewing the large candidates and recording a verdict gives labels that do not depend on the original citizen reports.

Inventory footprint. The paper does not state the inventory's completeness and its author was not reachable; of five likely-unmapped blocks, one shows excess candidates.

Class names. Land-use and lithology class names are inferred from public maps and not confirmed by the authors.


Dropped. A river-bank mask: scars sit near rivers and precision is highest there.

© 2026 rlukafiardi

Appendix: the pipeline runs from raw inventory and imagery to a frozen model, a once-only test score and a candidate file

Build, model and deliver phases; the test blocks are touched only in the deliver phase.

Architecture and data flow



Source: github.com/rlukafiardi/landslide-inventory-pacitan docs/architecture.svg

© 2026 rlukafiardi

Appendix: removing any one feature group moves precision at 80% recall by at most 1.9 points, and no interval excludes zero

Groups overlap in information and the intervals are wide: no group was shown to be needed, which is not the same as none mattering.

Variant	Precision at 80% recall	Change vs full	95% interval of change
Full model (all features)	**7.8%**	**–**	**–**
Without pixel NDVI features	6.7%	-1.1 pts	-2.1 to +1.3
Without neighbourhood NDVI features	5.9%	-1.9 pts	-4.2 to +0.7
Without terrain features	6.3%	-1.5 pts	-2.4 to +1.7
Without categorical site factors	8.2%	+0.4 pts	-1.5 to +2.5
Without distance to road	7.8%	+0.0 pts	-1.3 to +1.9
Without land use	7.1%	-0.7 pts	-2.7 to +1.3

Note: first cross-validation repeat; the 7.6% on page 8 is the mean of 5 repeats

© 2026 rlukafiardi

About

An independent analysis of public data, not an official product and not affiliated with the inventory's authors, BIG or Badan Geologi. Labels: Samodra et al. 2024 (CC BY 4.0). Sentinel-2: Copernicus. Satellite Embedding: Google and Google DeepMind (CC BY 4.0). Code is MIT licensed; data and reports keep their sources' terms.

Author

Radikal Lukafiardi, data professional, Jakarta

Related

https://github.com/rlukafiardi/landslide-inventory-pacitan

© 2026 rlukafiardi