

### Exercise 1.2

Show that the error probability is reduced by the use of  $R_3$  by computing the error probability of this code for a binary symmetric channel with noise level  $f$ .

#### Solution:

Without  $R_3$ , we have error probability  $f$ .

With  $R_3$ , we have error probability

$$\binom{3}{2}(1-f)f^2 + f^3 = 3f^2 - 3f^3 + f^3 = 3f^2 - 2f^3$$

Now,

$$\begin{aligned} f - (3f^2 - 2f^3) &= f - 3f^2 + 2f^3 \\ &= f(1 - 2f)(1 - f) > 0 \quad (\because f < \frac{1}{2}) \end{aligned}$$

so  $f > 3f^2 - 2f^3$  hence error probability reduced.

### Exercise 1.3

- (a) Show that the probability of error of  $R_N$ , the repetition code with  $N$  repetitions, is

$$p_b = \sum_{n=(N+1)/2}^N \binom{N}{n} f^n (1-f)^{N-n}, \quad (1.24)$$

for odd  $N$ .

- (b) Assuming  $f = 0.1$ , which of the terms in this sum is the biggest? How much bigger is it than the second-biggest term?
- (c) Use Stirling's approximation (p.2) to approximate the  $\binom{N}{n}$  in the largest term, and find, approximately, the probability of error of the repetition code with  $N$  repetitions.
- (d) Assuming  $f = 0.1$ , find how many repetitions are required to get the probability of error down to  $10^{-15}$ . [Answer: about 60.]

#### Solution:

- (a) If  $N$  is odd, error when  $\frac{N+1}{2}$  or more bits flipped. Hence,

$$p_b = \sum_{n=(N+1)/2}^N \binom{N}{n} f^n (1-f)^{N-n}$$

(b) Let each term  $a_n$ .

$$\begin{aligned} \frac{a_{n+1}}{a_n} &= \frac{N!n!(N-n)!f^{n+1}(1-f)^{N-n-1}}{(n+1)!(N-n-1)!N!f^n(1-f)^{N-n}} \\ &= \frac{(N-n)f}{(n+1)(1-f)} = \frac{N-n}{9(n+1)} \end{aligned}$$

Now,

$$\begin{aligned} \frac{N-n}{9(n+1)} &> 1 \\ \iff 10n+9 &< n \\ \iff n &< \frac{N-9}{10} \quad (< \frac{N+1}{2}) \end{aligned}$$

so ratio  $< 1$  for all  $n \geq \frac{N+1}{2}$ , meaning the series is decreasing, hence largest term is when  $n = \frac{N+1}{2}$ .

Ratio is  $\frac{N-\frac{N+1}{2}}{9(\frac{N+1}{2}+1)} = \frac{N-1}{9N+11} \simeq \frac{1}{9}$ .

The largest term is approximately 9 times larger than second-largest term.

(c)

$$\binom{N}{n} \approx 2^{NH_2(\frac{n}{N})}$$

Now,

$$\begin{aligned} H_2\left(\frac{\frac{N+1}{2}}{N}\right) &= \frac{N+1}{2N} \log_2\left(\frac{2N}{N+1}\right) + \frac{N-1}{2N} \log_2\left(\frac{2N}{N-1}\right) \\ &\simeq \frac{1}{2} \log_2(2) + \frac{1}{2} \log_2(2) = 1 \end{aligned}$$

So,

$$p_b \approx p_B = \binom{N}{\frac{N+1}{2}} f^{\frac{N+1}{2}} (1-f)^{N-\frac{N+1}{2}} \simeq 2^N f^{\frac{N}{2}} (1-f)^{\frac{N}{2}} = \{4f(1-f)\}^{\frac{N}{2}}$$

(d)

$$10^{-15} = \{4f(1-f)\}^{\frac{N}{2}}$$

$$N = 2 \frac{-15}{\log_{10} 4 \cdot 0.1 \cdot 0.9} = 67.91 \approx 68$$

or more accrately,

$$\begin{aligned} \ln \binom{N}{n} &\simeq (N-n) \log_2 \frac{N}{N-n} + n \log_2 \frac{N}{n} - \frac{1}{2} \log_2 (2\pi \frac{(N-n)n}{N}) \\ &= NH_2(\frac{n}{N}) - \frac{1}{2} \log(2\pi \frac{(N-n)n}{N}) \\ &\xrightarrow{n \simeq \frac{N}{2}} N - \frac{1}{2} \log_2(\frac{\pi N}{2}) \\ \binom{N}{n} &\simeq 2^{N - \frac{1}{2} \log_2(\frac{\pi N}{2})} = \frac{2^N}{\sqrt{\frac{\pi N}{2}}} \end{aligned}$$

Then,

$$\begin{aligned} p_b \approx p_B &\simeq \binom{N}{\frac{N+1}{2}} f^{\frac{N+1}{2}} (1-f)^{\frac{N-1}{2}} \\ &\simeq \frac{2^N}{\sqrt{\frac{\pi N}{2}}} f \{f(1-f)\}^{\frac{N-1}{2}} = \frac{2f}{\sqrt{\frac{\pi N}{2}}} f \{4f(1-f)\}^{\frac{N-1}{2}} \end{aligned}$$

so

$$10^{-15} = \frac{0.2}{\sqrt{\frac{\pi N}{2}}} \{4 \cdot 0.1 \cdot 0.9\}^{\frac{N-1}{2}}$$

$$\frac{N-1}{2} = \frac{-15 - \log_1 0 \frac{0.2}{\sqrt{\frac{\pi N}{2}}}}{\log_1 00.36}$$

Iterate from  $\hat{N}_1 = 67.6$

$$\begin{aligned} \frac{\hat{N}_2 - 1}{2} &= 29.95 \\ \hat{N}_2 &= 60.90 \\ \hat{N}_3 &= 61.00 \\ \hat{N}_3 &= 61.00 \end{aligned}$$

which results in 61.

Exercise

**Solution:**