

S-CALCF



NEW ZEALAND QUALIFICATIONS AUTHORITY
MANA TOHU MĀTAURANGA O AOTEAROA

Scholarship 2005

Mathematics with Calculus

FORMULAE AND TABLES BOOKLET

Refer to this booklet to answer the questions in the Question Booklet 93202Q.

Check that this booklet has pages 2–4 in the correct order and that none of these pages is blank.

YOU MAY KEEP THIS FORMULAE AND TABLES BOOKLET AT THE END OF THE EXAMINATION.

© New Zealand Qualifications Authority, 2005

All rights reserved. No part of this publication may be reproduced by any means without the prior permission of the New Zealand Qualifications Authority.

SCHOLARSHIP MATHEMATICS WITH CALCULUS – USEFUL FORMULAE

ALGEBRA

Quadratics

If $ax^2 + bx + c = 0$

$$\text{then } x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Logarithms

$$y = \log_b x \Leftrightarrow x = b^y$$

$$\log_b(xy) = \log_b x + \log_b y$$

$$\log_b \left(\frac{x}{y} \right) = \log_b x - \log_b y$$

$$\log_b(x^n) = n \log_b x$$

$$\log_b x = \frac{\log_a x}{\log_a b}$$

Complex Numbers

$$\begin{aligned} z &= x + iy \\ &= r \operatorname{cis} \theta \\ &= r(\cos \theta + i \sin \theta) \end{aligned}$$

$$\begin{aligned} \bar{z} &= x - iy \\ &= r \operatorname{cis}(-\theta) \\ &= r(\cos \theta - i \sin \theta) \end{aligned}$$

$$r = |z| = \sqrt{z\bar{z}} = \sqrt{(x^2 + y^2)}$$

$$\theta = \arg z$$

$$\text{where } \cos \theta = \frac{x}{r}$$

$$\text{and } \sin \theta = \frac{y}{r}$$

De Moivre's Theorem:
If n is any integer then
 $(r \operatorname{cis} \theta)^n = r^n \operatorname{cis}(n\theta)$

COORDINATE GEOMETRY

Straight Line

$$\text{Equation } y - y_1 = m(x - x_1)$$

Circle

$$(x - a)^2 + (y - b)^2 = r^2$$

has a centre (a, b) and radius r

Parabola

$$y^2 = 4ax \text{ or } (at^2, 2at)$$

Ellipse

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1 \text{ or } (a \cos \theta, b \sin \theta)$$

Hyperbola

$$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1 \text{ or } (a \sec \theta, b \tan \theta)$$

$$\text{asymptotes } y = \pm \frac{b}{a}x$$

CALCULUS

Differentiation

$y = f(x)$	$\frac{dy}{dx} = f'(x)$
$\ln x$	$\frac{1}{x}$
e^{ax}	ae^{ax}
$\sin x$	$\cos x$
$\cos x$	$-\sin x$
$\tan x$	$\sec^2 x$
$\sec x$	$\sec x \tan x$
$\operatorname{cosec} x$	$-\operatorname{cosec} x \cot x$
$\cot x$	$-\operatorname{cosec}^2 x$

Product rule

$$(f \cdot g)' = f \cdot g' + g \cdot f' \quad \text{or if } y = uv \text{ then } \frac{dy}{dx} = u \frac{dv}{dx} + v \frac{du}{dx}$$

Quotient rule

$$\left(\frac{f}{g}\right)' = \frac{g \cdot f' - f \cdot g'}{g^2} \quad \text{or if } y = \frac{u}{v} \text{ then } \frac{dy}{dx} = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$$

Integration

$f(x)$	$\int f(x) \, dx$
x^n	$\frac{x^{n+1}}{n+1} + c$
$\frac{1}{x}$	$\ln x + c$
$\frac{f'(x)}{f(x)}$	$\ln f(x) + c$

Composite Function or Chain Rule

$$\left(f(g)\right)' = f'(g) \cdot g'$$

$$\text{or if } y = f(u) \text{ and } u = g(x) \text{ then } \frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx}$$

Volume of Revolution

$$y = f(x) \text{ between } x = a \text{ and } x = b$$

rotated about the x -axis

$$\text{Volume} = \int_a^b \pi y^2 \, dx$$

NUMERICAL METHODS

Trapezium Rule

$$\int_a^b f(x) \, dx \approx \frac{1}{2} h [y_0 + y_n + 2(y_1 + y_2 + \dots + y_{n-1})]$$

$$\text{where } h = \frac{b-a}{n} \text{ and } y_r = f(x_r)$$

Simpson's Rule

$$\int_a^b f(x) \, dx \approx \frac{1}{3} h [y_0 + y_n + 4(y_1 + y_3 + \dots + y_{n-1}) + 2(y_2 + y_4 + \dots + y_{n-2})]$$

$$\text{where } h = \frac{b-a}{n}, y_r = f(x_r) \text{ and } n \text{ is even.}$$

TRIGONOMETRY



$$\operatorname{cosec} \theta = \frac{1}{\sin \theta}$$

$$\sec \theta = \frac{1}{\cos \theta}$$

$$\cot \theta = \frac{1}{\tan \theta}$$

$$\cot \theta = \frac{\cos \theta}{\sin \theta}$$

$$\tan \theta = \frac{\sin \theta}{\cos \theta}$$

**Sine Rule**

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

Cosine Rule

$$c^2 = a^2 + b^2 - 2ab \cos C$$

Identities

$$\cos^2 \theta + \sin^2 \theta = 1$$

$$\tan^2 \theta + 1 = \sec^2 \theta$$

$$\cot^2 \theta + 1 = \operatorname{cosec}^2 \theta$$

General Solutions

$$\text{If } \sin \theta = \sin \alpha \text{ then } \theta = n\pi + (-1)^n \alpha$$

$$\text{If } \cos \theta = \cos \alpha \text{ then } \theta = 2n\pi \pm \alpha$$

$$\text{If } \tan \theta = \tan \alpha \text{ then } \theta = n\pi + \alpha$$

where n is any integer

Compound Angles

$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$

$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$

$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$$

Double Angles

$$\sin 2A = 2 \sin A \cos A$$

$$\tan 2A = \frac{2 \tan A}{1 - \tan^2 A}$$

$$\begin{aligned} \cos 2A &= \cos^2 A - \sin^2 A \\ &= 2 \cos^2 A - 1 \\ &= 1 - 2 \sin^2 A \end{aligned}$$

Products

$$2 \sin A \cos B = \sin(A + B) + \sin(A - B)$$

$$2 \cos A \sin B = \sin(A + B) - \sin(A - B)$$

$$2 \cos A \cos B = \cos(A + B) + \cos(A - B)$$

$$2 \sin A \sin B = \cos(A - B) - \cos(A + B)$$

Sums

$$\sin C + \sin D = 2 \sin \frac{C+D}{2} \cos \frac{C-D}{2}$$

$$\sin C - \sin D = 2 \cos \frac{C+D}{2} \sin \frac{C-D}{2}$$

$$\cos C + \cos D = 2 \cos \frac{C+D}{2} \cos \frac{C-D}{2}$$

$$\cos C - \cos D = -2 \sin \frac{C+D}{2} \sin \frac{C-D}{2}$$

MEASUREMENT**Triangle**

$$\text{area} = \frac{1}{2} ab \sin C$$

Trapezium

$$\text{area} = \frac{1}{2} (a+b)h$$

Sector

$$\text{area} = \frac{1}{2} r^2 \theta$$

$$\text{arc length} = r\theta$$

Cylinder

$$\text{volume} = \pi r^2 h$$

$$\text{curved surface area} = 2\pi rh$$

Cone

$$\text{volume} = \frac{1}{3} \pi r^2 h$$

$$\text{curved surface area} = \pi r l \text{ where } l = \text{slant height}$$

Sphere

$$\text{volume} = \frac{4}{3} \pi r^3$$

$$\text{surface area} = 4\pi r^2$$