


$$\sqrt{1 + \sqrt{2 + \sqrt{3 + \sqrt{4 + \dots}}}}$$

$$1 - 1 + 1 - 1 + 1 \dots\dots\dots = ?$$



$$\sqrt{1 + \sqrt{2 + \sqrt{3 + \sqrt{4 + \dots}}}}$$

# Discrete mathematics



# Basic Structures: Sets, Functions, Sequences, Sums, and Matrices

$$\exists_{x \in \mathfrak{R}} \exists_{y \in \mathfrak{R}} ( \mathbf{x} = \mathbf{y} )$$



$$\forall_x (\mathfrak{R} / x)$$



$$\sum_{x=1}^{\infty} \frac{1}{x} = ?$$

$$\sum_{x=1}^{\infty} \mathbf{x} = ?$$

# Chapter 2

# RIZOAN TOUFIQ

ASSISTANT PROFESSOR

DEPARTMENT OF COMPUTER SCIENCE & ENGINEERING  
RAJSHAHI UNIVERSITY OF ENGINEERING & TECHNOLOGY





# Set Operations

## Section 2.2



# Section Summary

---

- ◆ Set Operations
  - Union
  - Intersection
  - Complementation
  - Difference
- ◆ More on Set Cardinality
- ◆ Set Identities
- ◆ Proving Identities
- ◆ Membership Tables

# Union


- ◆ **Definition:** Let  $A$  and  $B$  be sets. The *union* of the sets  $A$  and  $B$ , denoted by  $A \cup B$ , is the set:

$$\{x \mid x \in A \vee x \in B\}$$

- ◆ **Example:** What is  $\{1,2,3\} \cup \{3,4,5\}$ ?

**Solution:**  $\{1,2,3,4,5\}$

Venn Diagram for  $A \cup B$



# Intersection

- ◆ **Definition:** The *intersection* of sets  $A$  and  $B$ , denoted by  $A \cap B$ , is  $\{x | x \in A \wedge x \in B\}$

- ◆ Note if the intersection is empty, then  $A$  and  $B$  are said to be *disjoint*.


- ◆ **Example:** What is  $\{1,2,3\} \cap \{3,4,5\}$  ?

**Solution:**  $\{3\}$

- ◆ **Example:** What is  $\{1,2,3\} \cap \{4,5,6\}$  ?

**Solution:**  $\emptyset$

Venn Diagram for  $A \cap B$



# Complement

**Definition:** If  $A$  is a set, then the complement of the  $A$  (with respect to  $U$ ), denoted by  $\bar{A}$  is the set  $U - A$


$$\bar{A} = \{x \in U \mid x \notin A\}$$

(The complement of  $A$  is sometimes denoted by  $A^c$ .)

**Example:** If  $U$  is the positive integers less than 100, what is the complement of  $\{x \mid x > 70\}$

Solution:  $\{x \mid x \leq 70\}$

Venn Diagram for Complement




# Difference

- ◆ **Definition:** Let  $A$  and  $B$  be sets. The *difference* of  $A$  and  $B$ , denoted by  $A - B$ , is the set containing the elements of  $A$  that are not in  $B$ . The difference of  $A$  and  $B$  is also called the complement of  $B$  with respect to  $A$ .


$$A - B = \{x \mid x \in A \wedge x \notin B\} = A \cap \bar{B}$$

Venn Diagram for  $A - B$



# The Cardinality of the Union of Two Sets

- Inclusion-Exclusion  $|A \cup B| = |A| + |B| - |A \cap B|$



Venn Diagram for  $A, B, A \cap B, A \cup B$

- **Example:** Let  $A$  be the math majors in your class and  $B$  be the CS majors. To count the number of students who are either math majors or CS majors, add the number of math majors and the number of CS majors, and subtract the number of joint CS/math majors.



# Review Questions

---

**Example:**  $U = \{0,1,2,3,4,5,6,7,8,9,10\}$   $A = \{1,2,3,4,5\}$ ,  $B = \{4,5,6,7,8\}$

1.  $A \cup B$

**Solution:**  $\{1,2,3,4,5,6,7,8\}$

2.  $A \cap B$

**Solution:**  $\{4,5\}$

3.  $\bar{A}$

**Solution:**  $\{0,6,7,8,9,10\}$

4.  $\bar{B}$

**Solution:**  $\{0,1,2,3,9,10\}$

5.  $A - B$

**Solution:**  $\{1,2,3\}$

6.  $B - A$

**Solution:**  $\{6,7,8\}$

# Symmetric Difference

**Definition:** The *symmetric difference* of **A** and **B**, denoted by  $A \oplus B$  is the set

$$(A - B) \cup (B - A)$$

**Example:**


$$U = \{0,1,2,3,4,5,6,7,8,9,10\}$$

$$A = \{1,2,3,4,5\} \quad B = \{4,5,6,7,8\}$$

What is  $A \oplus B$

– **Solution:**  $\{1,2,3,6,7,8\}$

Venn Diagram



# Set Identities

---

- ◆ Identity laws

$$A \cup \emptyset = A \quad A \cap U = A$$

- ◆ Domination laws

$$A \cup U = U \quad A \cap \emptyset = \emptyset$$

- ◆ Idempotent laws

$$A \cup A = A \quad A \cap A = A$$

- ◆ Complementation law

$$\overline{(\overline{A})} = A$$

# Set Identities

---

- ◆ Commutative laws

$$A \cup B = B \cup A \quad A \cap B = B \cap A$$

- ◆ Associative laws

$$A \cup (B \cup C) = (A \cup B) \cup C$$

$$A \cap (B \cap C) = (A \cap B) \cap C$$

- ◆ Distributive laws

$$A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$$

$$A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$$

# Set Identities

---

- ◆ De Morgan's laws

$$\overline{A \cup B} = \overline{A} \cap \overline{B} \quad \overline{A \cap B} = \overline{A} \cup \overline{B}$$

- ◆ Absorption laws

$$A \cup (A \cap B) = A \quad A \cap (A \cup B) = A$$

- ◆ Complement laws

$$A \cup \overline{A} = U \quad A \cap \overline{A} = \emptyset$$



# Proving Set Identities

---

- ◆ Different ways to prove set identities:
  1. Prove that each set (side of the identity) is a subset of the other.
  2. Use set builder notation and propositional logic.
  3. Membership Tables: Verify that elements in the same combination of sets always either belong or do not belong to the same side of the identity. Use 1 to indicate it is in the set and a 0 to indicate that it is not.

# Proof of Second De Morgan Law

---

**Example:** Prove that  $\overline{A \cap B} = \overline{A} \cup \overline{B}$

**Solution:** We prove this identity by showing that:

$$1) \quad \overline{A \cap B} \subseteq \overline{A} \cup \overline{B} \quad \text{and}$$

$$2) \quad \overline{A} \cup \overline{B} \subseteq \overline{A \cap B}$$

# Proof of Second De Morgan Law

---

These steps show that:

$$x \in \overline{A \cap B}$$

$$x \notin A \cap B$$

$$\neg((x \in A) \wedge (x \in B))$$

$$\neg(x \in A) \vee \neg(x \in B)$$

$$x \notin A \vee x \notin B$$

$$x \in \overline{A} \vee x \in \overline{B}$$

$$x \in \overline{A} \cup \overline{B}$$

$$\overline{A \cap B} \subseteq \overline{A} \cup \overline{B}$$

by assumption

defn. of complement

defn. of intersection

1st De Morgan Law for Prop Logic

defn. of negation

defn. of complement

defn. of union

# Proof of Second De Morgan Law

---

These steps show that:

$$\overline{A} \cup \overline{B} \subseteq \overline{A \cap B}$$

$$x \in \overline{A} \cup \overline{B}$$

$$(x \in \overline{A}) \vee (x \in \overline{B})$$

$$(x \notin A) \vee (x \notin B)$$

$$\neg(x \in A) \vee \neg(x \in B)$$

$$\neg((x \in A) \wedge (x \in B))$$

$$\neg(x \in A \cap B)$$

$$x \in \overline{A \cap B}$$

by assumption

defn. of union

defn. of complement

defn. of negation

by 1st De Morgan Law for Prop Logic

defn. of intersection

defn. of complement



# Set-Builder Notation: Second De Morgan Law

---

$$\begin{aligned}\overline{A \cap B} &= \{x | x \notin A \cap B\} && \text{by defn. of complement} \\ &= \{x | \neg(x \in (A \cap B))\} && \text{by defn. of does not belong symbol} \\ &= \{x | \neg(x \in A \wedge x \in B)\} && \text{by defn. of intersection} \\ &= \{x | \neg(x \in A) \vee \neg(x \in B)\} && \text{by 1st De Morgan law} \\ &&& \text{for Prop Logic} \\ &= \{x | x \notin A \vee x \notin B\} && \text{by defn. of not belong symbol} \\ &= \{x | x \in \overline{A} \vee x \in \overline{B}\} && \text{by defn. of complement} \\ &= \{x | x \in \overline{A} \cup \overline{B}\} && \text{by defn. of union} \\ &= \overline{A} \cup \overline{B} && \text{by meaning of notation}\end{aligned}$$





# Membership Table

**Example:** Construct a membership table to show that the distributive law holds.

$$A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$$

**Solution:**

A	B	C	$B \cap C$	$A \cup (B \cap C)$	$A \cup B$	$A \cup C$	$(A \cup B) \cap (A \cup C)$
1	1	1	1	1	1	1	1
1	1	0	0	1	1	1	1
1	0	1	0	1	1	1	1
1	0	0	0	1	1	1	1
0	1	1	1	1	1	1	1
0	1	0	0	0	1	0	0
0	0	1	0	0	0	1	0
0	0	0	0	0	0	0	0

# Generalized Unions and Intersections

- ◆ Let  $A_1, A_2, \dots, A_n$  be an indexed collection of sets. We define:

$$\bigcup_{i=1}^n A_i = A_1 \cup A_2 \cup \dots \cup A_n$$

$$\bigcap_{i=1}^n A_i = A_1 \cap A_2 \cap \dots \cap A_n$$

These are well defined, since union and intersection are associative.

- ◆ For  $i = 1, 2, \dots$ , let  $A_i = \{i, i + 1, i + 2, \dots\}$ . Then,

$$\bigcup_{i=1}^n A_i = \bigcup_{i=1}^n \{i, i + 1, i + 2, \dots\} = \{1, 2, 3, \dots\}$$

$$\bigcap_{i=1}^n A_i = \bigcap_{i=1}^n \{i, i + 1, i + 2, \dots\} = \{n, n + 1, n + 2, \dots\} = A_n$$

# Computer Representation of Sets

---

- ◆ There are various ways to represent sets using a computer. One method is to store the elements of the set in an unordered fashion
- ◆ Assume that the universal set  $U$  is finite (and of reasonable size so that the number of elements of  $U$  is not larger than the memory size of the computer being used).
- ◆ First, specify an arbitrary ordering of the elements of  $U$ , for instance  $a_1, a_2, \dots, a_n$ . Represent a subset  $A$  of  $U$  with the bit string of length  $n$ , where the  $i$ th bit in this string is 1 if  $a_i$  belongs to  $A$  and is 0 if  $a_i$  does not belong to  $A$ .

# Computer Representation of Sets

- ◆ **Example:** Let  $U = \{1, 2, 3, 4, 5, 6, 7, 8, 9, 10\}$ , and the ordering of elements of  $U$  has the elements in increasing order; that is,  $a_i = i$ .
  - What bit strings represent
    - the subset of all odd integers in  $U$ ,
    - the subset of all even integers in  $U$ , and
    - the subset of integers not exceeding 5 in  $U$ ?
- ◆ **Solution:**
  - The bit string that represents the set of odd integers in  $U$ , namely,  $\{1, 3, 5, 7, 9\}$ , has a one bit in the first, third, fifth, seventh, and ninth positions, and a zero elsewhere. It is 10 1010 1010.
  - Similarly, we represent the subset of all even integers in  $U$ , namely,  $\{2, 4, 6, 8, 10\}$ , by the string 01 0101 0101.
  - The set of all integers in  $U$  that do not exceed 5, namely,  $\{1, 2, 3, 4, 5\}$ , is represented by the string 11 1110 0000.

# Query???



$$\sqrt{1 + \sqrt{2 + \sqrt{3 + \sqrt{4 \dots}}}}$$

$$\exists_{x \in \mathbb{R}} \exists_{y \in \mathbb{R}} (x = y) = ?$$

$$\sum_{x=1}^{\infty} x = ?$$

$$\sum_{x=1}^{\infty} \frac{1}{x} = ?$$

$$\forall_x (\mathbb{R} / x) = ?$$

$$\exists_{x \in \mathbb{R}} \exists_{y \in \mathbb{R}} (x = y) = ?$$



$$\sqrt{1 + \sqrt{2 + \sqrt{3 + \sqrt{4 \dots}}}} = ?$$

$$1 - 1 + 1 - 1 + 1 \dots \dots = ?$$

$$\sum_{x=1}^{\infty} \frac{1}{x} = ?$$