

Nanocarriers for targeted drug delivery

Benefits and challenges of nanotechnology for medicine

Russell Maguire

ENGI4131 Advanced Semiconductor Devices

Durham University

December 2, 2018

Nanocarrier delivery systems

Introduction

Structure, types and features





Figure: Structure of common types of nanocarriers

■ hydrophobic/lipophilic, ■ hydrophilic

Definition: nanocarrier

- Biocompatible nanoparticle encapsulating a drug

Characteristics

- Nanometer scale
- Biodegradable
- Non-immunogenic
- Long half-life
- Low toxicity
- Hydrophilic
- Soluble

Targeted drug delivery

Introduction

How is targeted drug delivery effective?




Figure: Goal is to increase drug concentration ratio for diseased tissue

Targets diseased tissue

- Maximise therapeutic effects
- Minimise effective drug dose
- Common targets are solid tumours, inflamed and infected tissue

Avoids healthy tissue

- Minimise toxic side-effects

Enhanced permeability and retention effect

Passive targeting

Basis for **first generation** nanomedicines [1]

Illustration of the EPR effect




Figure: *Lymphatic system drainage and blood vessel permeability in solid tumours*

Definition: EPR effect

- Tendency for nanoparticles to accumulate in **solid tumours** [2]

Accumulation procedure

- ① Molecules leak from blood into tissue due to enhanced vascular permeability
- ② Blocked lymphatic system due to dense rapidly growing tumour cells [3]
- ③ Large particles become trapped

[1] A. Wicki *et al.*, *Journal of controlled release* **200**, 138–157 (2015).

[2] H. Maeda *et al.*, *Journal of controlled release* **65**, 271–284 (2000).

[3] T. D. Roberts *et al.*, *Nature* **427**, 695 (2004).

Internal stimuli-responsive nanocarriers

Stimuli-responsive targeting

Example




Figure: Polymeric micelles with pH-responsive polymers for electrostatic release [4]

Bioindicators used to trigger release

- pH level
- Enzymes
- Redox potential
- Temperature

Target sites for pH-responsive nanocarriers

- Tumours have been observed with lower pH than healthy tissue [5, 6]
- Individual organelles within the cell maintain their own unique pH [1]

[1] A. Wicki *et al.*, *Journal of controlled release* **200**, 138–157 (2015).

[4] T. Sun *et al.*, *Angewandte Chemie International Edition* **53**, 12320–12364 (2014).

[5] I. F. Tannock, D. Rotin, *Cancer research* **49**, 4373–4384 (1989).

[6] L. E. Gerweck, K. Seetharaman, *Cancer research* **56**, 1194–1198 (1996).

External stimuli-responsive nanocarriers

Stimuli-responsive targeting

Example




Figure: *Liposomes modified with thermosensitive polymers for controlled release [7, 8]*

Mechanisms used to externally trigger release

- Electromagnetic fields
- Heat
- Ultrasound
- Light

Induced local hyperthermia

- Target temperature is 40–45 °C [9, 10]
- Direct heat can be applied to target
- Magnetic fields can be used to heat iron oxide nanoparticles at target [11]

[7] T. Ta, T. M. Porter, *Journal of controlled release* **169**, 112–125 (2013).

[8] K. Kono, *Advanced drug delivery reviews* **53**, 307–319 (2001).

[9] A. Jhaveri *et al.*, *Journal of controlled release* **190**, 352–370 (2014).

[10] S. Ganta *et al.*, *Journal of controlled release* **126**, 187–204 (2008).

[11] F. Scherer *et al.*, *Gene therapy* **9**, 102 (2002).

Nanocarriers with surface ligands

Active targeting

Receptor-mediated endocytosis of nanocarriers




Figure: Nanocarrier binding ①, endocytosis ②, endosome transport ③, and drug release ④

Definition: active targeting

- Targeted drug delivery to specific cell types by encouraging cell uptake

Cell uptake procedure

- ① Passive targeting delivers nanocarrier to solid tumour, eg. via EPR effect [2]
- ② Surface ligands on nanocarrier bind to surface receptors on target cell
- ③ Cell absorbs nanocarrier via receptor-mediated endocytosis [4]

[2] H. Maeda *et al.*, *Journal of controlled release* **65**, 271–284 (2000).

[4] T. Sun *et al.*, *Angewandte Chemie International Edition* **53**, 12320–12364 (2014).

Nanocarrier chemotherapy drugs on the market

Examples

Doxil[®]/Caelyx[®] [12]

- Pegylated liposomal *doxorubicin*
- Immunoliposome using *polyethylene glycol (PEG)* stealth coating
- Passively targeted *doxorubicin*
- Reduced cardiac side effects [1]
- Clinical trials are exploring *actively targeted variations* [13]

Abraxane[®] [14]

- Nanoparticle *albumin-bound paclitaxel*
- Protein-drug conjugate—*albumin* is a common transport protein in blood
- Passively targeted *paclitaxel*
- Improved solubility [1]
- Unavailable on NHS from 2015–2017, partly due to cost [15]

[1] A. Wicki *et al.*, *Journal of controlled release* **200**, 138–157 (2015).

[12] M. Harrison *et al.*, *Journal of clinical oncology* **13**, 914–920 (1995).

[13] C. Mamot *et al.*, *The lancet oncology* **13**, 1234–1241 (2012).

[14] N. Desai *et al.*, *Clinical cancer research* **12**, 1317–1324 (2006).

[15] National Institute for Health and Care Excellence, [Accessed: 2018-11-30] (<https://www.nice.org.uk/guidance/ta362>).

Thank you, any questions?

1. A. Wicki *et al.*, *Journal of controlled release* **200**, 138–157 (2015).
2. H. Maeda *et al.*, *Journal of controlled release* **65**, 271–284 (2000).
3. T. P. Padera *et al.*, *Nature* **427**, 695 (2004).
4. T. Sun *et al.*, *Angewandte Chemie International Edition* **53**, 12320–12364 (2014).
5. I. F. Tannock, D. Rotin, *Cancer research* **49**, 4373–4384 (1989).
6. L. E. Gerweck, K. Seetharaman, *Cancer research* **56**, 1194–1198 (1996).
7. T. Ta, T. M. Porter, *Journal of controlled release* **169**, 112–125 (2013).
8. K. Kono, *Advanced drug delivery reviews* **53**, 307–319 (2001).
9. A. Jhaveri *et al.*, *Journal of controlled release* **190**, 352–370 (2014).
10. S. Ganta *et al.*, *Journal of controlled release* **126**, 187–204 (2008).
11. F. Scherer *et al.*, *Gene therapy* **9**, 102 (2002).
12. M. Harrison *et al.*, *Journal of clinical oncology* **13**, 914–920 (1995).
13. C. Mamot *et al.*, *The lancet oncology* **13**, 1234–1241 (2012).
14. N. Desai *et al.*, *Clinical cancer research* **12**, 1317–1324 (2006).
15. National Institute for Health and Care Excellence, [Accessed: 2018-11-30] (<https://www.nice.org.uk/guidance/ta362>).