

ECHINOCOCCUS GRANULOSIS

CLASSIFICATION :

PHYLUM : PLATYHELMINTHES

CLASS : CESTODA

ORDER : CYCLOPHILLIDAE

FAMILY : TAENIDAE

GENUS : ECHINOCOCCUS

SPECIE : GRANULOSIS

CHARACTERS :

- 1.) THEIR EGGS LIBERATED IN FAECES FORM FROM SMALL INTESTINE OF DEFINITIVE HOST (DOG , WOLF)
- 2.) EGGS SMALL IN SIZE AND OVAL IN SHAPE AND RESEMBLE TO OTHER EGGS OF TAENICULA
- 3.) EGG IS ABOUT 30-55 Um IN SIZE AND 25-32 Um IN BREADTH
- 4.) CONTAINS HEXACANTH EMBRYO WITH 3 PAIRS OF HOOKS

SCHISTOSOMA JAPONICUM FEMALE

CLASSIFICATION :

PHYLUM : PLATYHELMINTHES

CLASS : TREMATODA

ORDER PROTOSTOMATA

FAMILY : SCHISTOSTOMATIDAE

GENUS : SCHISTOSOMA

SPECIE : JAPONICUM

CHARACTERS :

- 1.) FEMALE LARGER THEN MALE, DARKER IN COLOR
- 2.) UNBRANCHED OVARIES LIES IN POSTERIOR PART OF BODY
- 3.) UTERUS FILLS WITH 50-350 EGGS ARRANGED IN SINGLE ROW IN ANTERIOR POSITION IN BODY



FASCIOLA HEPATICA ADULT STAGE

CLASSIFICATION :

PHYLUM : PLATYHELMINTHES

CLASS : TREMATODA

ORDER : PROTOSTOMATA

FAMILY : FASCIDAE

GENUS : FASCIOLA

SPECIE : HEPATICA ADULT

CHARACTERS :

- 1.) ONE OF THE LARGEST FLUKE OF THE WORLD
- 2.) AVERAGE 30mm IN LENGTH & 13mm IN WIDTH
- 3.) DIGESTIVE SYSTEM IS INCOMPLETE, MOUTH PRESENT & ANUS IS ABSENT

SCHISTOSOMA MANSONI MALE

CLASSIFICATION :

PHYLUM : PLATYHELMINTHES

CLASS : TREMATODA

ORDER PROTOSTOMATA


FAMILY : SCHISTOSTOMATIDAE

GENUS : SCHISTOSOMA

SPECIE : MANSONI

CHARACTERS :

- 1.) ABOUT 2cm LONG AND 0.1cm WIDE
- 2.) FUNNEL SHAPED ORAL SURFACE & ITS ANTERIOR END IS FOLLOWED BY VENTRAL SUCKERS(2ND SUCKERS)
- 3.) EXTERNAL PART COMPOSED OF DOUBLE LAYER WHICH IS CALLED MEMBRANE LAYER




TRICHINELLA SPIRALIS ENCYSTED LARVAE

CLASSIFICATION :

PHYLUM : NEMATODA
 CLASS : ADENOPHORA
 ORDER : TRICHOCEPHALIDAE
 FAMILY : TRICHINELLIDAE
 GENUS : TRICHINELLA
 SPECIE : SPIRALIS

CHARACTERS :

- 1.) COILED LARVA ENCLOSED WITHIN A CYST IN STRIATED SKELETAL MUSCLES
- 2.) INFECTIVE STAGE FOR HUMANS WHEN INFECTED MEAT IS EATEN RAW OR UNDERCOOKED
- 3.) LIVES INSIDE A NURSE CELL, A MODIFIED MUSCLE CELL FORMED BY THE PARASITE
- 4.) CYST ARE OVAL AND LEMON SHAPED MEASURING ABOUT 0.25-0.5mm



ANCYCLOSTOMA CRANIUM

CLASSIFICATION :

PHYLUM : NEMATODA
 CLASS : CHROMODOREA
 ORDER : RHABDITIDA
 FAMILY : ANCYCLOSTOMATIDAE
 GENUS : ANCYCLOSTOMA
 SPECIE : CRANIUM

CHARACTERS :

- 1.) FEMALE TYPICALLY 14-16cm LONG & 0.5mm WIDE
- 2.) MALES ARE SMALLER AT 10-12cm IN LENGTH & 0.36mm WIDE
- 3.) VALVE OF ANCYCLOSTOMA CRANIUM FEMALE LOCATED AT THE BOUNDARY OF 2ND AND FINAL 3RD PART OF THE BODY

