

The University of Pennsylvania Smell Identification Test (UPSIT) on Insight46 phase 1

Tom Parker | 07/06/2019

Contents

Document details	2
Methods	2
Variables in this document	2

Document details

Categories of variables	The University of Pennsylvania Smell Identification Test (UPSIT)
Summary of work undertaken	- See methods below - All returned booklets were marked by a member of the study team and updated to XNAT - Any participants with missing scores: TP searched for any unmarked booklets to ensure all returned booklets were marked - Excel spreadsheet generated by TP
Names of source variables	Xnat, number), Upsit, out, of, missing, data, change_smell, subjective, change, in, sense, of, smell), reason, data, missing/test, not, done
Output variables	upsit_i46p1, change_smell_i46p1, upsit_missing_i46p1

Methods

Prior to olfactory testing participants were asked whether they had noticed a decline in their sense of smell over the past few years? In addition, participants were prospectively excluded and did not undertake smell testing if they provided a clear history one of the following: nasal bone fracture; nasal surgery; current diagnosis of rhinitis or nasal polyps; or another chronic nasal condition that could potentially affecting olfactory function.

The University of Pennsylvania Smell Identification Test (UPSIT) was used to asses olfactory function. The UPSIT is a commercially available, well-established, reliable, and standardized olfactory test that can be self-administered. The 'British' version was selected as the most culturally appropriate to the NSHD. Each test comprises four booklets with one odorant (embedded in 10–50-micron diameter microcapsules fixed in a proprietary binder and positioned on brown strips) at the bottom of each page (ten pages per booklet). Accompanying each strip is a multiple-choice question with four responses following an alternative forced choice paradigm. Packs were provided to participants to complete at home and returned using a stamped address envelope provided. All participants were advised if they had or developed an upper respiratory tract infection to complete the test after being symptom free for 2 weeks. If a participant only completed 38 or 39 items, a score of 0.25 was assigned for each missing item. If a participant completed 37 or less items, they were excluded from the analysis.

Variables in this document

2 high-confidence matches

Variable	Label	How it was found
Sub-Studies [57] › Insight46 [699] › Physical measures [6035] › Smell test [60355]		
change_smell_i46p1	Has participant noticed a recent change in their sense of smell? - Insight46 phase 1	topsheet field
upsit_i46p1	University of Pennsylvania Smell Identification Test (UPSIT) score - Insight46 phase 1	topsheet field