Vascular medication - Insight46 phase 1

Chris Lane | May 2019

Contents

Document details 2

Summary 2

Vascular medication outcome variables at 69-71y (Phase 1 Insight 46) (2019-CL) 2

Variables 2

Variables in this document 4

Document details

Categories of variables	Vascular medication
Summary of work undertaken	Cleaning Description/Rationale: Vascular medication (antihypertensive medication, diabetic medication and lipid-regulating medication) usage extracted from data collected at Phase 1 of Insight46.
Names of source variables	See, below
Output variables	vasc_medusge_i46p1, vasc_medhtn_i46p1, vasc_meddiuretic_i46p1, vasc_medbeta_i46p1, vasc_mednitrateccb_i46p1, vasc_bplwrmed_total_i46p1, vasc_cholmed_i46p1, vasc_dmmed_i46p1, vasc_bplwrmed_bin_i46p1, vasc_diabeticmed_i46p1, vasc_cholmed_bin_i46p1

Summary

Vascular medication outcome variables at 69-71y (Phase 1 Insight 46) (2019-CL)

This document details vascular medication variables derived from Insight 46 data. All participants had medication history (usage within the last 24 hours, including over the counter medication) collected in person by a Clinical Research Associate. All participants had been advised to bring a list of their medication to their study visit. Generic or trade names were recorded, depending on what was provided. Antihypertensive medication, diabetic and lipid-regulating medication usage was then coded by CL. In the case of antihypertensive medication, a coding system based on BNF groups was used, originally derived by Mark Rawle. It should be noted that information from all medication groups is required to capture all potential anti-hypertensive medications and therefore an additional variable was derived (**bploweringmed_bin_70**). There was no missing data.

Variables

(Please not original variables are used. See below for updated variable names)

medicationusage_70 – whether any medication used (not limited to vascular medication

Coded as:

0 = no medication 1 = medication used

medhtn_70 - Presence of Antihypertensive medication. These drugs are prescribed for high blood pressure or heart failure and includes angiotensin converting enzyme inhibitors (BNF255). Includes BNF groups 2.5.1, 2.5.2, 2.5.3, 2.5.4, 2.5.5, 2.5.5.1, 2.5.5.2 & 2.5.5.3. **meddiuretic_70 -** Presence of Diuretics. These drugs may be prescribed to promote renal function and offload fluid (for example in heart failure) or to lower blood pressure. The group includes thiazide like diuretics (BNF221), loop diuretics (BNF222), potassium sparing diuretics (BNF223, BNF224) and other diuretics. Includes BNF groups 2.2.1, 2.2.2, 2.2.3, 2.2.4, 2.2.5, 2.2.6, 2.2.7 & 2.2.8.

medbeta_70 - Presence of Beta Blockers. These drugs may be prescribed for heart failure, tachycardia, anxiety, hypertension or ischaemic heart disease. Includes BNF group 2.4.

mednitrateccb_70 - Presence of Nitrates. These drugs may be prescribed to lower blood pressure, or in heart failure or ischaemic heart disease. It includes calcium channel blockers (BNF262). Includes BNF groups 2.6.1, 2.6.2, 2.6.3 & 2.6.4.

bploweringmed_bin_70 – Presence of any potential blood pressure lowering medication. Derived from **medhtn_70, meddiuretic_70, medbeta_70** and **mednitrateccb_70.**

All coded as:

0 = no medication 1 = presence of medication

bploweringmed_total70 – Total number of potential antihypertensive medications being taken. Derived from original data. **dmmed70** – Total number of diabetic medications being taken. Based on BNF groups 6.1.1 (insulins) and 6.1.2 (oral antidiabetic drugs). **diabeticmed_bin_70** – Presence of diabetic medication. Binary variable derived from **dmmed70**. Based on BNF groups 6.1.1 (insulins) and 6.1.2 (oral antidiabetic drugs). **cholmed70** – Total number of lipid-regulating medications being taken. Based on BNF group 2.12. **cholmed_bin_70** – Presence of lipid-regulating medication. Binary variable derived from **cholmed70.** Based on BNF group 2.12.

Summary statistics

medicationusage_70	Frequency	Percent
0	89	17.73
1	413	82.27

medhtn_70	Frequency	Percent
0	369	73.51
1	133	26.49

meddiuretic_70	Frequency	Percent
0	438	87.25
1	64	12.75

medbeta_70	Frequency	Percent
0	443	88.25
1	59	11.75

mednitrateccb_70	Frequency	Percent
0	419	83.47
1	83	16.53

bploweringmed_bin_70	Frequency	Percent
0	301	59.96
1	201	40.04

bploweringmed_total70	Frequency	Percent
0	301	59.96
1	100	19.92
2	64	12.75
3	29	5.78
4	<10	-
5	<10	-

dmmed70	Frequency	Percent
0	464	92.43
1	25	4.98
2	9	1.79
3	4	0.80

diabeticmed_bin_70	Frequency	Percent
0	464	92.43
1	38	7.57

cholmed70	Frequency	Percent
0	320	63.75
1	177	35.26
2	<10	-

cholmed_bin_70	Frequency	Percent
0	320	63.75
1	182	36.25

Variables in this document

11 high-confidence matches

Variable	Label	How it was found
Sub-Studies [57] › Insight46 [699] › Health and medical history [6040] › Medications [60402]		
vasc_bplwrmed_bin_i46p1	Presence of any potential blood pressure lowering medication. Presence of any potential blood pressure lowering medication. Derived from medhtn_70, meddiuretic_70, medbeta_70 and mednitrateccb_70 - Insight46 phase 1	topsheet field
vasc_bplwrmed_total_i46p1	Total number of potential antihypertensive medications being taken. Derived from original data. - Insight46 phase 1	topsheet field
vasc_cholmed_bin_i46p1	Presence of lipid-regulating medication. Binary variable derived from cholmed70. Based on BNF group 2.12. - Insight46 phase 1	topsheet field
vasc_cholmed_i46p1	Total number of lipid-regulating medications being taken. Based on BNF group 2.12. - Insight46 phase 1	topsheet field
vasc_diabeticmed_i46p1	Presence of diabetic medication. Binary variable derived from dmmed70. Based on BNF groups 6.1.1 (insulins) and 6.1.2 (oral antidiabetic drugs). - Insight46 phase 1	topsheet field
vasc_dmmed_i46p1	Total number of diabetic medications being taken. Based on BNF groups 6.1.1 (insulins) and 6.1.2 (oral antidiabetic drugs). - Insight46 phase 1	topsheet field
vasc_medbeta_i46p1	Presence of Beta Blockers. These drugs may be prescribed for heart failure, tachycardia, anxiety, hypertension or ischaemic heart disease. Includes BNF group 2.4. - Insight46 phase 1	topsheet field
vasc_meddiuretic_i46p1	Presence of Diuretics. These drugs may be prescribed to promote renal function and offload fluid (for example in heart failure) or to lower blood pressure. The group includes thiazide like diuretics (BNF221), loop diuretics (BNF222), potassium sparing diuretics (BNF223, BNF224) and other diuretics. Includes BNF groups 2.2.1, 2.2.2, 2.2.3, 2.2.4, 2.2.5, 2.2.6, 2.2.7 & 2.2.8. - Insight46 phase 1	topsheet field
vasc_medhtn_i46p1	Presence of Antihypertensive medication. These drugs are prescribed for high blood pressure or heart failure and includes angiotensin converting enzyme inhibitors (BNF255). Includes BNF groups 2.5.1, 2.5.2, 2.5.3, 2.5.4, 2.5.5, 2.5.5.1, 2.5.5.2 & 2.5.5.3. - Insight46 phase 1	topsheet field
vasc_mednitrateccb_i46p1	Presence of Nitrates. These drugs may be prescribed to lower blood pressure, or in heart failure or ischaemic heart disease. It includes calcium channel blockers (BNF262). Includes BNF groups 2.6.1, 2.6.2, 2.6.3 & 2.6.4. - Insight46 phase 1	topsheet field
vasc_medusge_i46p1	Whether any medication used (not limited to vascular medication) - Insight46 phase 1	topsheet field

7 medium-confidence matches

Variable	Label	How it was found
Questionnaire data [55] › Health and medical history [510] › Medical conditions [5101] › Other medical conditions [51011]		
blood	Have you had anaemia in the last 10 years - at age 43 years	prose scan
Questionnaire data [55] › Medications [505] › Medication taken in last 24 hours before nurse interview [5051] › Derived medication taken in the last 24 hours [50512]		
bnf221	Derived: Medication (taken in last 24 hours) - Thiazides and related diuretics (nurse schedule) at age 60-64 years	prose scan
bnf222	Derived: Medication (taken in last 24 hours) - Loop diuretics (nurse schedule) at age 60-64 years	prose scan
bnf223	Derived: Medication (taken in last 24 hours) - Potassium-sparing diuretics and aldosterone antagonists (nurse schedule) at age 60-64 years	prose scan
bnf224	Derived: Medication (taken in last 24 hours) - Potassium-sparing diuretics with other diuretics (nurse schedule) at age 60-64 years	prose scan
bnf255	Derived: Medication (taken in last 24 hours) - Drugs affecting the renin-angiotensin system (nurse schedule) at age 60-64 years	prose scan
bnf262	Derived: Medication (taken in last 24 hours) - Calcium-channel blockers (nurse schedule) at age 60-64 years	prose scan