

BIOL 274 Homework 7

Ruby Krasnow

2024-03-28

Visualizing Solutions of Linear Systems with Constant Coefficients

For each of the following systems, state the general solution and sketch the phase portrait (with at least 4 different solutions / ICs shown):

Question 1

$$\begin{aligned}\dot{x} &= x + 2y \\ \dot{y} &= 3x + 2y\end{aligned}$$

$$A = \begin{bmatrix} 1 & 2 \\ 3 & 2 \end{bmatrix}$$

$$\det \begin{bmatrix} 1 - \lambda & 2 \\ 3 & 2 - \lambda \end{bmatrix} = 0$$

$$(1 - \lambda)(2 - \lambda) - 6 = 0$$

$$2 - \lambda - 2\lambda + \lambda^2 - 6 = 0$$

$$\lambda^2 - 3\lambda - 4 = (\lambda - 4)(\lambda + 1)$$

So we have found two distinct real eigenvalues, $\lambda_1 = 4$, $\lambda_2 = -1$. Now we need to solve $AX = \lambda X$, or $(A - \lambda I)X = 0$ with these two values of λ . Let's start with the first eigenvalue, 4.

$$A - 4I = \begin{bmatrix} 1 - 4 & 2 \\ 3 & 2 - 4 \end{bmatrix} = \begin{bmatrix} -3 & 2 \\ 3 & -2 \end{bmatrix} \rightarrow \begin{bmatrix} -3 & 2 \\ 0 & 0 \end{bmatrix}$$

$$-3x_1 + 2x_2 = 0 \rightarrow 2x_2 = 3x_1 \rightarrow v_1 = \begin{bmatrix} 2 \\ 3 \end{bmatrix}$$

And repeat for the second eigenvalue, -1:

$$A - (-I) = \begin{bmatrix} 1 + 1 & 2 \\ 3 & 2 + 1 \end{bmatrix} = \begin{bmatrix} 2 & 2 \\ 3 & 3 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}$$

$$x_1 + x_2 = 0 \quad \rightarrow \quad x_2 = -x_1 \quad \rightarrow \quad v_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

Solution

$$X = C_1 e^{4t} \begin{bmatrix} 2 \\ 3 \end{bmatrix} + C_2 e^{-t} \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

Check solution

$$X' = C_1 \begin{bmatrix} 8e^{4t} \\ 12e^{4t} \end{bmatrix} + C_2 \begin{bmatrix} -e^{-t} \\ e^{-t} \end{bmatrix}$$

$$x' = 8C_1 e^{4t} - C_2 e^{-t} = 2C_1 e^{4t} + C_2 e^{-t} + 2(3C_1 e^{4t} - C_2 e^{-t}) = x + 2y$$

$$y' = 12C_1 e^{4t} + C_2 e^{-t} = 3(2C_1 e^{4t} + C_2 e^{-t}) + 2(3C_1 e^{4t} - C_2 e^{-t}) = 3x + 2y$$

Question 2

$$\dot{x} = 4x + 2y$$

$$\dot{y} = -x + y$$

$$A = \begin{bmatrix} 4 & 2 \\ -1 & 1 \end{bmatrix}$$

$$\det \begin{bmatrix} 4 - \lambda & 2 \\ -1 & 1 - \lambda \end{bmatrix} = 0$$

$$(4 - \lambda)(1 - \lambda) + 2 = 0$$

$$\lambda^2 - 5\lambda + 6 = 0$$

$$(\lambda - 3)(\lambda - 2) = 0$$

So we have found two distinct real eigenvalues, $\lambda_1 = 3$, $\lambda_2 = 2$. Now we need to solve $AX = \lambda X$, or $(A - \lambda I)X = 0$ with these two values of λ . Let's start with the first eigenvalue, 3.

$$A - 3I = \begin{bmatrix} 4-3 & 2 \\ -1 & 1-3 \end{bmatrix} = \begin{bmatrix} 1 & 2 \\ -1 & -2 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 2 \\ 0 & 0 \end{bmatrix}$$

$$x_1 + 2x_2 = 0 \quad \rightarrow \quad 2x_2 = -x_1 \quad \rightarrow \quad v_1 = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$$

And repeat for the second eigenvalue, 2:

$$A - 2I = \begin{bmatrix} 4-2 & 2 \\ -1 & 1-2 \end{bmatrix} = \begin{bmatrix} 2 & 2 \\ -1 & -1 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}$$

$$x_1 + x_2 = 0 \quad \rightarrow \quad x_2 = -x_1 \quad \rightarrow \quad v_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

Solution

$$X = C_1 e^{3t} \begin{bmatrix} 2 \\ -1 \end{bmatrix} + C_2 e^{2t} \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

Check solution

$$x' = 6C_1 e^{3t} + 2C_2 e^{2t} = 4(2C_1 e^{3t} + C_2 e^{2t}) + 2(-C_1 e^{3t} - C_2 e^{2t}) = 4x + 2y$$

$$y' = -3C_1 e^{3t} - 2C_2 e^{2t} = -(2C_1 e^{3t} + C_2 e^{2t}) + (-C_1 e^{3t} - C_2 e^{2t}) = -x + y$$

Question 3

$$\dot{x} = 3x - 5y$$

$$\dot{y} = 5x + 3y$$

$$A = \begin{bmatrix} 3 & -5 \\ 5 & 3 \end{bmatrix}$$

$$\det \begin{bmatrix} 3 - \lambda & -5 \\ 5 & 3 - \lambda \end{bmatrix} = 0$$

$$(3 - \lambda)(3 - \lambda) + 25 = 0$$

$$\lambda^2 - 6\lambda + 34 = 0$$

$$\frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \rightarrow \frac{6 \pm \sqrt{36 - 136}}{2} = \frac{6 \pm 10i}{2} = 3 \pm 5i$$

So we have complex eigenvalues, $\lambda_1 = 3 + 5i$, $\lambda_2 = 3 - 5i$. Now we need to solve $AX = \lambda X$, or $(A - \lambda I)X = 0$ with these two values of λ .

$$A - (3 + 5i)I = \begin{bmatrix} 3 - 3 - 5i & -5 \\ 5 & 3 - 3 - 5i \end{bmatrix} = \begin{bmatrix} -5i & -5 \\ 5 & -5i \end{bmatrix} \rightarrow \begin{bmatrix} i & 1 \\ 0 & 0 \end{bmatrix}$$

$$ix_1 + x_2 = 0 \rightarrow ix_1 = -x_2 \rightarrow v_1 = \begin{bmatrix} i \\ 1 \end{bmatrix}$$

Which means the eigenvector for the other eigenvalue is the conjugate of this vector, $v_2 = \begin{bmatrix} -i \\ 1 \end{bmatrix}$.

Now we use Euler's formula:

$$e^{(3+5i)t} \begin{bmatrix} i \\ 1 \end{bmatrix} = e^{3t} e^{5it} \begin{bmatrix} i \\ 1 \end{bmatrix} = e^{3t} (\cos(5t) + i \sin(5t)) \begin{bmatrix} i \\ 1 \end{bmatrix}$$
$$e^{3t} \begin{bmatrix} i \cos(5t) - \sin(5t) \\ \cos(5t) + i \sin(5t) \end{bmatrix} = e^{3t} \left(\begin{bmatrix} -\sin(5t) \\ \cos(5t) \end{bmatrix} + i \begin{bmatrix} \cos(5t) \\ \sin(5t) \end{bmatrix} \right)$$

Having found our real and imaginary parts, we can now write our general solution as a linear combination of them:

Solution

$$X = C_1 e^{3t} \begin{bmatrix} -\sin 5t \\ \cos 5t \end{bmatrix} + C_2 e^{3t} \begin{bmatrix} \cos 5t \\ \sin 5t \end{bmatrix}$$

Check solution

$$x = -C_1 e^{3t} \sin 5t + C_2 e^{3t} \cos 5t$$

$$y = C_1 e^{3t} \cos 5t + C_2 e^{3t} \sin 5t$$

$$\begin{aligned} \dot{x} &= -5C_1 e^{3t} \cos 5t - 3C_1 e^{3t} \sin 5t - 5C_2 e^{3t} \sin 5t + 3C_2 e^{3t} \cos 5t \\ &= 3(-C_1 e^{3t} \sin 5t + C_2 e^{3t} \cos 5t) - 5(C_1 e^{3t} \cos 5t + C_2 e^{3t} \sin 5t) \\ &= 3x - 5y \end{aligned}$$

$$\begin{aligned} \dot{y} &= -5C_1 e^{3t} \sin 5t + 3C_1 e^{3t} \cos 5t + 5C_2 e^{3t} \cos 5t + 3C_2 e^{3t} \sin 5t \\ &= 5(-C_1 e^{3t} \sin 5t + C_2 e^{3t} \cos 5t) + 3(C_1 e^{3t} \cos 5t + C_2 e^{3t} \sin 5t) \\ &= 5x + 3y \end{aligned}$$

Question 4

$$\dot{x} = 6x - 8y$$

$$\dot{y} = -3x + 4y$$

$$A = \begin{bmatrix} 6 & -8 \\ -3 & 4 \end{bmatrix}$$

$$\det \begin{bmatrix} 6 - \lambda & -8 \\ -3 & 4 - \lambda \end{bmatrix} = 0$$

$$(6 - \lambda)(4 - \lambda) - 24 = 0$$

$$\lambda^2 - 10\lambda = \lambda(\lambda - 10) = 0$$

So we have found two distinct real eigenvalues, $\lambda_1 = 0$, $\lambda_2 = 10$. Now we need to solve $AX = \lambda X$, or $(A - \lambda I)X = 0$ with these two values of λ . Let's start with the first eigenvalue, 0.

$$A - 0I = \begin{bmatrix} 6 & -8 \\ -3 & 4 \end{bmatrix} \rightarrow \begin{bmatrix} 3 & -4 \\ 0 & 0 \end{bmatrix}$$

$$3x_1 - 4x_2 = 0 \rightarrow 3x_1 = 4x_2 \rightarrow v_1 = \begin{bmatrix} 4 \\ 3 \end{bmatrix}$$

And repeat for the second eigenvalue, 10:

$$A - 10I = \begin{bmatrix} 6 - 10 & -8 \\ -3 & 4 - 10 \end{bmatrix} = \begin{bmatrix} -4 & -8 \\ -3 & -6 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 2 \\ 0 & 0 \end{bmatrix}$$

$$x_1 + 2x_2 = 0 \rightarrow x_1 = -2x_2 \rightarrow v_2 = \begin{bmatrix} -2 \\ 1 \end{bmatrix}$$

Solution

$$X = C_1 \begin{bmatrix} 4 \\ 3 \end{bmatrix} + C_2 e^{10t} \begin{bmatrix} -2 \\ 1 \end{bmatrix}$$

Check solution

$$\begin{aligned} x' &= -20C_2 e^{10t} = 6(4C_1 - 2C_2 e^{10t}) - 8(3C_1 + C_2 e^{10t}) = 6x - 8y \\ y' &= 10C_2 e^{10t} = -3(4C_1 - 2C_2 e^{10t}) + 4(3C_1 + C_2 e^{10t}) = -3x + 4y \end{aligned}$$

Question 5

$$\dot{x} = 4x - 2y$$

$$\dot{y} = 2x$$

$$A = \begin{bmatrix} 4 & -2 \\ 2 & 0 \end{bmatrix}$$

$$\det \begin{bmatrix} 4 - \lambda & -2 \\ 2 & -\lambda \end{bmatrix} = 0$$

$$(4 - \lambda)(-\lambda) + 4 = 0$$

$$\lambda^2 - 4\lambda + 4 = (\lambda - 2)^2 = 0$$

So we have a repeated eigenvalue, $\lambda_1 = \lambda_2 = 2$.

If the eigenvalues of A are repeated (i.e. $\lambda_1 = \lambda_2$), then the general solution to the system $\dot{X} = AX$ is $X = e^{\lambda t} v_0 + t e^{\lambda t} v_1$, where $v_0 = \begin{bmatrix} x_0 \\ y_0 \end{bmatrix}$ and $v_1 = (A - \lambda I)v_0$.

$$\begin{aligned} v_1 &= (A - 2I)v_0 \\ &= \begin{bmatrix} 2 & -2 \\ 2 & -2 \end{bmatrix} \begin{bmatrix} x_0 \\ y_0 \end{bmatrix} \\ &= \begin{bmatrix} 2x_0 - 2y_0 \\ 2x_0 - 2y_0 \end{bmatrix} \end{aligned}$$

Which means the general solution is:

Solution

$$X = e^{2t} \begin{bmatrix} x_0 \\ y_0 \end{bmatrix} + 2t e^{2t} \begin{bmatrix} x_0 - y_0 \\ x_0 - y_0 \end{bmatrix}$$

Check solution

$$\begin{aligned}\dot{x} &= 2e^{2t}x_0 + 4te^{2t}(x_0 - y_0) + 2e^{2t}(x_0 - y_0) \\ &= 4e^{2t}x_0 + 4te^{2t}(x_0 - y_0) - 2e^{2t}y_0 \\ &= 4(e^{2t}x_0 + 2te^{2t}(x_0 - y_0)) - 2(e^{2t}y_0 + 2te^{2t}(x_0 - y_0)) \\ &= 4x - 2y\end{aligned}$$

$$\begin{aligned}\dot{y} &= 2e^{2t}y_0 + 4te^{2t}(x_0 - y_0) + 2e^{2t}(x_0 - y_0) \\ &= 2e^{2t}x_0 + 4te^{2t}(x_0 - y_0) \\ &= 2(e^{2t}x_0 + 2te^{2t}(x_0 - y_0)) \\ &= 2x\end{aligned}$$

Question 6

$$\begin{aligned}\dot{x} &= -4x + y \\ \dot{y} &= 3x - 2y\end{aligned}$$

$$A = \begin{bmatrix} -4 & 1 \\ 3 & -2 \end{bmatrix}$$

$$\det \begin{bmatrix} -4 - \lambda & 1 \\ 3 & -2 - \lambda \end{bmatrix} = 0$$

$$(-4 - \lambda)(-2 - \lambda) - 3 = 0$$

$$\lambda^2 + 6\lambda + 5 = (\lambda + 5)(\lambda + 1) = 0$$

So we have found two distinct real eigenvalues, $\lambda_1 = -5$, $\lambda_2 = -1$. Now we need to solve $AX = \lambda X$, or $(A - \lambda I)X = 0$ with these two values of λ . Let's start with the first eigenvalue, 0.

$$A + 5I = \begin{bmatrix} -4 + 5 & 1 \\ 3 & -2 + 5 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 3 & 3 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}$$

$$x_1 + x_2 = 0 \rightarrow x_1 = -x_2 \rightarrow v_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

$$A + I = \begin{bmatrix} -4 + 1 & 1 \\ 3 & -2 + 1 \end{bmatrix} = \begin{bmatrix} -3 & 1 \\ 3 & -1 \end{bmatrix} \rightarrow \begin{bmatrix} 3 & -1 \\ 0 & 0 \end{bmatrix}$$

$$3x_1 - x_2 = 0 \rightarrow 3x_1 = x_2 \rightarrow v_2 = \begin{bmatrix} 1 \\ 3 \end{bmatrix}$$

Solution

$$X = C_1 e^{-5t} \begin{bmatrix} 1 \\ -1 \end{bmatrix} + C_2 e^{-t} \begin{bmatrix} 1 \\ 3 \end{bmatrix}$$

Check solution

$$\begin{aligned} x' &= -5C_1 e^{-5t} - C_2 e^{-t} = -4(C_1 e^{-5t} + C_2 e^{-t}) + (-C_1 e^{-5t} + 3C_2 e^{-t}) = -4x + y \\ y' &= 5C_1 e^{-5t} - 3C_2 e^{-t} = 3(C_1 e^{-5t} + C_2 e^{-t}) - 2(-C_1 e^{-5t} + 3C_2 e^{-t}) = 3x - 2y \end{aligned}$$

Question 7

$$\begin{aligned} \dot{x} &= -9x - 3y \\ \dot{y} &= 3x + y \end{aligned}$$

$$A = \begin{bmatrix} -9 & -3 \\ 3 & 1 \end{bmatrix}$$

$$\det \begin{bmatrix} -9 - \lambda & -3 \\ 3 & 1 - \lambda \end{bmatrix} = 0$$

$$(-9 - \lambda)(1 - \lambda) + 9 = 0$$

$$\lambda^2 + 8\lambda = \lambda(\lambda + 8) = 0$$

So we have found two distinct real eigenvalues, $\lambda_1 = 0$, $\lambda_2 = -8$. Now we need to solve $AX = \lambda X$, or $(A - \lambda I)X = 0$ with these two values of λ . Let's start with the first eigenvalue, 0.

$$A - 0I = \begin{bmatrix} -9 & -3 \\ 3 & 1 \end{bmatrix} \rightarrow \begin{bmatrix} 3 & 1 \\ 0 & 0 \end{bmatrix}$$

$$3x_1 + x_2 = 0 \rightarrow 3x_1 = -x_2 \rightarrow v_1 = \begin{bmatrix} 1 \\ -3 \end{bmatrix}$$

And repeat for the second eigenvalue, -8:

$$A + 8I = \begin{bmatrix} -9+8 & -3 \\ 3 & 1+8 \end{bmatrix} = \begin{bmatrix} -1 & -3 \\ 3 & 9 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 3 \\ 0 & 0 \end{bmatrix}$$

$$x_1 + 3x_2 = 0 \rightarrow x_1 = -3x_2 \rightarrow v_2 = \begin{bmatrix} 3 \\ -1 \end{bmatrix}$$

Solution

$$X = C_1 \begin{bmatrix} 1 \\ -3 \end{bmatrix} + C_2 e^{-8t} \begin{bmatrix} 3 \\ -1 \end{bmatrix}$$

Check solution

$$\begin{aligned}x' &= -24C_2e^{-8t} = -9(C_1 + 3C_2e^{-8t}) - 3(-3C_1 - C_2e^{-8t}) = -9x - 3y \\y' &= 8C_2e^{-8t} = 3(C_1 + 3C_2e^{-8t}) + (-3C_1 - C_2e^{-8t}) = 3x + y\end{aligned}$$

Question 8

$$\begin{aligned}\dot{x} &= -3x + 10y \\ \dot{y} &= -x + 3y\end{aligned}$$

$$A = \begin{bmatrix} -3 & 10 \\ -1 & 3 \end{bmatrix}$$

$$\det \begin{bmatrix} -3 - \lambda & 10 \\ -1 & 3 - \lambda \end{bmatrix} = 0$$

$$(-3 - \lambda)(3 - \lambda) + 10 = 0$$

$$\lambda^2 + 1 = 0$$

So we have complex eigenvalues, $\lambda_1 = i$, $\lambda_2 = -i$. Now we need to solve $AX = \lambda X$, or $(A - \lambda I)X = 0$ with these two values of λ .

$$A - iI = \begin{bmatrix} -3 - i & 10 \\ -1 & 3 - i \end{bmatrix} \rightarrow \begin{bmatrix} -1 & 3 - i \\ 0 & 0 \end{bmatrix}$$

$$-x_1 + (3 - i)x_2 = 0 \rightarrow x_1 = (3 - i)x_2 \rightarrow v_1 = \begin{bmatrix} 3 - i \\ 1 \end{bmatrix}$$

Which means the eigenvector for the other eigenvalue is the conjugate of this vector, $v_2 = \begin{bmatrix} 3 + i \\ 1 \end{bmatrix}$.

Now we use Euler's formula:

$$e^{it} \begin{bmatrix} 3 - i \\ 1 \end{bmatrix} = (\cos(t) + i \sin(t)) \begin{bmatrix} 3 - i \\ 1 \end{bmatrix}$$

$$\begin{bmatrix} (3 - i) \cos t + (3i + 1) \sin t \\ \cos t + i \sin t \end{bmatrix} = \begin{bmatrix} 3 \cos t + \sin t \\ \cos t \end{bmatrix} + i \begin{bmatrix} 3 \sin t - \cos t \\ \sin t \end{bmatrix}$$

Having found our real and imaginary parts, we can now write our general solution as a linear combination of them:

Solution

$$X = C_1 \begin{bmatrix} 3 \cos t + \sin t \\ \cos t \end{bmatrix} + C_2 \begin{bmatrix} 3 \sin t - \cos t \\ \sin t \end{bmatrix}$$

Check solution

$$\begin{aligned}x' &= -3C_1 \sin t + C_1 \cos t + 3C_2 \cos t + C_2 \sin t \\&= -3(3C_1 \cos t + C_1 \sin t + 3C_2 \sin t - C_2 \cos t) + 10(C_1 \cos t + C_2 \sin t) \\&= -3x + 10y\end{aligned}$$

$$\begin{aligned}y' &= -C_1 \sin t + C_2 \cos t \\&= -(3C_1 \cos t + C_1 \sin t + 3C_2 \sin t - C_2 \cos t) + 3(C_1 \cos t + C_2 \sin t) \\&= -x + 3y\end{aligned}$$

Phase Portraits

```
fun <- function(t, y, parameters) {  
  x <- y[1]  
  y <- y[2]  
  
  a <- parameters[1]  
  b <- parameters[2]  
  c <- parameters[3]  
  d <- parameters[4]  
  
  dy <- numeric(2)  
  
  dy[1] <- a*x + b*y  
  dy[2] <- c*x + d*y  
  
  return(list(dy))  
}
```

```
p1+theme_classic()+  
  geom_abline(slope=-1, color="gray", linewidth=1)+  
  geom_abline(slope=3/2, color="gray", linewidth=1)+  
  geom_path(data=z, aes(x=x, y=y, color=pt), linewidth=1.5)+  
  geom_point(data=y2, aes(x=x,y=y, color=pt), shape=21, fill="white",size=5, stroke=2)+labs(color="IC")
```

