



[Course](#) > [Section...](#) > [1.3 Qu...](#) > [Definiti...](#)

Audit Access Expires Mar 24, 2020

You lose all access to this course, including your progress, on Mar 24, 2020.

Upgrade by Feb 18, 2020 to get unlimited access to the course as long as it exists on the site. [Upgrade now](#)

Definition of quantiles

Definition of quantiles

Quantiles are cutoff points that divide a dataset into intervals with set probabilities. The q th quantile is the value at which $q\%$ of the observations are equal to or less than that value.

Using the quantile function

Given a dataset `data` and desired quantile q , you can find the q th quantile of `data` with:

```
quantile(data, q)
```

Percentiles

Percentiles are the quantiles that divide a dataset into 100 intervals each with 1% probability. You can determine all percentiles of a dataset `data` like this:

```
p <- seq(0.01, 0.99, 0.01)
quantile(data, p)
```

Quartiles

Quartiles divide a dataset into 4 parts each with 25% probability. They are equal to the 25th, 50th and 75th percentiles. The 25th percentile is also known as the *1st quartile*, the 50th percentile is also known as the *median*, and the 75th percentile is also known as the *3rd quartile*.



The `summary()` function returns the minimum, quartiles and maximum of a vector.

Examples

Load the heights dataset from the **dslabs** package:

```
library(dslabs)
data(heights)
```

Use `summary()` on the `heights$height` variable to find the quartiles:

```
summary(heights$height)
```

Find the percentiles of `heights$height`:

```
p <- seq(0.01, 0.99, 0.01)
percentiles <- quantile(heights$height, p)
```

Confirm that the 25th and 75th percentiles match the 1st and 3rd quartiles. Note that `quantile()` returns a named vector. You can access the 25th and 75th percentiles like this (adapt the code for other percentile values):

```
percentiles[names(percentiles) == "25%"]
percentiles[names(percentiles) == "75%"]
```

[Learn About Verified Certificates](#)

© All Rights Reserved

