



[Course](#) > [Section...](#) > [1.2 Intr...](#) > Norma...

Audit Access Expires Mar 24, 2020

You lose all access to this course, including your progress, on Mar 24, 2020.

Upgrade by Feb 18, 2020 to get unlimited access to the course as long as it exists on the site. **Upgrade now**

Normal Distribution: Standard Units and Z-scores

Standard units

For data that are approximately normal, *standard units* describe the number of standard deviations an observation is from the mean. Standard units are denoted by the variable z and are also known as *z-scores*.

For any value x from a normal distribution with mean μ and standard deviation σ , the value in standard units is:

$$z = \frac{x - \mu}{\sigma}$$

Standard units are useful for many reasons. Note that the formula for the normal distribution is simplified by substituting z in the exponent:

$$\Pr(a < x < b) = \int_a^b \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{1}{2}z^2} dx$$

When $z = 0$, the normal distribution is at a maximum, the mean μ . The function is defined to be symmetric around $z = 0$.

The normal distribution of z-scores is called the *standard normal distribution* and is defined by $\mu = 0$ and $\sigma = 1$.



Z-scores are useful to quickly evaluate whether an observation is average or extreme. Z-scores near 0 are average. Z-scores above 2 or below -2 are significantly above or below the mean, and z-scores above 3 or below -3 are extremely rare.

We will learn more about benchmark z-score values and their corresponding probabilities below.

Code: Converting to standard units

The `scale` function converts a vector of approximately normally distributed values into z-scores.

```
z <- scale(x)
```


You can compute the proportion of observations that are within 2 standard deviations of the mean like this:

```
mean(abs(z) < 2)
```

The 68-95-99.7 Rule

The normal distribution is associated with the 68-95-99.7 rule. This rule describes the probability of observing events within a certain number of standard deviations of the mean.






The probability distribution function for the normal distribution is defined such that:

- About 68% of observations will be within one standard deviation of the mean ($\mu \pm \sigma$). In standard units, this is equivalent to a z-score of $|z| \leq 1$.




Probability of an observation within 1 SD of mean



- About 95% of observations will be within one standard deviation of the mean ($\mu \pm 2\sigma$). In standard units, this is equivalent to a z-score of $|z| \leq 2$.




Probability of an observation within 2 SD of mean



- About 99.7% of observations will be within one standard deviation of the mean ($\mu \pm 3\sigma$). In standard units, this is equivalent to a z-score of $|z| \leq 3$.



Probability of an observation within 3 SD of mean



- We will learn how to compute these exact probabilities in a later section, as well as probabilities for other intervals.

[Learn About Verified Certificates](#)

edX Verified Certificate

Why upgrade?

- Official proof of completion
- Easily shareable certificate
- Proven motivator to complete the course
- Certificate purchases help edX continue to offer free courses

How it works

- Pay the Verified Certificate upgrade fee
- Verify your identity with a webcam and government issued ID
- Study hard and pass the course



- Share your certificate with friends, employers, and others

edX Learner Stories



My certificate has helped me showcase my knowledge on my resume - I feel like this certificate could really help me land my dream job!

- Christina Fong, edX Learner



I wanted to include a verified certificate on my resume and my profile to illustrate that I am working towards this goal I have and that I have achieved something while I was unemployed.

- Cheryl Troell, edX Learner

Upgrade (\$49)

© All Rights Reserved

