



[Course](#) > [Section...](#) > [1.3 Qu...](#) > [Quantil...](#)

Audit Access Expires Mar 24, 2020

You lose all access to this course, including your progress, on Mar 24, 2020.

Upgrade by Feb 18, 2020 to get unlimited access to the course as long as it exists on the site. [Upgrade now](#)

Quantile-Quantile Plots

Quantile-Quantile Plots

quantile-
quantile plots

for the normal
distribution,

then it must be because
the data is approximated
by a normal distribution.

To obtain the quantiles
for the data, we can use
the quantile function

in `r` like this.

So we're going to define
an object called `observe`
quantiles, and calculate

the quantiles for `x` at the
series of values of `p`,

which is stored in the
object `p`.

To obtain the theoretical
normal distribution
quantiles

with the corresponding
average and standard
deviation,

we use the `qnorm`
function, like this.

We're going to find an



Video

[Download video file](#)

Transcripts

[Download SubRip \(.srt\) file](#)

[Download Text \(.txt\) file](#)

Textbook link

This video corresponds to the [textbook section on quantile-quantile plots](#).

Key points

- Quantile-quantile plots, or QQ-plots, are used to check whether distributions are well-approximated by a normal distribution.
- Given a proportion p , the quantile q is the value such that the proportion of values in the data below q is p .
- In a QQ-plot, the sample quantiles in the observed data are compared to the theoretical quantiles expected from the normal distribution. If the data are well-approximated by the normal distribution, then the points on the QQ-plot will fall near the identity line (sample = theoretical).
- Calculate sample quantiles (observed quantiles) using the `quantile()` function.
- Calculate theoretical quantiles with the `qnorm()` function. `qnorm()` will calculate quantiles for the standard normal distribution ($\mu = 0, \sigma = 1$) by default, but it can calculate quantiles for any normal distribution given `mean()` and `sd()` arguments. We will learn more about `qnorm()` in the probability course.
- Note that we will learn alternate ways to make QQ-plots with less code later in the series.

Code



```
# define x and z
library(tidyverse)
library(dslabs)
data(heights)
index <- heights$sex=="Male"
x <- heights$height[index]
z <- scale(x)

# proportion of data below 69.5
mean(x <= 69.5)

# calculate observed and theoretical quantiles
p <- seq(0.05, 0.95, 0.05)
observed_quantiles <- quantile(x, p)
theoretical_quantiles <- qnorm(p, mean = mean(x), sd = sd(x))

# make QQ-plot
plot(theoretical_quantiles, observed_quantiles)
abline(0,1)

# make QQ-plot with scaled values
observed_quantiles <- quantile(z, p)
theoretical_quantiles <- qnorm(p)
plot(theoretical_quantiles, observed_quantiles)
abline(0,1)
```

Learn About Verified Certificates

© All Rights Reserved

