

**Anomalous temperature and injury mortality in the USA: age-, sex- and injury-specific impacts**

Robbie M Parks<sup>1,2</sup>, James E Bennett<sup>1,2</sup>, Helen Tamura-Wicks<sup>1,2</sup>, Vasilis Kontis<sup>1,2</sup>, Ralf Toumi<sup>3</sup>, Goodarz Danaei<sup>4</sup>, Majid Ezzati<sup>1,2,5\*</sup>

<sup>1</sup>MRC-PHE Centre for Environment and Health, Imperial College London, London, United Kingdom

<sup>2</sup>Department of Epidemiology and Biostatistics, School of Public Health, Imperial College London, London, United Kingdom

<sup>3</sup>Space and Atmospheric Physics, Imperial College London, London, United Kingdom

<sup>4</sup>Harvard T.H. Chan School of Public Health, Boston, Massachusetts, USA.

<sup>5</sup>WHO Collaborating Centre on NCD Surveillance and Epidemiology, Imperial College London, London, United Kingdom

Robbie M Parks: [robbie.parks@imperial.ac.uk](mailto:robbie.parks@imperial.ac.uk)

James E Bennett: [j.e.bennett@imperial.ac.uk](mailto:j.e.bennett@imperial.ac.uk)

Helen Tamura-Wicks: [helen.tamura-wicks07@imperial.ac.uk](mailto:helen.tamura-wicks07@imperial.ac.uk)

Vasilis Kontis: [v.kontis@imperial.ac.uk](mailto:v.kontis@imperial.ac.uk)

Ralf Toumi: [r.toumi@imperial.ac.uk](mailto:r.toumi@imperial.ac.uk)

Goodarz Danaei: [gdanaei@hsph.harvard.edu](mailto:gdanaei@hsph.harvard.edu)

Majid Ezzati: [majid.ezzati@imperial.ac.uk](mailto:majid.ezzati@imperial.ac.uk)

\*corresponding author:

Majid Ezzati

Imperial College London, Norfolk Place, London W2 1PG, UK

Tel: +44 (0)20 7594 0767

Temperatures which deviate from long-term norms will be more frequent as the global climate changes, and may be associated with adverse health consequences.<sup>1-4</sup> There is limited data on how such deviations affect deaths from different injuries, especially by type of injury, month of year, age and sex. Here, we used data on mortality and temperature over a 37-year period (1980-2016) in the entire contiguous USA and formulated a Bayesian spatio-temporal model to estimate how anomalous temperatures, defined as deviations from the long-term norm of monthly temperature, affect deaths from different intentional (transport, falls and drownings) and unintentional (assault and intentional self-harm) injuries by age groups and sex. We found that a 1°C anomalously warm year would be associated with an estimated 941 (95% credible interval 831-1,053) additional injury deaths in the contiguous USA. 87% of deaths would occur in males, concentrated mostly in adolescent to middle ages. These excess deaths would comprise of increases in drowning, transport injuries, assault and intentional self-harm, offset partly by an overall decline in deaths from falls in older ages. The findings demonstrate the need for targeted public health interventions against injuries during periods of anomalously high temperatures, especially as these episodes increase with global climate change.

The potential health impacts of anthropogenic climate change are one of the key drivers for efforts to mitigate greenhouse gas emissions and for pursuing adaptation measures.<sup>3-5</sup> Current assessments of the health effects of climate change largely focus on parasitic and infectious diseases, and cardiorespiratory and other chronic diseases.<sup>2-7</sup> Less research has been conducted on injuries,<sup>8-10</sup> especially in a consistent way across injury types and demographic subgroups of the population, even though death rates from injuries vary seasonally,<sup>11,12</sup> which means that temperature may play a role in their pathogenesis. Our aim was to evaluate how deaths from

various injuries may be affected by changes in temperature that could arise as a result of global climate change in a national study.

We used vital registration data on all injury deaths in the contiguous USA from 1980 to 2016, with information on sex, age at death, underlying cause of death and state of residence. During this period, 4,006,454 boys and men and 1,757,862 girls and women died from an injury in the contiguous USA (i.e., excluding Alaska and Hawaii), accounting for 9.2% and 4.2% of all male and female deaths respectively. 95.6% of male injury deaths and 93.9% of female injury deaths were in those aged 15 years and older, and over half (52.6%) of male injury deaths were in those aged 15-44 years (Figure 1). In contrast with males, there was less of an age gradient in females after 15 years of age.

Injuries from transport, falls, drownings, assault, and intentional self-harm accounted for 79.0% of injury deaths in males and 72.1% in females. The remainder were from a heterogeneous group of “other unintentional injuries” (Figure 1), within which the type of injury that led to death varied by sex and age group. Transport was the leading injury cause of death in women younger than 75 years and men younger than 35 years. Between 35 and 74 years of age, more men died of intentional self-harm than any other injury. Above 75 years of age, falls were the largest cause of death in both men and women.

There was a decline in age-standardised death rates of three out of five major injuries (transport, drownings and assault) from 1980 to 2016, although assault deaths have shown a recent increase since 2014 (Figure 2). In contrast, age-standardised death rates from falls increased over time while those from intentional self-harm initially decreased followed by an increase to surpass 1980 levels. The largest overall decline over time was for transport deaths, which

declined by over 50% from 1980 to 2016. Age-standardised death rates for transport injuries and drownings peaked in summer months but deaths from other major injuries did not have clear seasonal patterns.

With few exceptions,<sup>8,13</sup> current climate change risk assessments typically extrapolate from changes in mortality in relation to daily temperature.<sup>1,6,7,14,15</sup> Climate change, however, will fundamentally modify weather, including seasonal weather patterns, compared to long-term norms, and hence can disrupt long-term adaptation. To mimic the conditions that may arise with global climate change, we developed methodology to examine how deviations from long-term norm temperature may impact injury death rates.

We first defined a measure of anomalous temperature for each state and month relative to long-term norm temperature of the state in that month (Figure 3). In this approach, a state with higher, but more stable, temperature in a specific month has smaller anomalies than one with lower but more inter-annually variable temperature. Average size of anomaly over the study period (1980-2016), a measure of how variable temperatures are around their central state-month long-term norm, ranged from 0.4°C for Florida in September, to 3.4°C for North Dakota in February (Figure 4). The average size of anomaly had a median value of 1.2°C across all states and months, with 27% less than 1°C and 90% less than 2°C (Figure 4). Temperature anomalies were largest in January and December and smallest in August and September. They were larger in northern and central states than in southern and coastal ones.

We then analysed the association of monthly injury death rates with anomalous temperature using a Bayesian spatio-temporal model, described in Methods. The model accounted for systematic variations in death rates across states and months, through state-, month- and state-

month-specific random intercepts, and for their long-term trends. These terms together remove the effects of space and time varying factors other than temperature that affect injuries. Analyses were done separately by injury type, sex and age group. We used the resultant risk estimates and the age-sex-specific death rates from each injury in 2016, to calculate additional deaths if each month in each state were +1°C above its long-term norm, by type of injury, sex, age group, state and month.

Based on these calculations, there would be an estimated 941 (95% credible interval 831, 1,053) excess injury deaths, equivalent to 0.47% of all injury deaths in 2016, in each year in which each month in each state were +1°C above its long-term norm (Figure 5). Deaths from drowning, transport, assault and intentional self-harm would be predicted to increase, partly offset by a decline in deaths from falls in middle and older ages and in winter months (Figure 5). Most excess deaths would be from transport injuries (448) followed by intentional self-harm (315). 87% of the excess deaths would occur in males and 13% in females. 80% of all male excess deaths would occur in those aged 15-64 years, who have higher rates of deaths from transport injuries. In those aged 85 years and older, there would be an estimated decline in injury deaths, because deaths from falls are expected to decline in a warmer year.

Proportionally, deaths from drownings are predicted to increase more than those of other injury types, by as much 8.3% (7.3, 9.3) in men aged 15-24 years (Supplementary Figure 1); the smallest proportional increase was that of assault and intentional self-harm (less than 2% in all age and sex groups). There was a larger percent increase in transport deaths for males than for females, especially in young and middle-ages (~e.g., 1.25% (0.90, 1.60) for 25-34 year old men versus 0.23% (-0.28, 0.76) for women of the same age) (Supplementary Figure 1).

While there are no previous studies of how deviations of monthly temperature from long-term norm are associated with injury mortality, our results are broadly in agreement with those that have analysed associations with absolute temperature and for specific injury types. A study of suicide in US counties over 37 years (1968-2004) estimated that 1°C higher monthly temperature would lead to a 0.7% rise in suicides,<sup>8</sup> compared to our findings of 0.44-1% in males and 0.39-1.47% in females in different ages. In a study of six French heatwaves during 1971-2003, mortality from unintentional injuries rose by up to 4% during a heatwave period compared to a non-heatwave baseline.<sup>9</sup> A study of daily mortality from all injuries from Estonia found a 1.24% increase in mortality when daily maximum temperature went from the 75<sup>th</sup> to 99<sup>th</sup> percentile of long-term distribution.<sup>10</sup>

That anomalously warm temperature influences deaths from drowning, although not previously quantified, is highly plausible because swimming is likely to be more common when monthly temperature is higher. The higher relative and absolute impacts on men compared with women may reflect differences in behaviour. For example, over half of swimming deaths for males occur in natural water, compared to about quarter for females,<sup>16</sup> which may lead to a larger rise in the former in warmer weather. Similarly, the decline in deaths from falls, which are mostly in older ages, may be because falls in older people are more likely to be due to slipping on ice than in younger ages.<sup>17-19</sup>

The pathways from anomalous temperature to transport injury are more varied. Firstly, driving performance deteriorates at higher temperatures.<sup>20-23</sup> Further, alcohol consumption increases during warm temperature anomalies,<sup>24</sup> potentially also explaining why teenagers, who are more likely than other age groups to crash while intoxicated,<sup>25</sup> experience a larger proportional rise in deaths from transport than older ages when temperatures are anomalously warm. Lastly,

warmer temperatures generally increase road traffic in North America;<sup>26–29</sup> With more people generally outdoors in warmer weather,<sup>30</sup> this could lead to more fatal collisions.

Pathways linking anomalously high temperatures and deaths from assault and self-harm are less established. One hypothesis is that, similar to transport, more time spent outdoors in anomalously warmer temperatures leads to an increased number of face-to-face interactions, and hence arguments, confrontations, and ultimately assaults.<sup>31,32</sup> These effects could be compounded by the greater anger levels linked to higher temperatures.<sup>33,34</sup> Regarding intentional self-harm, higher temperature has been hypothesised as associated with higher levels of distress in younger people.<sup>35</sup> Nonetheless, links between temperature and mental health requires further investigation,<sup>36</sup> including whether the relationship varies by age and sex, as indicated by our results.

The major strength of our study is that we have comprehensively modelled the association of temperature anomaly with injury by type of injury, month, age group and sex. Our measure of temperature anomaly internalises long-term historical experience of each state, and is closer to what climate change may bring about than solely examining daily episodes, or average temperature to which people have adapted. To utilise this metric, we integrated two large disparate national datasets on mortality (US vital statistics) and meteorology (ERA-Interim<sup>37</sup>), and developed a bespoke Bayesian spatio-temporal model. A limitation of our study is that, like all observation studies, we cannot rule out confounding of results due to other factors, although it is unlikely that such factors will have the same anomalies as temperature, even if their average space and time patterns are the same.

Our work highlights how deaths from injuries are currently susceptible to temperature anomalies and could also be modified by rising temperatures resulting from climate change, unless countered by social and health system interventions that mitigate these impacts. Though absolute impacts on mortality are modest, some groups, especially men in young to middle-ages, will experience larger impacts. Therefore, a combination of public health interventions that broadly target injuries in these groups – for example targeted messaging for younger males on the risks of transport injury and drowning – and those that trigger in relation to forecasted high temperature periods – for example more targeted blood alcohol level checks – should be a public health priority.

## **Methods**

### *Data sources*

We used data on deaths by sex, age, underlying cause of death and state of residence in the contiguous USA from 1980 to 2016 through the National Center for Health Statistics (NCHS) ([https://www.cdc.gov/nchs/nvss/dvs\\_data\\_release.htm](https://www.cdc.gov/nchs/nvss/dvs_data_release.htm)) and on population from the NCHS bridged-race dataset for 1990 to 2016 ([https://www.cdc.gov/nchs/nvss/bridged\\_race.htm](https://www.cdc.gov/nchs/nvss/bridged_race.htm)) and from the US Census Bureau prior to 1990 (<https://www.census.gov/data/tables/time-series/demo/popest/1980s-county.html>). We calculated monthly population counts through linear interpolation, assigning each yearly count to July.

The underlying cause of death was coded according to the international classification of diseases (ICD) system (9<sup>th</sup> revision from 1980 to 1998 and 10<sup>th</sup> revision thereafter). The 5.7 million injury deaths fell into six categories: transport, falls, drownings, assault, intentional self-harm and an aggregate set of other unintentional injuries. We report the results of all of these categories except other unintentional injuries (1,329,200 deaths or 23% of total injury



deaths during 1980-2016), because the composition of this aggregate group varies by sex, age group, state and time.

We obtained data on temperature from ERA-Interim, which combines predictions from a physical model with in-situ and satellite measurements.<sup>37</sup> We used gridded four-times-daily estimates at a resolution of 80 km to generate monthly population-weighted temperature by state throughout the analysis period.

#### *Anomalous temperature metric*

To calculate the magnitude of temperature anomaly by state and month, we first calculated 30-year (long-term) norm temperatures (from 1980-2009) for each month in each state. We calculated for 30 years because it is the duration used in climate assessments.<sup>38</sup> We subtracted these long-term norm temperatures from respective monthly temperature values to generate a temperature anomaly time series for each month and year in each state (Figure 3). The temperature anomaly metric measures the extent that temperature experienced in a specific month, year and state is warmer or cooler than the long-term norm to which the population of each state has acclimatised. These values can be different for different months in the same state, and different states in the same month.

#### *Statistical methods*

We formulated a Bayesian spatio-temporal model to estimate the effect of temperature anomaly on injury deaths rates. The outcome was deaths from several types of injury. We carried out all analyses separately by sex and age group (0-4 years, 10-year age groups from 5 to 84 years, and 85+ years) because injury deaths rates vary by age group and sex,<sup>11,12,39</sup> as might their associations with temperature.

We modelled the number of deaths in each year as following a Poisson distribution:

$$deaths_{state-time} \sim \text{Poisson}(death\ rate_{state-time} \cdot population_{state-time})$$

with log-transformed death rates modelled as a sum of components that depend on location (state) of death, month of year, overall time and temperature anomaly:

$$\begin{aligned} \log(deaths_{state-time}) = & \\ & \alpha_0 + \beta_0 \cdot time + \\ & \alpha_{state} + \beta_{state} \cdot time + \\ & \alpha_{month} + \beta_{month} \cdot time + \\ & \zeta_{state-month} + \\ & \nu_{time} + \\ & \gamma_{month} \cdot Anomaly_{state-time} + \\ & \epsilon_{state-time} \end{aligned}$$

The model contained terms that represent the overall level and trend in mortality, with  $\alpha_0$  as the common intercept and  $\beta_0$  the common time slope. Death rates also vary by month, which may be partly related to temperature and partly due to other monthly factors; monthly variations tend to be smooth across adjacent months.<sup>11</sup> Therefore, we allowed each month of the year to systematically have a different mortality level and trend, with  $\alpha_{month}$  the month-specific intercept and  $\beta_{month}$  the month-specific time slope. We used a random walk for the month terms to smooth the coefficients, widely used to characterise smoothly varying associations.<sup>40</sup> The random walk had a cyclic structure, so that December was adjacent to January.

We also included state random intercepts and slopes for death rates, with  $\alpha_{state}$  as the state-specific intercept and  $\beta_{state}$  the state-specific time slope. These terms measure deviations of each state from national values, and allow variation in level and trend in mortality by state. In addition, death rates in neighbouring states may be more similar than in those further away, modelled using a Conditional Autoregressive (CAR) spatial model.<sup>41</sup> This allows mortality

levels and trends of states to be estimated based on their own data as well as using those of their neighbours. The extent to which information is shared between neighbouring states depends on the uncertainty of death rates in a state and the empirical similarity of death rates in neighbouring states. We also included state-month interactions for intercepts and slopes ( $\zeta_{state-month}$ ), to allow variation in mortality levels and trends in a particular state for different months and vice-versa. Non-linear change over time was captured by a first-order national random walk,  $v_{time}$ .<sup>40</sup>

Finally, we included a term that relates log-transformed death rate to the above-defined state-month temperature anomaly,  $\gamma_{month} \cdot Anomaly_{state-time}$ . The coefficients of  $\gamma_{month}$  represent the logarithm of the monthly death rate ratio per 1°C increase in anomaly. There was a separate coefficient for each month which means that an anomaly of the same magnitude could have different associations with injury mortality in different months. As with the month-specific intercepts and trends, we used a cyclic random walk to smooth the coefficient of the temperature anomaly across months. An over-dispersion term ( $\epsilon_{state-time}$ ) captured the variation unaccounted for by other terms in the model, modelled as  $N(0, \sigma_{\epsilon}^2)$ . We fitted the models using integrated nested Laplace approximation (INLA), using the R-INLA software, which offers orders of computational efficiency improvement in Bayesian inference compared to traditional MCMC.<sup>42</sup>

We estimated the mortality impact of a national year-round temperature anomaly of 1°C in each month and state, realistic in our lifetimes under current projections of global climate change,<sup>43</sup> as well as within the range of anomaly size experienced by some states (Figure 4). For this calculation, we multiplied the actual death counts for each month, sex, state and age group in 2016 by the corresponding excess relative risk, which was calculated as the

exponential of the coefficient of the temperature anomaly term from the above analysis. The uncertainty in our results were obtained from 5000 draws from the posterior marginal of each month's excess relative risk. The reported 95% credible intervals, quoted in brackets where appropriate, are the 2.5<sup>th</sup> to 97.5<sup>th</sup> percentiles of the sampled values.

### *Sensitivity analysis*

We conducted sensitivity analyses to assess how much our results might depend on the temperature metric used to generate anomalous temperature. First, instead of building our monthly temperature anomalies based on daily mean temperatures, we used daily maxima and minima. These measures were strongly correlated to those generated from daily means (Supplementary Table 2), and therefore we did not run models using these alternatives.

Together with temperature anomaly based on daily mean temperatures, we also included a second measure of anomaly in the model. The additional measures were related to more extreme anomalous situations:

- temperature anomaly calculated based on 90<sup>th</sup> percentile (°C) of daily mean temperatures within a month, compared to 30-year (long-term) norm of 90<sup>th</sup> percentile for each state and month
- number of days in a month above the long-term 90<sup>th</sup> percentile of norm temperature for each state and month (adjusted for length of month)
- number of 3+ day episodes above the long-term 90<sup>th</sup> percentile of norm temperature for each state and month (adjusted for length of month)

The correlations among these variables and anomaly based on mean were between 0.60 and 0.89 (Supplementary Table 3). The estimated rate ratios of temperature anomaly based on daily

305 means (i.e., the anomaly measure used in the main analysis) were robust to the addition of  
306 alternative measures of anomaly, while the coefficients of the additional measures were  
307 generally not significant and with large credible intervals. Therefore, we did not include the  
308 alternative additional measures of extreme anomalous temperature in the main analysis.

## **Acknowledgments**

Robbie Parks is supported by a Wellcome Trust ISSF Studentship. The development of statistical methods is supported by grants from the Wellcome Trust (grants 205208/Z/16/Z and 209376/Z/17/Z). Work on the US mortality data is supported by a grant from US Environmental Protection Agency. This paper has not been formally reviewed by EPA. The views expressed in this document are solely those of authors and do not necessarily reflect those of the Agency. EPA does not endorse any products or commercial services mentioned in this publication.

## **Author contributions**

All authors contributed to study concept, analytical approach, and interpretation of results. RP, GD and ME collated and organised mortality files. RP performed the analysis, with input from other authors. RP and ME wrote the first draft of the paper; other authors contributed to revising and finalising the paper.

## **Competing interests statement**

ME reports a charitable grant from AstraZeneca Young Health Programme, and personal fees from Prudential, Scor, and Third Bridge, all outside the submitted work; all other authors declare no competing interests.


- 329    1.    Gasparrini, A. *et al.* Mortality risk attributable to high and low ambient temperature: A  
330    multicountry observational study. *Lancet* **386**, 369–375 (2015).
- 331    2.    Watts, N. *et al.* The 2018 report of the Lancet Countdown on health and climate  
332    change: shaping health of nations for centuries to come. *Lancet* **6736**, 1–4 (2018).
- 333    3.    McMichael, A. J., Woodruff, R. E. & Hales, S. Climate change and human health:  
334    Present and future risks. *Lancet* (2006). doi:10.1016/S0140-6736(06)68079-3
- 335    4.    Haines, A. & Ebi, K. The imperative for climate action to protect health. *N. Engl. J.*  
336    *Med.* 263–273 (2019).
- 337    5.    Smith, K. R. *et al.* Human health: Impacts, adaptation, and co-benefits. in *Climate*  
338    *Change 2014 Impacts, Adaptation and Vulnerability: Part A: Global and Sectoral*  
339    *Aspects* (2015). doi:10.1017/CBO9781107415379.016
- 340    6.    Huang, C. *et al.* Projecting future heat-related mortality under climate change  
341    scenarios: A systematic review. *Environmental Health Perspectives* (2011).  
342    doi:10.1289/ehp.1103456
- 343    7.    Gasparrini, A. *et al.* Projections of temperature-related excess mortality under climate  
344    change scenarios. *Lancet Planet. Heal.* (2017). doi:10.1016/S2542-5196(17)30156-0
- 345    8.    Burke, M. *et al.* Higher temperatures increase suicide rates in the United States and  
346    Mexico. *Nat. Clim. Chang.* (2018). doi:10.1038/s41558-018-0222-x
- 347    9.    Rey, G. *et al.* The impact of major heat waves on all-cause and cause-specific  
348    mortality in France from 1971 to 2003. *Int. Arch. Occup. Environ. Health* (2007).  
349    doi:10.1007/s00420-007-0173-4
- 350    10.    Orru, H. & Åström, D. O. Increases in external cause mortality due to high and low  
351    temperatures: evidence from northeastern Europe. *Int. J. Biometeorol.* (2017).  
352    doi:10.1007/s00484-016-1270-4
- 353    11.    Parks, R. M., Bennett, J. E., Foreman, K. J., Toumi, R. & Ezzati, M. National and  
354    regional seasonal dynamics of all-cause and cause-specific mortality in the USA from  
355    1980 to 2016. *Elife* **7**, (2018).
- 356    12.    Rau, R. Seasonality in human mortality. A demographic approach. *Wirtschafts- und*  
357    *Sozialwissenschaftlichen Fak. PhD*, 361 (2004).
- 358    13.    Shi, L., Kloog, I., Zanobetti, A., Liu, P. & Schwartz, J. D. Impacts of temperature and  
359    its variability on mortality in New England. *Nat. Clim. Chang.* **5**, 988–991 (2015).
- 360    14.    Ye, X. *et al.* Ambient temperature and morbidity: a review of epidemiological  
361    evidence. *Environ. Health Perspect.* **120**, 19–28 (2012).
- 362    15.    Basu, R. High ambient temperature and mortality: A review of epidemiologic studies  
363    from 2001 to 2008. *Environ. Heal. A Glob. Access Sci. Source* **8**, 40 (2009).
- 364    16.    Xu, J. Unintentional drowning deaths in the United States, 1999-2010. *NCHS Data*  
365    *Brief* (2014).
- 366    17.    Ambrose, A. F., Paul, G. & Hausdorff, J. M. Risk factors for falls among older adults:  
367    A review of the literature. *Maturitas* (2013). doi:10.1016/j.maturitas.2013.02.009
- 368    18.    Bobb, J. F. *et al.* Time-course of cause-specific hospital admissions during  
369    snowstorms : An analysis of electronic medical records from major hospitals in  
370    Boston, Massachusetts. *Am. J. Epidemiol.* **185**, 283–294 (2017).
- 371    19.    Kelsey, J. L. *et al.* Indoor and outdoor falls in older adults are different: The  
372    maintenance of balance, independent living, intellect, and zest in the elderly of boston  
373    study. *J. Am. Geriatr. Soc.* (2010). doi:10.1111/j.1532-5415.2010.03062.x
- 374    20.    Daanen, H. A. M., Van De Vliert, E. & Huang, X. Driving performance in cold, warm,  
375    and thermoneutral environments. *Appl. Ergon.* (2003). doi:10.1016/S0003-  
376    6870(03)00055-3

21. Zlatoper, T. J. Determinants of motor vehicle deaths in the United States: A cross-sectional analysis. *Accid. Anal. Prev.* (1991). doi:10.1016/0001-4575(91)90062-A
22. Mackie, R. R. & Hanlon, J. F. . O. A study of the combined effects of extended driving and heat stress on driver arousal and performance. in *Symposium on relationships among theory, physiological correlates, and operational performance* (1976). doi:10.1007/978-1-4684-2529-1\_25
23. Wyon, D. P., Wyon, I. & Norin, F. Effects of moderate heat stress on driver vigilance in a moving vehicle. *Ergonomics* (1996). doi:10.1080/00140139608964434
24. Opinium. Brits drink more alcohol in warmer weather. *Opinium.co.uk* (2018). Available at: <https://www.opinium.co.uk/brits-drink-more-alcohol-in-warmer-weather/>. (Accessed: 10th January 2019)
25. Voas, R. B., Torres, P., Romano, E. & Lacey, J. H. Alcohol-related risk of driver fatalities: An update using 2007 data. *J. Stud. Alcohol Drugs* (2012). doi:10.15288/jsad.2012.73.341
26. Datla, S., Sahu, P., Roh, H.-J. & Sharma, S. A comprehensive analysis of the association of highway traffic with winter weather conditions. *Procedia - Soc. Behav. Sci.* (2013). doi:10.1016/j.sbspro.2013.11.143
27. Roh, H.-J., Sahu, P. K., Sharma, S., Datla, S. & Mehran, B. Statistical investigations of snowfall and temperature interaction with passenger car and truck traffic on primary highways in Canada. *J. Cold Reg. Eng.* (2016). doi:10.1061/(ASCE)CR.1943-5495.0000099
28. Roh, H.-J., Datla, S. & Sharma, S. Effect of snow, temperature and their interaction on highway truck traffic. *J. Transp. Techn.* (2013). doi:10.4236/jtts.2013.31003
29. Roh, H. J., Sharma, S. & Sahu, P. K. Modeling snow and cold effects for classified highway traffic volumes. *KSCE J. Civ. Eng.* (2016). doi:10.1007/s12205-015-0236-0
30. Graff Zivin, J. & Neidell, M. Temperature and the allocation of time: Implications for climate change. *J. Labor Econ.* (2014). doi:10.1086/671766
31. Glaeser, E. L., Sacerdote, B. & Scheinkman, J. A. Crime and social interactions. *Q. J. Econ.* (1996). doi:10.2307/2946686
32. Rotton, J. & Cohn, E. G. Global warming and U.S. crime rates: An application of routine activity theory. *Environ. Behav.* (2003). doi:10.1177/0013916503255565
33. Anderson, C. A. Temperature and aggression: Ubiquitous effects of heat on occurrence of human violence. *Psychol. Bull.* (1989). doi:10.1037/0033-2909.106.1.74
34. Baron, R. A. & Bell, P. A. Aggression and heat: The influence of ambient temperature, negative affect, and a cooling drink on physical aggression. *J. Pers. Soc. Psychol.* (1976). doi:10.1037/0022-3514.33.3.245
35. Majeed, H. & Lee, J. The impact of climate change on youth depression and mental health. *Lancet Planet. Heal.* **1**, e94–e95 (2017).
36. Berry, H. L., Waite, T. D., Dear, K. B. G., Capon, A. G. & Murray, V. The case for systems thinking about climate change and mental health. *Nature Climate Change* (2018). doi:10.1038/s41558-018-0102-4
37. Dee, D. P. *et al.* The ERA-Interim reanalysis: configuration and performance of the data assimilation system. *Q. J. R. Meteorol. Soc.* **137**, 553–597 (2011).
38. Wallace, J. M. & Hobbs, P. V. *Atmospheric science: An introductory survey: Second edition. Atmospheric Science: An Introductory Survey: Second Edition* (2006). doi:10.1016/C2009-0-00034-8
39. Lozano, R. *et al.* Global and regional mortality from 235 causes of death for 20 age groups in 1990 and 2010: A systematic analysis for the Global Burden of Disease Study 2010. *Lancet* (2012). doi:10.1016/S0140-6736(12)61728-0
40. Rue, H. & Held, L. *Gaussian Markov random fields. Theory and applications.*




- 427 *Chapman & Hall* (2005). doi:10.1007/s00184-007-0162-3  
428 41. Besag, J. Spatial interaction and the statistical analysis of lattice systems. *J. R. Stat.*  
429 *Soc. Ser. B (Statistical Methodol.* (1974). doi:10.2307/2984812  
430 42. Rue, H., Martino, S. & Chopin, N. Approximate Bayesian inference for latent  
431 Gaussian models by using integrated nested Laplace approximations. *J. R. Stat. Soc.*  
432 *Ser. B Stat. Methodol.* (2009). doi:10.1111/j.1467-9868.2008.00700.x  
433 43. IPCC. IPCC special report on the impacts of global warming of 1.5 °C - Summary for  
434 policy makers. (2018).  
435


436 **Figure 1.** Number of injury deaths, by type of unintentional (transport, falls, drownings, and  
437 other) and intentional (assault and intentional self-harm) injury, by sex and age group in the  
438 contiguous USA for 1980-2016.




439 **Figure 2.** National age-standardised death rates from 1980 to 2016, by type of injury and  
440 month.




441 **Figure 3.** Graphic representation of temperature anomaly measure used in the analysis. The  
442 graph shows how monthly temperatures in July two example states (Florida in red and  
443 Minnesota in blue) (left panel) for 1980-2016 are used to calculate temperature anomalies. As  
444 seen, a warmer state like Florida (top right) can have a smaller inter-annual variation in a  
445 particular month (here, July) compared with a cooler state like Minnesota (bottom right).



Florida July anomalies




Minnesota July anomalies



446 **Figure 4.** Average size of temperature anomaly ( $^{\circ}\text{C}$ ) from 1980 to 2016, by state and month.  
447 The value for each state and month is the mean of the absolute size of anomaly, be it cold or  
448 warm, and hence gives an indication of the scale of anomalies around the norm local  
449 temperatures.





450 **Figure 5.** Additional annual injury deaths for the 2016 US population in year in which each  
451 month was +1°C warmer compared with 1980-2009 norm temperatures. The top row shows  
452 breakdown by type of injury, sex and age group. The bottom row shows the break down by  
453 type of injury, sex and month. Black dots represent net changes in deaths for each set of bars.

Additional deaths associated with a 1 degree warmer year (based on 2016 population)

