

Epidemiology

Long-term traffic-related air pollutant exposure and amyotrophic lateral sclerosis diagnosis in Denmark: A Bayesian hierarchical analysis --Manuscript Draft--

Manuscript Number:	EDE21-0753R3
Full Title:	Long-term traffic-related air pollutant exposure and amyotrophic lateral sclerosis diagnosis in Denmark: A Bayesian hierarchical analysis
Article Type:	Original Article
Corresponding Author:	Robbie M Parks, Ph.D. Columbia University NY, New York UNITED STATES
Corresponding Author Secondary Information:	
Corresponding Author's Institution:	Columbia University
Corresponding Author's Secondary Institution:	
First Author:	Robbie M Parks, Ph.D.
First Author Secondary Information:	
Order of Authors:	Robbie M Parks, Ph.D. Yanelli Nunez, PhD Arin A Balalian, MD, MPH Elizabeth A Gibson, PhD Johnni Hansen, PhD Ole Raaschou-Nielsen, PhD Matthias Ketzel, PhD Jibran Khan, PhD Jørgen Brandt, PhD Roel Vermeulen, PhD Susan Peters, PhD Jeff Goldsmith, PhD Diane B Re, PhD Marc G Weisskopf, PhD, ScD Marianthi-Anna Kioumourtoglou, ScD
Order of Authors Secondary Information:	
Abstract:	<p>Background: Amyotrophic lateral sclerosis (ALS) is a fatal neurodegenerative disease. Limited evidence suggests ALS diagnosis may be associated with air pollution exposure and specifically traffic-related pollutants.</p> <p>Methods: In this population-based case-control study, we used 3,937 ALS cases from the Danish National Patient Register diagnosed during 1989–2013 and matched on age, sex, year of birth and vital status to 19,333 population-based controls free of ALS at index date. We used validated predictions of elemental carbon (EC), nitrogen oxides (NOx), carbon monoxide (CO), and fine particles (PM2.5) to assign 1-, 5-, and 10-year average exposures pre-ALS diagnosis at study participants' present and historical residential addresses. We used a Bayesian hierarchical conditional logistic model, adjusting for potential confounders, to estimate individual pollutant associations, well as joint and average associations for the traffic-related pollutants (EC, NOx, CO).</p>

	<p>Results: For a standard deviation (SD) increase in 5-year average concentrations, EC (SD=0.42µg/m³) had a high probability of being individually associated with an increase in odds of ALS (11.5%; 95% credible interval[CrI]:-1.0%,25.6%; 96.3% posterior probability of a positive association), with negative associations for NO_x (SD=20µg/m³) (-4.6%;95%CrI-18.1%,8.9%; 27.8% posterior probability of a positive association), CO (SD=106µg/m³) (-3.2%;95%CrI-14.4%,10.0%; 26.7% posterior probability of a positive association) and a null association for non-EC PM_{2.5} (SD=2.37µg/m³) (0.7%;95%CrI-9.2%,12.4%). We found no association between ALS and joint or average traffic pollution concentrations.</p> <p>Conclusions: This study found a high probability of a positive association between ALS diagnosis and EC concentration. Further work is needed to understand the role of traffic-related air pollution on ALS pathogenesis.</p>
Additional Information:	
Question	Response
Word count, abstract (Enter "0" if not applicable.)	251
Word count, main text. Enter the number of words in the main text. For research articles, main text typically includes the introduction, methods, results, and discussion. Do not include words in title page, bibliography, tables, figure legends, or figures. Do not paste the text from the document itself into this box.	4154
Total word count. Enter the total word count for your submission, excluding the title page and abstract. Include the following in the total word count: (a) main text (for research articles, this typically includes the introduction, methods, results, and discussion), (b) bibliography, (c) tables, (d) figure legends, and (e) figures (calculated as 250 words per figure, each panel in a panel figure should be separately counted). Do not count words in Supplemental Digital Content. Do not paste the text from the document itself into this box.	6784
<p>Funding sources All relevant sources of funding should be included here and on the title page of the manuscript with the heading "Source of Funding."</p> <p>-Enter "none" if the work was completed without specific funding support.</p> <p>-If the result reported in the submission corresponds directly to the specific aims of a source (or sources) of funding, then describe that source of funding as: "The results reported herein correspond to specific aims of grant (or other source of support) XXX to investigator YYY from ZZZ", where XXX is a grant or project</p>	<p>Robbie M Parks was supported by the NIEHS K99 ES033742 and the Earth Institute post-doctoral research fellowship at Columbia University. Funding was also provided by the National Institute of Environmental Health Sciences (NIEHS) grants R01 ES030616, R01 ES028805, R01 AG066793, R21 ES028472, P30 ES009089, and P30 ES000002.</p>

<p>number, YYY is the Principal Investigator of the grant or project, and ZZZ is the funding agency.</p> <p>-Describe all other sources of support as: "This work was (also) supported by grant(s) (or other source of support) XXX from ZZZ", where '(also)' is inserted only if the listed support is in addition to support corresponding directly to a specific aim, XXX is a grant or project number, and ZZZ is a funding agency. Additional sources of support should be added serially (e.g., grants XXX1 from ZZZ1, XXX2 from ZZZ2, and XXX3 from ZZZ3." Sources of support can include general salary support, which may not have a grant or project number.</p> <p>Grant or project numbers should be provided in a format that allows interested parties to find the grant in publicly available databases provided by many funding agencies.</p>	
<p>Describe the process by which someone else could obtain the data and computing code required to replicate the results reported in your submission. If one or the other is not available, please write "The [data, computing code, or both] are not available for replication because" and add an explanation for why one or the other is not available. Authors of submissions without original results, such as commentaries or letters, should enter "Not applicable." Your answer to this question will not affect the editors' decision about the submission's suitability for publication. If your paper is published, your response to this question will be included in the manuscript footnotes.</p>	<p>Danish patient records are available via the Danish National Patient Register (https://econ.au.dk/the-national-centre-for-register-based-research/danish-registers/the-national-patient-register/browse). Danish population records are available via the Danish Civil Registration System (https://econ.au.dk/the-national-centre-for-register-based-research/danish-registers/the-danish-civil-registration-system-cpr/browse). Exposure data are available via the DEHM-UBM-AirGIS website (https://envs.au.dk/en/research-areas/air-pollution-emissions-and-effects/the-monitoring-program/air-pollution-models/airgis/about-airgis/). All code for analysis, results from analysis, and visualization presented in this manuscript will be publicly available via GitHub at https://github.com/rmp15/multipollutants_and_als_code_review.</p>
<p>RETAINED RIGHTS: Except for copyright, other proprietary rights related to the Work (e.g., patent or other rights to any process or procedure) shall be retained by the author. To reproduce any text, figures, tables, or illustrations from this Work in future works of their own, the author must obtain written permission from Wolters Kluwer Health, Inc. ("WKH").</p>	<p>I agree</p>

ORIGINALITY: Each author warrants that his or her submission to the Work is original, does not infringe upon, violate, or misappropriate any copyright or other intellectual property rights, or any other proprietary right, contract or other right or interest of any third party, and that he or she has full power to enter into this agreement. Neither this Work nor a similar work has been published nor shall be submitted for publication elsewhere while under consideration by this Publication.

AUTHORSHIP RESPONSIBILITY: Each author warrants that he or she has participated sufficiently in the intellectual content, the analysis of data, if applicable, and the writing of the Work to take public responsibility for it. Each has reviewed the final version of the Work, believes it represents valid work, and approves it for publication. Moreover, should the editors of the Publication request the data upon which the work is based, they shall produce it.

PREPRINTS: Upon acceptance of the article for publication, each author warrants that he/she will promptly remove any prior versions of this Work (normally a preprint) that may have been posted to an electronic server.

DISCLAIMER: Each author warrants that this Work contains no libelous or unlawful statements and does not infringe or violate the publicity or privacy rights of any third party, libel or slander any third party, contain any scandalous, obscene, or negligently prepared information, or infringe or violate any other personal or proprietary right of others. Each author

warrants that the Work does not contain any fraudulent, plagiarized or incorrectly attributed material. Each author warrants that all statements contained in the Work purporting to be facts are true, and any formula or instruction contained in the Work will not, if followed accurately, cause any injury, illness, or damage to the user. If excerpts (e.g., text, figures, tables, illustrations, or audio/video files) from copyrighted works are included, a written release will be secured by the author prior to submission, and credit to the original publication will be properly acknowledged. Each author further warrants that he or she has obtained, prior to submission, written releases from patients whose names or likenesses are submitted as part of the Work. Should the Editor or WKH request copies of such written releases, the author shall provide them in a timely manner.

DISCLOSURES/CONFLICT OF INTEREST

Each author must identify any financial interests or affiliations with institutions, organizations, or companies relevant to the manuscript by completing the form below. Additionally, any financial associations involving a spouse, partner or children must be disclosed as well.

Note: Some sections below come from the ICMJE Uniform Disclosure Form for Potential Conflicts of Interest at http://www.icmje.org/downloads/coi_disclosure.pdf (dated July 2010).

Did you or your institution at any time receive payment or support in kind for any aspect of the submitted work (including but not limited to grants, consulting fee or honorarium, support for travel to meetings for the study or other purposes, fees for participation in review activities such as data monitoring boards, statistical analysis, end point committees, and the

No

like, payment for writing or reviewing the manuscript, provision of writing assistance, medicines, equipment, or administrative support, etc...)?	
Other: Did you or your institution at any time receive additional payments or support in kind for any aspect of the submitted work?	No
Please indicate whether you have financial relationships (regardless of amount of compensation) with entities. You should report relationships that were present during the 36 months prior to submission including board membership, consultancy, employment, expert testimony, grants/grants pending, payment for lectures including service on speakers bureaus, payment for manuscript preparation, patents (planned, pending or issued), royalties, payment for development of educational presentations, stock/stock options, travel/accommodations/meeting expenses unrelated to activities listed (for example, if you report a consultancy above there is no need to report travel related to that consultancy), etc.	No
Other (err on the side of full disclosure): Please indicate whether you have any additional financial relationships (regardless of amount of compensation) with entities. You should report relationships that were present during the 36 months prior to submission.	None
Other Relationships Are there other relationships or activities that readers could perceive to have influenced, or that give the appearance of potentially influencing, what you wrote in the submitted work?	No other relationships/conditions/circumstances that present potential conflict of interest
AUTHOR'S OWN WORK: In consideration of WKH's publication of the Work, the author hereby transfers, assigns, and otherwise conveys all his/her copyright ownership worldwide, in all languages, and in all forms of media now or hereafter known, including electronic media such as CD-ROM, Internet, and Intranet, to WKH. If WKH should decide	I agree

for any reason not to publish the Work, WKH shall give prompt notice of its decision to the corresponding author, this agreement shall terminate, and neither the author nor WKH shall be under any further liability or obligation. Each author grants WKH the rights to use his or her name and biographical data (including professional affiliation) in the Work and in its or the journal's promotion.

Notwithstanding the foregoing, this paragraph shall not apply, and any transfer made pursuant to this paragraph shall be null and void if (i) the Work has been accepted by WKH for publication, and (ii) the author chooses to have the Work published by WKH as an open access publication.

WORK MADE FOR HIRE: If this Work or any element thereof has been commissioned by another person or organization, or if it has been written as part of the duties of an employee, an authorized representative of the commissioning organization or employer must also sign this form stating his or her title in the organization.

GOVERNMENT EMPLOYEES: If the Work or a portion of it has been created in the course of any author's employment by the United States Government, check the "Government" box at the end of this form. A work prepared by a government employee as part of his or her official duties is called a "work of the U.S. Government" and is not subject to copyright. If it is not prepared as part of the employee's official duties, it may be subject to copyright.

INSTITUTIONAL REVIEW

BOARD/ANIMAL CARE COMMITTEE

APPROVAL: Each author warrants that his or her institution has approved the protocol for any investigation involving humans or animals and that all experimentation was conducted in

<p>conformity with ethical and humane principles of research.</p> <p>WARRANTIES: Each author warranty made in this form is for the benefit of WKH and the Editor; each author agrees to defend, indemnify, and hold harmless those parties for any breach of such warranties.</p>	
<p>The journal will permit the author(s) to deposit for display a "final peer-reviewed manuscript" (the final manuscript after peer-review and acceptance for publication but prior to the publisher's copyediting, design, formatting, and other services) 12 months after publication of the final article on the author's personal web site, university's institutional repository or employer's intranet, subject to the following:</p> <p>* You may only deposit the final peer-reviewed manuscript.</p> <p>* You may not update the final peer-reviewed manuscript text or replace it with a proof or with the final published version.</p> <p>* You may not include the final peer-reviewed manuscript or any other version of the article on any commercial site or in any repository owned or operated by any third party. For authors of articles based on research funded by the National Institutes of Health ("NIH"), Wellcome Trust, Howard Hughes Medical Institute ("HHMI"), or other funding agency, see below for the services that WKH will provide on your behalf to comply with "Public Access Policy" guidelines.</p> <p>* You may not display the final peer-reviewed manuscript until twelve months after publication of the final article.</p>	<p>I agree</p>

<p>* You must attach the following notice to the final peer-reviewed manuscript: "This is a non-final version of an article published in final form in (provide complete journal citation)".</p> <p>* You shall provide a link in the final peer-reviewed manuscript to the journal website.</p>	
<p>"Public Access Policy" Funding Disclosure Please disclose below if you have received funding for research on which your article is based from any of the following organizations:</p>	National Institutes of Health (NIH)
<p>Please select:</p>	Author's Own Work
<p>Any additional comments?</p>	None
<p><u>Compliance with RCUK and Wellcome Trust Open Access Policies</u></p> <p>Both the Research Councils UK (RCUK) and the Wellcome Trust have adopted policies regarding Open Access to articles that have been funded by grants from the RCUK or the Wellcome Trust. If either "Wellcome Trust" or "Research Councils UK (RCUK)" has been selected above, and the authors of the applicable article choose to have the article published as an open access publication, the following policies will apply:</p> <p>* If the article is to be published pursuant to the "Gold" route of Open Access, both the RCUK and the Wellcome Trust require that WKH make the article freely available immediately pursuant to the Attribution 4.0 Creative Commons License, currently found at http://creativecommons.org/licenses/by/4.0/legalcode (the "CC BY License"). The CC BY License is the most accommodating of the Creative Commons licenses and allows others to distribute, remix,</p>	I agree

tweak, and build upon the article, even commercially, as long as they credit the authors for the original creation.

* If the article is to be published pursuant to the "Green" route of Open Access, both the RCUK and the Wellcome Trust require that WKH make the article freely available within six months pursuant to the Attribution-NonCommercial 4.0 Creative Commons License, currently found at <http://creativecommons.org/licenses/by-nc/4.0/legalcode> (the "CC BY-NC License"). The CC BY-NC License allows others to remix, tweak, and build upon the article non-commercially, and although their new works must also acknowledge the authors for the original creation and be non-commercial, they don't have to license their derivative works on the same terms.

As a service to our authors, WKH will identify the National Library of Medicine (NLM) articles that require deposit pursuant to the RCUK and Wellcome Trust policies described in this section. This Copyright Transfer Agreement provides the mechanism for identifying such articles.

WKH will transmit the final peer-reviewed manuscript of an article based on research funded in whole or in part by either RCUK or the Wellcome Trust to Pub Med Central.

Upon NIH request, it remains the legal responsibility of the author to confirm with NIH the provenance of his/her manuscript for purposes of deposit. Author will not deposit articles him/herself. Author will not alter the final peer-reviewed manuscript already transmitted to NIH.

<p>With respect to the “Green” route of Open Access, author will not authorize the display of the final peer-reviewed manuscript prior to 6 months following publication of the final article.</p> <p>Authors of articles that have been funded from grants from the RCUK or the Wellcome Trust are required to sign the WKH Open Access License Agreement prior to publication of the applicable article. Please contact the Editorial Office of the applicable journal to receive the Open Access License Agreement that is to be signed in connection with the publication of the article.</p>	
<p>I am the person in question for this submission or otherwise have approval to complete this agreement.</p>	<p>I agree</p>
<p>CME/CE Disclosure</p> <p>Each author must identify and disclose any financial associations involving a spouse, partner or children by completing the Family Disclosure question below, and whether any off-label uses or unapproved drugs or devices are discussed in his/her manuscript by completing the Off-Label Use/Unapproved Drugs or Products question below. In the event that the Work is published as a continuing education or continuing medical education article, this information will be provided to the accrediting body and may be included in the published article. When applicable, articles accepted for publication may need to comply with additional standards related to CME or CE accreditation. Please refer to guidelines for authors for details.</p> <p>WKH and its affiliates reserve the right to publish the manuscript as a continuing education article.</p>	<p>I agree</p>
<p>Family Disclosure</p>	<p>No other relationships/conditions/circumstances that present potential conflict of interest</p>

<p>Do your children or your spouse or partner have financial relationships with entities that have an interest in the content of the submitted work?</p>	
<p>Off-Label Use/Unapproved Drugs or Products</p> <p>If your manuscript discusses an unlabeled use of a commercial product or device or an investigational use of a product or device not yet approved by the FDA for any purpose, you must specifically disclose in the manuscript that the product is not labeled for the use under discussion or that the product is still investigational. Please check the item below that applies to you</p>	<p>I will not discuss unlabeled/investigational uses of any commercial product or device</p>

THE EARTH INSTITUTE
COLUMBIA UNIVERSITY

The Earth Institute
Columbia University
Hogan Hall
MC 3277
2910 Broadway
NY, NY 10025

1 June 2022

Robbie M Parks

Post-doctoral research fellow
Earth Institute, Columbia University
robbie.parks@columbia.edu
www.robbiemparks.github.io

Dr Timothy L. Lash,
Editor-in-Chief, *Epidemiology*

Dr Stefanie Ebelt
Editor, *Epidemiology*

Dear Dr Lash, Dr Ebelt:

Please find enclosed our revised manuscript entitled “Long-term traffic-related air pollutant exposure and amyotrophic lateral sclerosis diagnosis in Denmark: A Bayesian hierarchical analysis” (DE21-0753). We have revised the text according to the comments and the suggestions of the Editors and Reviewers, as outlined in our responses.

We look forward to your response and would be glad to address any further comments you may have on the manuscript.

Amyotrophic lateral sclerosis (ALS) is a devastating and fatal neurodegenerative disease, currently without a cure. Approximately half of patients die within three years of symptom onset. Annually, there are nearly 30,000 cases of ALS in Europe and over 200,000 worldwide. Known inherited genetic variants only account for 5–10% of ALS cases. Environmental factors, therefore, are likely important in ALS pathogenesis. Ambient air pollution, especially urban air pollution, is a ubiquitous exposure that has been associated with several other neurodegenerative disorders, and is consistently linked to systemic inflammation, oxidative stress, and neuroinflammation, all of which, in turn, have been reported as key pathways to ALS pathogenesis. Despite the compelling plausibility, few studies to date have evaluated the association between air pollution and ALS.

With this study, *the largest case-control study of ALS and air pollution to date*, we used data on 3,939 ALS cases from the Danish National Patient Register diagnosed between 1989 – 2013 and matched on, sex, year of birth and vital status to 19,298 population-based controls free of ALS at index date. We used predictions of nitrogen oxides (NO_x), carbon monoxide (CO), elemental carbon (EC), and fine particles (PM_{2.5}) from validated spatio-temporal models to assign 1-, 5-, and 10-year average exposures pre-ALS diagnosis at present and historical residential addresses of study participants. Although the last year of data in our analyses is from eight years ago (2013), our findings remain timely due to the uniquely large number of ALS cases we have collected, that air pollution remains so pervasive, and that ALS prevalence is projected to increase nearly 70% by 2040.

We found that an increase in the joint exposure to traffic-related pollutants was associated with an increase in odds of ALS diagnosis, significant for elemental carbon for a 1-year average standard deviation increase, though not significant at the 95% credible interval level for other pollutants. Our results indicate that sources of air pollution with elemental carbon, such as diesel engines and woodburning stoves, might contribute to development of ALS.

This manuscript has not been previously published and is not under review in any other journal. All authors have contributed to the paper, have approved its submission, and take responsibility for its contents. The authors have no actual or potential competing financial interests. There is no closely related paper included with this submission.

Sincerely,

A handwritten signature in black ink, appearing to read 'RM Parks', written in a cursive style.

Robbie M. Parks, PhD

Type of Manuscript: Original Research Article

Manuscript Title: Long-term traffic-related air pollutant exposure and amyotrophic lateral sclerosis diagnosis in Denmark: A Bayesian hierarchical analysis

Authors: Robbie M Parks^{1,2*}, Yanelli Nunez¹, Arin A Balalian³, Elizabeth A Gibson^{1,4}, Johnni Hansen⁵, Ole Raaschou-Nielsen^{5,6}, Matthias Ketzel^{6,7}, Jibran Khan⁶, Jørgen Brandt^{6,8}, Roel Vermeulen⁹, Susan Peters⁹, Jeff Goldsmith¹⁰, Diane B. Re¹, Marc G. Weisskopf¹¹, Marianthi-Anna Kioumourtoglou¹

¹Department of Environmental Health Sciences, Mailman School of Public Health, Columbia University, New York, New York, USA

²The Earth Institute, Columbia University, New York, New York, USA

³Department of Epidemiology, Mailman School of Public Health, Columbia University, New York, New York, USA

⁴Department of Biostatistics, Harvard TH Chan School of Public Health, Boston, Massachusetts, USA

⁵Danish Cancer Society Research Center, Copenhagen, Denmark

⁶Department of Environmental Science, Aarhus University, Roskilde, Denmark

⁷Global Centre for Clean Air Research (GCARE), University of Surrey, Guildford, United Kingdom

⁸iClimate – interdisciplinary Center for Climate Change, Aarhus University, Denmark

⁹Institute for Risk Assessment Sciences, Utrecht University, Utrecht, the Netherlands

¹⁰Department of Biostatistics, Mailman School of Public Health, Columbia University, New York, New York, USA

¹¹Departments of Environmental Health and Epidemiology, T. H. Chan School of Public Health, Harvard University, Boston, Massachusetts, USA

***Corresponding Author:**

Robbie M Parks

Department of Environmental Health Sciences

Columbia University Mailman School of Public Health

722 West 168th Street, #1104

New York, New York, 10032

Email: robbie.parks@columbia.edu

Suggestions for running head: Traffic-related air pollutants and ALS

The authors declare they have no actual or potential competing financial interests.

Sources of financial support: Robbie M Parks was supported by the NIEHS K99 ES033742 and the Earth Institute post-doctoral research fellowship at Columbia University. Funding was also provided by the National Institute of Environmental Health Sciences (NIEHS) grants R01 ES030616, R01 ES028805, R01 AG066793, R21 ES028472, P30 ES009089, and P30 ES000002.

Description of the process by which someone else could obtain the data and computing code required to replicate the results reported in your submission (or explanation why data or code are not available): Danish patient records are available via the Danish National Patient Register (<https://econ.au.dk/the-national-centre-for-register-based-research/danish-registers/the-national-patient-register/browse>). Danish population records are available via the Danish Civil Registration System (<https://econ.au.dk/the-national-centre-for-register-based-research/danish-registers/the-danish-civil-registration-system-cpr/browse>). Exposure data are available via the DEHM-UBM-AirGIS website (<https://envs.au.dk/en/research-areas/air-pollution-emissions-and-effects/the-monitoring-program/air-pollution-models/airgis/about-airgis/>). All code for analysis, results from analysis, and visualization presented in this manuscript will be publicly available via GitHub at https://github.com/rmp15/multipollutants_and_als_code_review.

Abstract

Background: Amyotrophic lateral sclerosis (ALS) is a fatal neurodegenerative disease. Limited evidence suggests ALS diagnosis may be associated with air pollution exposure and specifically traffic-related pollutants.

Methods: In this population-based case-control study, we used 3,937 ALS cases from the Danish National Patient Register diagnosed during 1989–2013 and matched on age, sex, year of birth and vital status to 19,333 population-based controls free of ALS at index date. We used validated predictions of elemental carbon (EC), nitrogen oxides (NO_x), carbon monoxide (CO), and fine particles (PM_{2.5}) to assign 1-, 5-, and 10-year average exposures pre-ALS diagnosis at study participants' present and historical residential addresses. We used a Bayesian hierarchical conditional logistic model, adjusting for potential confounders, to estimate individual pollutant associations, well as joint and average associations for the traffic-related pollutants (EC, NO_x, CO).

Results: For a standard deviation (SD) increase in 5-year average concentrations, EC (SD=0.42µg/m³) had a high probability of being individually associated with an increase in odds of ALS (11.5%; 95% credible interval[CrI]:-1.0%,25.6%; 96.3% posterior probability of a positive association), with negative associations for NO_x (SD=20µg/m³) (-4.6%;95%CrI-18.1%,8.9%; 27.8% posterior probability of a positive association), CO (SD=106µg/m³) (-3.2%;95%CrI-14.4%,10.0%; 26.7% posterior probability of a positive association) and a null association for non-EC PM_{2.5} (SD=2.37µg/m³) (0.7%;95%CrI-9.2%,12.4%). We found no association between ALS and joint or average traffic pollution concentrations.

Conclusions: This study found a high probability of a positive association between ALS diagnosis and EC concentration. Further work is needed to understand the role of traffic-related air pollution on ALS pathogenesis.

Abbreviations:

ALS	Amyotrophic lateral sclerosis
BKMR	Bayesian kernel machine regression
BMI	Body mass index
CO	Carbon monoxide
CrI	Credible interval
DEHM-UBM-AirGIS	Spatio-temporal air pollution modelling system used in study
EC	Elemental carbon
ICD	International Classification of Diseases
IQR	Interquartile range
IR	Incidence ratio
Non-EC PM _{2.5}	Non-elemental carbon fine particles
NO _x	Nitrogen oxides
O ₃	Ozone
PM _{2.5}	Fine particles
SD	Standard deviation
SES	Socioeconomic status

Introduction

Amyotrophic lateral sclerosis (ALS) is a devastating and fatal neurodegenerative disease,¹ currently without a cure.² Approximately half of patients die within three years of symptom onset.³ Annually, there are nearly 30,000 cases of ALS in Europe and over 200,000 worldwide.⁴ Known inherited genetic variants only account for 5–10% of ALS cases.^{5,6} Environmental factors, therefore, are likely important in ALS pathogenesis.⁷ However, because the disease is relatively rare, it is challenging to conduct large-scale prospective studies. There is a recognized need for more evidence of the environmental contributors of ALS.^{5,8}

Although air pollution is commonly studied in association with respiratory- and cardiovascular-related outcomes, e.g., references^{9–14}, epidemiologic and toxicological studies also support several plausible biological mechanisms in association with the nervous system and neurodegeneration, e.g., references^{15–34}. Ambient air pollution, especially urban air pollution, is a ubiquitous exposure that has been associated with several other neurodegenerative disorders, e.g., references^{16–21,35,36}, and is consistently linked to systemic inflammation,^{22–24} oxidative stress,^{25–28} and neuroinflammation,^{15,29} all of which, in turn, have been reported as key pathways to ALS pathogenesis, e.g., references^{30–34}.

Despite the compelling plausibility, few studies to date have evaluated the association between air pollution and ALS.^{35,37–39} A study in 2021 found that traffic-related air pollutants may be driving observed associations.³⁸ Another study of ALS and PM_{2.5} in Denmark examining critical windows of exposure found that more recent exposure to PM_{2.5} (i.e., the previous 1 to 5 years) may be the most important driver of the potential association, though the constituents of PM_{2.5} were not

analyzed, neither together nor separately.⁴⁰ No study has hitherto attempted to understand the individual, joint, and average associations of the pollutants in a single model. Air pollutants have been consistently associated with adverse health, primarily in single pollutant analyses.^{13,17,41–43} However, they are highly correlated with one another.⁴¹ It is therefore a mixture modelling challenge to infer the association of multiple air pollutants and health outcomes.⁴⁴ Using three air pollutants commonly used in health studies as traffic-related emissions tracers— elemental carbon (EC), nitrogen oxides (NO_x), and carbon monoxide (CO)— we aimed to assess whether exposure to (a) each individual air pollutant is independently associated with ALS diagnosis, and estimate their (b) joint and (c) average traffic-related emissions associations. This study pairs with and complements the work of Nunez et al.⁴⁰

Methods

Study Population and Outcome Assessment

We used data from the Danish National Patient Register during 1989-2013, through which details on demographic characteristics and certain health outcomes of all Danish residents can be linked based on a unique personal identifier.⁴⁵ The Register was established in 1977 and is comprehensive, including nationwide clinical and administrative records for all inpatient data, with outpatient data available since 1995.⁴⁶

We identified ALS cases based on their International Classification of Diseases (ICD) discharge diagnoses, i.e., ICD-8 code 348.0 (ALS) until 1993 and ICD-10 code G12.2 (motor neuron disease) thereafter, using the date of the first relevant code as the diagnosis date. This was the index date. We only included patients who were at least 20 years old when diagnosed because (i) cases

1
2
3
4 92 younger than 20 years old were at a greater chance of misclassification, since ALS has been
5
6 93 predominantly diagnosed in older adults,⁴⁷ and (ii) the very few juvenile ALS cases have been
7
8
9 94 explained to a much larger degree by genetic mutations (~40%).⁴⁸ In our validation study, Register
10
11 95 data for ALS ascertainment were highly reliable; working with a specialist ALS neurologist to
12
13 96 review medical records and comparing to death certificates and hospital discharges, the Danish
14
15
16 97 National Patient Register was found to have an overall predictive value for ALS of 82%.⁴⁹
17
18
19 98

20
21 99 We obtained controls through the Danish Civil Registration System, established in 1968 and
22
23 100 updated daily, which includes administrative records (e.g., date and place of birth, sex, vital status,
24
25
26 101 and history of civil status and addresses since 1971) on all persons living in Denmark; records are
27
28
29 102 kept even when a person dies or emigrates.⁵⁰ We randomly matched five controls per case by age,
30
31 103 sex, year of birth, and vital status. Controls were alive and free of diagnosed ALS at the ALS
32
33 104 diagnosis date of the matched case (index date). The control-sampling scheme followed a risk-set
34
35
36 105 matching pattern, so cases could have served as controls before diagnosis of ALS.⁵¹
37
38 106

39
40
41 107 We obtained all addresses of cases and controls from January 1st 1979 onwards from the Danish
42
43 108 Civil Registration System,⁵⁰ including the dates of moving to and from each address, before the
44
45
46 109 index date. We then obtained the geographical coordinates at the door of each house of the
47
48 110 residential history of the participants, with previous evidence of the high accuracy of this method
49
50
51 111 of geocoding of addresses in Denmark.¹⁷
52
53 112

54
55 113 This study was approved by the Institutional Review Board Committee at Columbia University
56
57
58 114 and the Danish Data Protection Agency.
59
60
61
62
63
64
65

Exposure data

We obtained predictions on monthly concentrations of elemental carbon (EC), nitrogen oxides (NO_x), carbon monoxide (CO), and fine particles (PM_{2.5}) (as well as ozone, O₃, for a sensitivity analysis, usually negatively correlated with other pollutants due to its chemistry⁵²), at residential addresses of study participants from the validated spatio-temporal air pollution modelling system DEHM-UBM-AirGIS that provides full space and time coverage over the study period, described in detail elsewhere.^{53–56} In brief, DEHM-UBM-AirGIS is a human exposure modelling system for traffic pollution, developed for application in Danish air pollution epidemiological studies. The modelling system integrates air pollution dispersion models, digital maps, national and local administrative databases, concentrations of air pollutants at regional, urban background and street level, meteorological data, and a Geographic Information System (GIS). The modelling system is therefore able to generate street configuration and traffic data based on digital maps and national databases, which enables estimation of air quality levels at a large number of addresses in an automatic and effective way. These predicted pollutant concentrations have been extensively used in previous air pollution epidemiologic studies in Denmark.^{17,40,57–59} The models have good predictive accuracy, with average monthly correlations between measured and modelled results of 0.79 for EC, 0.85 for NO_x, 0.91 for CO, 0.92 for O₃, and 0.83 for annual concentrations of PM_{2.5}.^{53,56} Because traffic is a major source of PM_{2.5} and EC one of the main PM_{2.5} components in urban environments,⁶⁰ we removed the EC concentration from the total PM_{2.5} mass concentration (non-EC PM_{2.5}) by subtraction to avoid overadjustment when including both in the models simultaneously; this was valid since the DEHM-UBM-AirGIS modelling system constructed PM_{2.5} concentrations by adding from specific species of pollutants, one of which was EC.^{53–56}

Based on the residential history of each case or control, we calculated 1-, 5-, and 10-year average exposure to each pollutant ending at one year before the index date, as diagnosis has been shown previously to occur at a median of 12 months after symptoms onset.⁶¹ Specifically, each case or control average value (1-, 5- or 10-year) was calculated as the mean of all concentrations recorded across time at the recorded addresses within each time window. A small number of Danish residents lack a complete address history (1.7%; lack of house number). To ensure we were including participants with adequately complete exposure records, we set the following minimum criteria for number of complete exposure record months to include cases and controls: (i) 1-year averages: 9 of 12 months, at least one measurement in each season; (ii) 5-year averages (main exposure): 30 of 60 months; and (iii) 10-year averages: 60 of 120 months.

Covariate data

We included a set of covariates based on as close as possible to index date to account for potential confounding bias, including household socioeconomic status (SES) based on last-reported job title at index date; civil status at index date, last reported place of residence at index date, and place of birth. We used a five-category individual-level SES definition developed by the Danish Institute of Social Sciences, based on job titles from income tax forms, which has been associated with ALS diagnosis in Denmark,⁶² as well as how quickly one is identified as having ALS in the Danish Civil Registration System.⁶³ Group 1 (highest status) includes corporate managers and academics; group 2: proprietors, managers of small businesses and teachers; group 3: technicians and nurses; group 4: skilled workers; and group 5: unspecialized workers, such as entry-level positions within food and retail environments. We also included a group for participants whose job title was unknown

(group 9). For each married participant, we used the higher of the couple's individual SES categories, when available. We also used information on civil status (never married, married, divorced, widowed) due to the influence that a spouse may have on visiting a family physician,⁶⁴ last reported place of residence from postcode (Greater Copenhagen, big cities of Denmark, rest of Denmark, Greenland) to account for various local environmental and behavioral stressors,⁷ and place of birth (Greater Copenhagen, big cities of Denmark, rest of Denmark, Greenland, foreign, unknown) to adjust for other potential family-specific, location-specific, and early-life confounders, which may have an impact on the probability of developing ALS.⁶⁵ Ultimately, we were limited by what was available in the Danish Civil Registration System.⁶³ As part of a sensitivity analysis, we also included parish-level SES, measured by percentage of residents with greater than high-school education, in the model. In Denmark, parishes are administrative units with an average population of ~2,500 residents.

Statistical analysis

We analyzed the association between ALS diagnosis (binary) and exposure to traffic-related pollutants by applying a Bayesian formulation of the conditional logistic model, with Bayesian hierarchy on the traffic-related pollutants (EC, NO_x, CO).^{66,67} The conditional approach examines contrasts within matched strata, i.e., groupings of case and matched controls, implicitly adjusting for matching factors (age, sex, year of birth, vital status) within each matched stratum.⁶⁶ Matching by finer scale than year of birth was not possible. Bayesian inference allows for full distributional estimation of parameters of interest.⁶⁷ We employed a Bayesian hierarchical formulation because it enables estimates of (a) independent pollutant-outcome associations, (b) a joint association of the three pollutants (i.e., total percentage change in odds of ALS diagnosis with a simultaneous

increase in each of EC, NO_x, CO), and (c) an average traffic association (i.e., average percentage change in odds of ALS diagnosis with increases in each of EC, NO_x, CO), while accounting for the variance-covariance structure between the highly-correlated exposures and their coefficients.⁶⁷ We included a linear term for each included pollutant and adjusted for individual- and parish-level SES, civil status, last reported place of residence, and place of birth.

Specifically, via a logit function, we modelled the log-odds of ALS diagnosis, as follows:

$$\begin{aligned} \text{logit}[\text{Pr}(Y_{ci} = 1)] = & \alpha_c + \\ & \beta_{EC} EC_{ci} + \beta_{NO_x} NO_{xci} + \beta_{CO} CO_{ci} + \\ & \beta_{PM_{2.5}} (\text{non-EC } PM_{2.5ci}) + \\ & \beta_{SES} SES_{ci} + \beta_{Civil\ Status} Civil\ Status_{ci} + \beta_{Residence} Residence_{ci} + \beta_{Birth} Birth_{ci}, \end{aligned}$$

where Y_{ci} denotes whether subject i in matched stratum c was diagnosed with ALS, i.e., c represents a case and its matched controls; α_c the matched stratum-specific intercepts (not estimated in conditional logistic models); β_{EC} , β_{NO_x} , β_{CO} , $\beta_{PM_{2.5}}$ the individual pollutant coefficients (log-odds) per standard deviation (SD) increase in concentration of EC, NO_x, CO, non-EC PM_{2.5} respectively, scaled by their respective SDs and centered at their means, with each β an individual pollutant association adjusted by other terms in the model and the rest as coefficients for subject-specific covariates. Interquartile Range (IQR) could equivalently be used to scale pollutant concentrations. If other sources of air pollution are associated with ALS, then including non-EC PM_{2.5} adjusts for PM_{2.5} from other sources,⁶⁸ as well as indicating whether pollution from other sources not explicitly quantified might also have associations with ALS. Therefore, $\beta_{PM_{2.5}}$ is interpreted as the association with air pollutants not specifically included in our analysis. In urban European environments, traffic-related pollutants typically represent on-

average 14% of PM_{2.5} concentrations.⁶⁹ In a sensitivity analysis, we included O₃ in the model, as O₃ concentrations have been associated with many adverse health outcomes,⁷⁰ and were negatively correlated with traffic-related pollutants, and added $ns(SES_{\text{parish}_{ct}})$, as a natural spline with three degrees of freedom.

In our model, β_{EC} , β_{NO_x} , and β_{CO} represent the independent individual pollutant associations with ALS diagnosis. In the same model, we estimated the joint association between these three pollutants and ALS diagnosis as:

$$Traffic_{\text{Joint}} = \sum_{p=EC,NO_x,CO} \beta_p p.$$

This sum quantifies the association (log-odds) with ALS of a one-SD increase in the three pollutants simultaneously.

Finally, we assumed that the traffic-related individual pollutant associations arise from a distribution of the average traffic association with ALS diagnosis. We placed a hierarchy on the traffic-specific individual pollutant terms in the model to account for the fact that the traffic-related pollutants, EC, NO_x, CO, originate from common sources and primarily traffic in urban environments:

$$\beta_{\text{Traffic}} = [\beta_{EC}, \beta_{NO_x}, \beta_{CO}],$$

$$\beta_{\text{Traffic}} \sim MVN(\mu, \Sigma),$$

$$\mu \sim N(\lambda, \sigma_\lambda),$$

$$\Sigma = \tau \Omega \tau,$$

where λ denotes the average one-SD association of traffic-related pollution with variance σ_λ . Σ , the estimated variance-covariance matrix among individual pollutant estimates, was expressed as a decomposition into a positive-definite correlation matrix Ω and scale matrix τ .⁷¹

We used weakly-informative priors so that data drove parameter estimation. Hyper-priors for coefficients on non-EC $\text{PM}_{2.5}$ and covariates were $N(0,10)$; for σ_λ and τ we used Half-Cauchy(0,10), as recommended by Gelman, Polson and Scott as a weakly-informative prior;^{72,73} Ω was defined by the weakly-informative prior $\text{LKJCorr}(1)$.⁷⁴ The exception to this was the prior for λ , the average association of traffic-related pollutants, for which estimates became unrealistically high (approaching infinity and not converging with further iterations) with a more weakly-informative prior. We therefore used a prior of $N(0,0.1)$, which did not affect estimates of other parameters. We conducted sensitivity analyses to understand the influence of priors and the robustness of the results.

We present all results as percentage change in odds of ALS diagnosis per SD increase in pollutant concentration (calculated via e.g., $e^{\beta_{\text{EC}}} - 1$, etc. obtained in the modelling process). Due to the risk-set matching pattern of our case-control study, odds ratios are also equivalently incidence ratios (IRs).⁶⁶ We ran each model with four chains with a sample size of 1,000 each, after a warm-up of 1,000 samples, for 4,000 total samples. We assessed whether the models converged by checking that the Gelman-Rubin potential scale reduction statistic⁷⁵ was below 1.1 for all estimated model parameters. The reported 95% credible intervals (CrI) are the 2.5th to 97.5th percentiles of each parameter's posterior marginal distribution. To calculate the probability that an association estimate was greater than null, we used the 4,000 samples of the posterior distribution and took

the proportion of samples which were above the null. A 50% probability means that it is as likely as not that the marginal estimate is null, a probability closer to 100% indicates that the association is more likely to be truly positive, with closer to 0% indicating more likely to be truly negative.

We conducted statistical analyses using the R Statistical Software, version 4.1.1⁷⁶ and R-STAN, version 2.21.2.⁶⁷ All code for analysis, results from analysis, and visualization presented in this manuscript is publicly available via GitHub at https://github.com/rmp15/traffic_air_pollution_als_denmark_epidemiology.

We assessed the sensitivity of our results to hyper-prior assignment; running more iterations and warm-up per chain; inclusion of O₃; single traffic-related pollutant models adjusting for non-EC PM_{2.5}; as well as adjusting by parish-level SES. From the parish-level SES sensitivity analysis we excluded those who lived in areas without parish-level SES data, namely: (i) 819 participants for 1-year average exposure; (ii) 826 participants for 5-year average exposure; and (iii) 838 participants for 10-year average exposure.

Results

After filtering the original 4,011 cases and 20,055 controls based on completeness of exposure records, we used information on 3,934 (98.1% of total) cases and 19,298 (96.2% of total) controls for 5-year average exposure. We also used 3,937 cases, 19,333 controls for 1-year average exposure and 3,929 cases, 19,250 controls for 10-year average exposure. Descriptive statistics of included cases and controls for 5-year average exposure can be found in Table 1. Descriptive statistics of controls for 5-year exposure by socioeconomic status, civil status, residence, and place of birth are

found in eTables 1-4. For the main results, we present 5-year average exposure associations as a balance between representation of most recent exposure as well as long-term concentration.

The 5-year average traffic-related pollutant concentrations were 0.85 $\mu\text{g}/\text{m}^3$ for EC (SD=0.42 $\mu\text{g}/\text{m}^3$), 27 $\mu\text{g}/\text{m}^3$ for NO_x (SD=20 $\mu\text{g}/\text{m}^3$), and 238 $\mu\text{g}/\text{m}^3$ for CO (SD=106 $\mu\text{g}/\text{m}^3$) (Table 2). Figure 1 shows Spearman correlations between pollutants for 1-, 5-, and 10-year average exposures. Traffic-related pollutants (EC, NO_x , CO) were highly correlated in cases, controls and overall, ranging from correlations of 0.91 to 0.96. Otherwise, non-EC $\text{PM}_{2.5}$ was most highly correlated with CO, ranging from 0.67 to 0.7. O_3 was negatively correlated with other pollutants, ranging from -0.54 to -0.89. Maps of average concentration of included pollutants (EC, NO_x , CO, $\text{PM}_{2.5}$, O_3) across Denmark for a representative year (2000; middle of study period 1989-2013) are also available in eFigure 1.

For 5-year average pollutant concentrations, we observed the largest overall increase in odds of ALS diagnosis for the individual SD increase in EC (11.5%; 95% CrI: -1.0%, 25.6% per 0.42 $\mu\text{g}/\text{m}^3$; 96.3% posterior probability of positive association) (Figure 2). SD increases were associated with a decrease in odds of ALS diagnosis in NO_x (-4.6%; 95% CrI: -18.1%, 8.9% per 20 $\mu\text{g}/\text{m}^3$; 27.8% posterior probability of positive association) and CO (-3.2%; 95% CrI: -14.4%, 10.0% per 106 $\mu\text{g}/\text{m}^3$; 26.7% posterior probability of positive association). Non-EC $\text{PM}_{2.5}$ was not associated with ALS diagnosis (0.7%; 95% CrI: -9.2%, 12.4% per 2.37 $\mu\text{g}/\text{m}^3$; 54.1% posterior probability of positive association). 1-year EC average exposure was associated with an increase in odds of ALS diagnosis (15.4%; 95% CrI: 1.6%, 25.6% per 0.42 $\mu\text{g}/\text{m}^3$; 98.9% posterior probability of positive association). Single-pollutant models for each traffic-related pollutant

adjusting for non-EC PM_{2.5} (eFigure 1; single traffic-related pollutant models D, E and F) resulted in positive associations for each of EC, NO_x, CO, with positive associations for non-EC PM_{2.5} in all but the model with EC. The 95% credible interval for EC in the single-pollutant model (eFigure 1; model F) overlapped with the credible intervals of the EC term in the multi-pollutant models (eFigure 1; models A to C, G to P). The joint association of traffic-related pollutants (EC, NO_x, CO) was 2.3% (95% CrI: -3.3%, 7.7%), with an 77.8% posterior probability of a positive association. The average traffic association was null (-0.1%; 95% CrI: -17.4%, 20.8%; 45.5% posterior probability of positive association). Compared to the 1- and 5-year results, the 10-year average exposure results were attenuated, as associations tended further to the null. Results from variations of the main model in the sensitivity analyses were robust to prior choices, inclusion of O₃, and inclusion of parish-level SES (eFigure 2).

Discussion

In the largest case-control study of ALS and traffic-related air pollution to date, we found that EC had the largest-in-magnitude independent association with ALS diagnosis, while associations with NO_x and CO were negative with credible intervals overlapping the null, and smaller in magnitude. A joint increase in concentrations of traffic-related pollutants had a high probability of being associated with an increase in odds of ALS diagnosis. Sensitivity analyses demonstrated that for single pollutant models, the association for EC was smaller than for our main multi-pollutant model, which took into account the variance-covariance structure of traffic-related pollutants. Overall conclusions for the association between EC and ALS diagnosis were similar from the single- or multi-pollutant models. The inconsistent associations for NO_x and CO in the multi- and single-pollutant models and the consistency of the EC association suggest that EC concentrations

may have been more relevant than NO_x and CO for ALS diagnosis, though further study is required. Our results indicate that traffic-related pollutants, hazardous in many ways,^{9–21,41–43} may also be associated with ALS diagnosis. Our finding—that increases in EC, are potentially positively associated with ALS diagnosis—is plausible. A case-control study in the Netherlands from 2021 reported that ultrafine particles, another traffic emissions-related surrogate, were associated with ALS diagnosis,³⁸ while another based in Catalonia, Spain found ALS cases clustered around key road infrastructure.⁷⁷ Although we did not find an association with non-EC PM_{2.5} in our study, our results are not directly comparable to those of the other studies, as our PM_{2.5} effect estimates capture the PM_{2.5} components not accounted for by other pollutants in the analysis. A study examining critical windows of exposure of PM_{2.5} and ALS diagnosis in Denmark found that concentrations 1 to 5 years before exposure may be driving the association with ALS onset,⁴⁰ consistent with our findings that the most recent 1-year average EC concentration exhibited the largest association.

Our results indicate that EC exposure—a large part of which comes from diesel combustion and small combustion sources (such as wood stoves) in European urban centers, where prevalence of diesel cars is high⁷⁸—has a high probability of a positive association with ALS diagnosis. In our previous study of ALS and occupational exposures in Denmark we found that those working in agriculture and construction, associated with exposure to diesel engine exhausts, were at higher relative risk than those in other employments.⁶² Truck drivers, for whom diesel exposure is common, are also at increased risk of sporadic ALS.⁷⁹ EC exposure has been associated with inflammation,⁸⁰ mitochondrial dysfunction⁸¹ and DNA damage,^{81,82} all of which are plausible

pathways of neurodegeneration. These factors have also previously been identified as particular pathways to pathogenesis of ALS.^{30–34}

We did not find a high probability of a positive association with NO_x in our analyses, in contrast with a previous study, though that study did not include EC.³⁸ NO_x is also highly correlated with EC (0.94 to 0.96 in our study), which is expected given that they are both combustion products commonly associated with emissions in urban environments. EC exposure was more strongly associated with 1-year than for 5-/10-year average concentrations, which may indicate that the previous year of exposure may be the most relevant exposure window relevant to traffic-related exposures and ALS; this is biologically plausible, as this critical exposure window would be at the pre-symptomatic stage of underlying ALS progression, where traffic-related pollution exposure may add to the ongoing cellular or molecular process of the disease, to the point where the body can no longer compensate and subsequently enters the clinical phase.^{83–85} We do not expect that these results are attributed to reverse causation, as we have lagged these 1-year exposures by one year already prior to diagnosis, and there was likely little substantial residential movement in the year before ALS diagnosis.⁸⁶ We do not expect that calendar time was a potential source of confounding, as the controls were matched on age and year of birth. The null joint association, combined with the largest associations from traffic-related pollutant in all models found with EC, further indicates that EC may be driving the association of air pollution with ALS, though further analysis will be necessary to confirm this.

Our study used one the largest number of ALS patients ever included in an environmental health study. Another strength of our study is that we leveraged highly correlated traffic pollutants and

Bayesian hierarchical modeling and were able to estimate independent, joint, and average traffic-related pollutant associations. Although we have adjusted implicitly (by matching; age, sex, year of birth, vital status) and explicitly for many common covariates (SES, civil status, residence, place of birth), we cannot rule out residual confounding. Information on individual-level variables, such as body mass index (BMI) and smoking status is not currently available through the Danish Civil Registration System. These variables, while potential risk factors for ALS, are not likely confounders in this analysis as they are not expected to be associated with pollutant concentrations in a manner independent of neighborhood SES. If this information were available, it could be used to further adjust for SES.⁸⁶ To the extent that the variables we included in our models to adjust for household- and neighborhood-level SES are adequate, we would expect any residual SES-related confounding to be minimal. Exposure measurement error is inevitable, as any modelled exposure will be inaccurate to some degree. However, any error is not likely correlated with ALS diagnosis, and therefore any bias would be towards null.⁸⁷ While a previous study found that ALS ascertainment from the Danish National Patient Register was highly reliable,⁴⁹ outcome misclassification cannot be ruled out, nor can the possibility that date of diagnosis and symptom onset were irregularly aligned. While our analysis adjusted for marital status and household SES, many couples in Denmark cohabit. This would not be captured by our analysis, and ALS diagnosis in relation to cohabitation status should be further investigated.⁸⁸

Future research might use larger cohort data to understand the importance of each respective pollutant in a single model. Other mixture model methods, such as Bayesian Kernel Machine Regression (BKMR)⁸⁹ might be useful in further exploring the robustness of joint associations in a different framework, though BKMR was not appropriate for our particular research question,

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

388 since BKMR is currently not available for case-control study applications. The timing of exposure
389 will continue to be an important study route. ALS is projected to increase in prevalence over the
390 next few decades all over the world.⁴ Understanding ALS pathogenesis and identifying modifiable
391 risk factors is critical for preventive action.

References

1. Rowland LP, Shneider NA. Amyotrophic lateral sclerosis. *New England Journal of Medicine*. 2001;344(22):1688-1700.
2. Chio A, Logroscino G, Hardiman O, et al. Prognostic factors in ALS: A critical review. *Amyotrophic Lateral Sclerosis*. 2009;10(5-6):310-323.
3. Mitchell JD, Borasio GD. Amyotrophic lateral sclerosis. *The Lancet*. 2007;369(9578):2031-2041.
4. Arthur KC, Calvo A, Price TR, Geiger JT, Chio A, Traynor BJ. Projected increase in amyotrophic lateral sclerosis from 2015 to 2040. *Nature Communications*. 2016;7(1):1-6.
5. Al-Chalabi A, Hardiman O. The epidemiology of ALS: A conspiracy of genes, environment and time. *Nature Reviews Neurology*. 2013;9(11):617-628.
6. Hardiman O, Al-Chalabi A, Chio A, et al. Amyotrophic lateral sclerosis. *Nature reviews Disease primers*. 2017;3(1):1-19.
7. Oskarsson B, Horton DK, Mitsumoto H. Potential environmental factors in amyotrophic lateral sclerosis. *Neurologic Clinics*. 2015;33(4):877-888.
8. Longinetti E, Fang F. Epidemiology of amyotrophic lateral sclerosis: An update of recent literature. *Current Opinion In Neurology*. 2019;32(5):771.
9. Dominici F, Peng RD, Bell ML, et al. Fine particulate air pollution and hospital admission for cardiovascular and respiratory diseases. *JAMA*. 2006;295(10):1127-1134.
10. Bennett JE, Tamura-Wicks H, Parks RM, et al. Particulate matter air pollution and national and county life expectancy loss in the USA: A spatiotemporal analysis. *PLOS Medicine*. 2019;16(7):e1002856. doi:10.1371/journal.pmed.1002856
11. Schwartz J. Particulate air pollution and chronic respiratory disease. *Environmental Research*. 1993;62(1):7-13.
12. Schwartz J. The distributed lag between air pollution and daily deaths. *Epidemiology*. 2000;11(3):320-326.
13. Brook RD, Rajagopalan S, Pope III CA, et al. Particulate matter air pollution and cardiovascular disease: An update to the scientific statement from the American Heart Association. *Circulation*. 2010;121(21):2331-2378.
14. Dockery DW, Pope CA, Xu X, et al. An association between air pollution and mortality in six U.S. cities. *New England Journal of Medicine*. 1993;329(24):1753-1759. doi:10.1056/NEJM199312093292401

15. Block ML, Elder A, Auten RL, et al. The outdoor air pollution and brain health workshop. *Neurotoxicology*. 2012;33(5):972-984.
16. Zanobetti A, Dominici F, Wang Y, Schwartz JD. A national case-crossover analysis of the short-term effect of PM 2.5 on hospitalizations and mortality in subjects with diabetes and neurological disorders. *Environmental Health*. 2014;13(1):1-11.
17. Ritz B, Lee PC, Hansen J, et al. Traffic-related air pollution and Parkinson's disease in Denmark: A case-control study. *Environmental Health Perspectives*. 2016;124(3):351-356.
18. Kioumourtzoglou MA, Schwartz JD, Weisskopf MG, et al. Long-term PM2.5 exposure and neurological hospital admissions in the northeastern United States. *Environmental health perspectives*. 2016;124(1):23-29.
19. Levesque S, Surace MJ, McDonald J, Block ML. Air pollution & the brain: Subchronic diesel exhaust exposure causes neuroinflammation and elevates early markers of neurodegenerative disease. *Journal of Neuroinflammation*. 2011;8(1):1-10.
20. Heusinkveld HJ, Wahle T, Campbell A, et al. Neurodegenerative and neurological disorders by small inhaled particles. *Neurotoxicology*. 2016;56:94-106.
21. Power MC, Weisskopf MG, Alexeeff SE, Coull BA, Spiro III A, Schwartz J. Traffic-related air pollution and cognitive function in a cohort of older men. *Environmental Health Perspectives*. 2011;119(5):682-687.
22. Dubowsky SD, Suh H, Schwartz J, Coull BA, Gold DR. Diabetes, obesity, and hypertension may enhance associations between air pollution and markers of systemic inflammation. *Environmental Health Perspectives*. 2006;114(7):992-998.
23. Ruckerl R, Ibal-Mulli A, Koenig W, et al. Air pollution and markers of inflammation and coagulation in patients with coronary heart disease. *American Journal of Respiratory and Critical Care Medicine*. 2006;173(4):432-441.
24. Hoffmann B, Moebus S, Dragano N, et al. Chronic residential exposure to particulate matter air pollution and systemic inflammatory markers. *Environmental Health Perspectives*. 2009;117(8):1302-1308.
25. Kelly FJ. Oxidative stress: Its role in air pollution and adverse health effects. *Occupational and Environmental Medicine*. 2003;60(8):612-616.
26. Chuang KJ, Chan CC, Su TC, Lee CT, Tang CS. The effect of urban air pollution on inflammation, oxidative stress, coagulation, and autonomic dysfunction in young adults. *American journal of respiratory and critical care medicine*. 2007;176(4):370-376.
27. Li N, Sioutas C, Cho A, et al. Ultrafine particulate pollutants induce oxidative stress and mitochondrial damage. *Environmental Health Perspectives*. 2003;111(4):455-460.

28. Sørensen M, Daneshvar B, Hansen M, et al. Personal PM2.5 exposure and markers of oxidative stress in blood. *Environmental health perspectives*. 2003;111(2):161-166.
29. Block ML, Calderón-Garcidueñas L. Air pollution: Mechanisms of neuroinflammation and CNS disease. *Trends in neurosciences*. 2009;32(9):506-516.
30. Perry VH, Cunningham C, Holmes C. Systemic infections and inflammation affect chronic neurodegeneration. *Nature Reviews Immunology*. 2007;7(2):161-167.
31. Bergeron C. Oxidative stress: Its role in the pathogenesis of amyotrophic lateral sclerosis. *Journal of the neurological sciences*. 1995;129:81-84.
32. Mhatre M, Floyd RA, Hensley K. Oxidative stress and neuroinflammation in Alzheimer's disease and amyotrophic lateral sclerosis: Common links and potential therapeutic targets. *Journal of Alzheimer's disease*. 2004;6(2):147-157.
33. D'Amico E, Factor-Litvak P, Santella RM, Mitsumoto H. Clinical perspective on oxidative stress in sporadic amyotrophic lateral sclerosis. *Free radical biology and medicine*. 2013;65:509-527.
34. Perry VH, Nicoll JA, Holmes C. Microglia in neurodegenerative disease. *Nature Reviews Neurology*. 2010;6(4):193-201.
35. Nunez Y, Boehme AK, Weisskopf MG, et al. Fine particle exposure and clinical aggravation in neurodegenerative diseases in New York State. *Environmental Health Perspectives*. 2021;129(2):027003.
36. Nunez Y, Boehme AK, Li M, et al. Parkinson's disease aggravation in association with fine particle components in New York State. *Environmental Research*. 2021;201:111554.
37. Malek AM, Barchowsky A, Bowser R, et al. Exposure to hazardous air pollutants and the risk of amyotrophic lateral sclerosis. *Environmental Pollution*. 2015;197:181-186.
38. Yu Z, Peters S, van BL, et al. Long-Term Exposure to Ultrafine Particles and Particulate Matter Constituents and the Risk of Amyotrophic Lateral Sclerosis. *Environmental Health Perspectives*. 2021;129(9):097702. doi:10.1289/EHP9131
39. Seelen M, Toro CRA, Veldink JH, et al. Long-term air pollution exposure and amyotrophic lateral sclerosis in Netherlands: A population-based case-control study. *Environmental Health Perspectives*. 2017;125(9):097023. doi:10.1289/EHP1115
40. Nunez Y, Balalian A, Parks RM, et al. Exploring relevant time windows in the association between PM2.5 exposure and Amyotrophic Lateral Sclerosis: A case-control study in Denmark. *American Journal of Epidemiology*. Published online Under Revision.
41. Strak M, Weinmayr G, Rodopoulou S, et al. Long term exposure to low level air pollution and mortality in eight European cohorts within the ELAPSE project: Pooled analysis. *BMJ*. 2021;374:n1904. doi:10.1136/bmj.n1904

42. Hamra GB, Laden F, Cohen AJ, Raaschou-Nielsen O, Brauer M, Loomis D. Lung cancer and exposure to nitrogen dioxide and traffic: A systematic review and meta-analysis. *Environmental Health Perspectives*. 2015;123(11):1107-1112.
43. Chen H, Kwong JC, Copes R, et al. Living near major roads and the incidence of dementia, Parkinson's disease, and multiple sclerosis: A population-based cohort study. *The Lancet*. 2017;389(10070):718-726.
44. Gibson EA, Nunez Y, Abuawad A, et al. An overview of methods to address distinct research questions on environmental mixtures: An application to persistent organic pollutants and leukocyte telomere length. *Environmental Health*. 2019;18(1):1-16.
45. Frank L. When an entire country is a cohort. *Science*. 2000;287(5462):2398-2399.
46. Schmidt M, Schmidt SAJ, Sandegaard JL, Ehrenstein V, Pedersen L, Sørensen HT. The Danish National Patient Registry: A review of content, data quality, and research potential. *Clinical epidemiology*. 2015;7:449.
47. Trabjerg BB, Garton FC, van Rheenen W, et al. ALS in Danish registries: heritability and links to psychiatric and cardiovascular disorders. *Neurology Genetics*. 2020;6(2).
48. Mathis S, Goizet C, Soulages A, Vallat JM, Le Masson G. Genetics of amyotrophic lateral sclerosis: A review. *Journal of the Neurological Sciences*. 2019;399:217-226.
49. Kioumourtzoglou MA, Seals RM, Himmelslev L, Gredal O, Hansen J, Weisskopf MG. Comparison of diagnoses of amyotrophic lateral sclerosis by use of death certificates and hospital discharge data in the Danish population. *Amyotrophic Lateral Sclerosis and Frontotemporal Degeneration*. 2015;16(3-4):224-229.
50. Pedersen CB. The Danish civil registration system. *Scandinavian journal of public health*. 2011;39(7_suppl):22-25.
51. Langholz B, Goldstein L. Risk set sampling in epidemiologic cohort studies. *Statistical Science*. Published online 1996:35-53.
52. Sillman S. The relation between ozone, NOx and hydrocarbons in urban and polluted rural environments. *Atmospheric Environment*. 1999;33(12):1821-1845.
53. Khan J, Kakosimos K, Raaschou-Nielsen O, et al. Development and performance evaluation of new AirGIS—a GIS based air pollution and human exposure modelling system. *Atmospheric environment*. 2019;198:102-121.
54. Brandt J, Christensen JH, Frohn LM, Palmgren F, Berkowicz R, Zlatev Z. Operational air pollution forecasts from European to local scale. *Atmospheric Environment*. 2001;35:S91-S98.

55. Brandt J, Christensen J, Frohn L, Berkowicz R. Air pollution forecasting from regional to urban street scale—implementation and validation for two cities in Denmark. *Physics and Chemistry of the Earth, Parts A/B/C*. 2003;28(8):335-344.
56. Frohn LM, Ketzel M, Christensen JH, et al. Modelling ultrafine particle number concentrations at address resolution in Denmark from 1979-2018—Part 1: Regional and urban scale modelling and evaluation. *Atmospheric Environment*. 2021;264:118631.
57. Raaschou-Nielsen O, Andersen ZJ, Hvidberg M, et al. Lung cancer incidence and long-term exposure to air pollution from traffic. *Environmental health perspectives*. 2011;119(6):860-865.
58. Raaschou-Nielsen O, Sørensen M, Ketzel M, et al. Long-term exposure to traffic-related air pollution and diabetes-associated mortality: A cohort study. *Diabetologia*. 2013;56(1):36-46.
59. Sørensen M, Hoffmann B, Hvidberg M, et al. Long-term exposure to traffic-related air pollution associated with blood pressure and self-reported hypertension in a Danish cohort. *Environmental health perspectives*. 2012;120(3):418-424.
60. Seinfeld J, Pandis S. Atmospheric chemistry and physics. 1997. New York. Published online 2008.
61. Galvin M, Gaffney R, Corr B, Mays I, Hardiman O. From first symptoms to diagnosis of amyotrophic lateral sclerosis: Perspectives of an Irish informal caregiver cohort—a thematic analysis. *BMJ Open*. 2017;7(3). doi:10.1136/bmjopen-2016-014985
62. Dickerson AS, Hansen J, Kioumourtzoglou MA, Specht AJ, Gredal O, Weisskopf MG. Study of occupation and amyotrophic lateral sclerosis in a Danish cohort. *Occup Environ Med*. 2018;75(9):630-638. doi:10.1136/oemed-2018-105110
63. Roberts AL, Johnson NJ, Chen JT, Cudkowicz ME, Weisskopf MG. Race/ethnicity, socioeconomic status, and ALS mortality in the United States. *Neurology*. 2016;87(22):2300-2308.
64. Bucher BT, Shi J, Pettit RJ, Ferraro J, Chapman WW, Gundlapalli A. Determination of marital status of patients from structured and unstructured electronic healthcare data. In: *AMIA Annual Symposium Proceedings*. Vol 2019. American Medical Informatics Association; 2019:267.
65. Norman RE, Carpenter DO, Scott J, Brune MN, Sly PD. Environmental exposures: an underrecognized contribution to noncommunicable diseases. *Reviews on environmental health*. 2013;28(1):59-65.
66. Rothman KJ, Greenland S, Lash TL, others. *Modern Epidemiology*. Vol 3. Wolters Kluwer Health/Lippincott Williams & Wilkins Philadelphia; 2008.

67. Gelman A, Carlin JB, Stern HS, Dunson DB, Vehtari A, Rubin DB. *Bayesian Data Analysis, Third Edition*. CRC Press; 2013.
68. Mostofsky E, Schwartz J, Coull BA, et al. Modeling the association between particle constituents of air pollution and health outcomes. *American journal of epidemiology*. 2012;176(4):317-326.
69. Thunis P, Degraeuwe B, Pisoni E, et al. PM2.5 source allocation in European cities: A SHERPA modelling study. *Atmospheric Environment*. 2018;187:93-106.
70. Nuvolone D, Petri D, Voller F. The effects of ozone on human health. *Environmental Science and Pollution Research*. 2018;25(9):8074-8088.
71. Martin R, Peters G, Wilkinson J. Symmetric decomposition of a positive definite matrix. *Numerische Mathematik*. 1965;7(5):362-383.
72. Polson NG, Scott JG. On the half-Cauchy prior for a global scale parameter. *Bayesian Analysis*. 2012;7(4):887-902.
73. Gelman A. Prior distributions for variance parameters in hierarchical models (comment on article by Browne and Draper). *Bayesian Anal*. 2006;1(3):515-534. doi:10.1214/06-BA117A
74. Lewandowski D, Kurowicka D, Joe H. Generating random correlation matrices based on vines and extended onion method. *Journal of multivariate analysis*. 2009;100(9):1989-2001.
75. Gelman A, Rubin DB. Inference from iterative simulation using multiple sequences. *Statistical science*. 1992;7(4):457-472.
76. R Core Team. R: A language and environment for statistical computing. Published online 2013.
77. Povedano M, Saez M, Martinez-Matos JA, Barceló MA. Spatial assessment of the association between long-term exposure to environmental factors and the occurrence of amyotrophic lateral sclerosis in Catalonia, Spain: A population-based nested case-control study. *Neuroepidemiology*. 2018;51(1-2):33-49.
78. von Schneidemesser E, Mar KA, Saar D. Black carbon in Europe: Targeting an air Pollutant and climate forcer. Published online 2017.
79. Pamphlett R, Rikard-Bell A. Different occupations associated with amyotrophic lateral sclerosis: Is diesel exhaust the link? *PloS One*. 2013;8(11):e80993.
80. Zhang R, Dai Y, Zhang X, et al. Reduced pulmonary function and increased pro-inflammatory cytokines in nanoscale carbon black-exposed workers. *Part Fibre Toxicol*. 2014;11:73. doi:10.1186/s12989-014-0073-1

81. Gao X, Xu H, Shang J, et al. Ozonized carbon black induces mitochondrial dysfunction and DNA damage. *Environ Toxicol*. 2017;32(3):944-955. doi:10.1002/tox.22295
82. Kyjovska ZO, Jacobsen NR, Saber AT, et al. DNA damage following pulmonary exposure by instillation to low doses of carbon black (Printex 90) nanoparticles in mice. *Environ Mol Mutagen*. 2015;56(1):41-49. doi:10.1002/em.21888
83. Benatar M, Turner MR, Wu J. Defining pre-symptomatic amyotrophic lateral sclerosis. *Amyotrophic Lateral Sclerosis and Frontotemporal Degeneration*. 2019;20(5-6):303-309.
84. Benatar M, Wu J, McHutchison C, et al. Preventing amyotrophic lateral sclerosis: insights from pre-symptomatic neurodegenerative diseases. Published online 2021.
85. Eisen A, Kiernan M, Mitsumoto H, Swash M. Amyotrophic lateral sclerosis: A long preclinical period? *Journal of Neurology, Neurosurgery & Psychiatry*. 2014;85(11):1232-1238.
86. Weisskopf MG, Webster TF. Trade-offs of personal vs. more proxy exposure measures in environmental epidemiology. *Epidemiology (Cambridge, Mass)*. 2017;28(5):635.
87. Carroll RJ, Ruppert D, Stefanski LA, Crainiceanu CM. *Measurement Error in Nonlinear Models: A Modern Perspective*. CRC press; 2006.
88. Frisch M, Simonsen J. Marriage, cohabitation and mortality in Denmark: national cohort study of 6.5 million persons followed for up to three decades (1982–2011). *International Journal of Epidemiology*. 2013;42(2):559-578.
89. Bobb JF, Valeri L, Claus Henn B, et al. Bayesian kernel machine regression for estimating the health effects of multi-pollutant mixtures. *Biostatistics*. 2015;16(3):493-508.

Table 1. Demographic characteristics of cases and controls for 5-year average exposure group.

Characteristic	Overall, N = 23,232 ^a	Case, N = 3,934 ^a	Control, N = 19,298 ^a
Average age (years)	66 (12)	66 (12)	66 (12)
Sex			
Female	10,973 (47%)	1,854 (47%)	9,119 (47%)
Male	12,259 (53%)	2,080 (53%)	10,179 (53%)
Socioeconomic status (SES)			
Group 1 (Highest)	2,337 (10%)	451 (11%)	1,886 (9.8%)
Group 2	2,839 (12%)	499 (13%)	2,340 (12%)
Group 3	4,360 (19%)	785 (20%)	3,575 (19%)
Group 4	6,598 (28%)	1,076 (27%)	5,522 (29%)
Group 5 (Lowest)	4,419 (19%)	717 (18%)	3,702 (19%)
Group 9 (Unknown)	2,679 (12%)	406 (10%)	2,273 (12%)
Place of birth			
Greater Copenhagen	4,858 (21%)	831 (21%)	4,027 (21%)
Big cities of Denmark	7,923 (34%)	1,357 (34%)	6,566 (34%)
Rest of Denmark	9,009 (39%)	1,548 (39%)	7,461 (39%)
Greenland	243 (1.0%)	53 (1.3%)	190 (1.0%)
Foreign	1,065 (4.6%)	122 (3.1%)	943 (4.9%)
Unknown	134 (0.6%)	23 (0.6%)	111 (0.6%)
Civil status			
Married	14,158 (61%)	2,411 (61%)	11,747 (61%)
Divorced	2,703 (12%)	433 (11%)	2,270 (12%)
Widowed	4,224 (18%)	726 (18%)	3,498 (18%)
Never married	2,147 (9.2%)	364 (9.3%)	1,783 (9.2%)
Last reported place of residence			
Greater Copenhagen	1,887 (8.1%)	335 (8.5%)	1,552 (8.0%)
Big cities of Denmark	9,385 (40%)	1,590 (40%)	7,795 (40%)
Rest of Denmark	11,954 (51%)	2,008 (51%)	9,946 (52%)
Greenland	6 (<0.1%)	1 (<0.1%)	5 (<0.1%)

^aMean (SD); n (%)

Table 2. Summary of 1-, 5-, and 10-year average pollutant concentrations (all in $\mu\text{g}/\text{m}^3$).

	Pollutant	Overall, N = 23,270 ^a	Case, N = 3,937 ^a	Control, N = 19,333 ^a
1-year average	EC	0.81 (0.42)	0.83 (0.44)	0.81 (0.42)
	NO _x	26 (19)	26 (20)	26 (19)
	CO	224 (97)	226 (101)	224 (96)
	non-EC PM _{2.5}	11.17 (2.32)	11.20 (2.34)	11.17 (2.31)
	O ₃	52.6 (6.1)	52.4 (6.2)	52.6 (6.1)
	Pollutant	Overall, N = 23,232 ^a	Case, N = 3,934 ^a	Control, N = 19,298 ^a
5-year average	EC	0.85 (0.42)	0.86 (0.45)	0.85 (0.42)
	NO _x	27 (20)	28 (21)	27 (20)
	CO	238 (106)	239 (112)	237 (105)
	non-EC PM _{2.5}	11.76 (2.37)	11.78 (2.41)	11.76 (2.37)
	O ₃	51.9 (6.0)	51.9 (6.1)	52.0 (6.0)
	Pollutant	Overall, N = 23,179 ^a	Case, N = 3,929 ^a	Control, N = 19,250 ^a
10-year average	EC	0.89 (0.43)	0.89 (0.46)	0.88 (0.43)
	NO _x	29 (20)	29 (22)	29 (20)
	CO	253 (115)	255 (122)	253 (113)
	non-EC PM _{2.5}	12.53 (2.55)	12.55 (2.59)	12.52 (2.55)
	O ₃	51.3 (6.0)	51.3 (6.1)	51.4 (6.0)

^aMean (SD)


Figure Captions

Figure 1. Spearman correlation of 1-, 5-, and 10-year average pollutant concentrations.

Figure 2. Percentage change in odds of ALS diagnosis per 1-, 5- and 10-year average standard deviation (SD) increase for each pollutant. Results are from the Bayesian hierarchical model including each of EC, NO_x CO, and non-EC PM_{2.5} together, and were additionally adjusted by age, sex, year of birth, vital status, socioeconomic status, civil status, last reported place of residence, and place of birth.


Figure 1

Overall




1-year average

Cases




Controls




5-year average


Overall



Cases




Controls




10-year average

Overall



Cases



Controls







Figure 2





[Click here to access/download](#)

Supplemental Digital Content
supplemental digital content.pdf



Comments from the editors

We thank the Editors and Reviewers for their thoughtful and constructive suggestions. We have revised the manuscript in response to their comments, as detailed below.

All page/line/reference numbers refer to the clean revised manuscript.

Comments from reviewers and editors' concerns

Editors' concerns

Dear authors, thank you for your responses and revision. There are just a few more minor comments to handle prior to moving this manuscript forward:

1. The abstract reads much better than before. A few additional suggested changes:

a. Abstract Results, line 17 – please add ‘with an increase in odds of ALS’.

We have added this to the revised manuscript (P. 1, Lines 16-18):

For a standard deviation (SD) increase in 5-year average concentrations, EC ($SD=0.42\mu\text{g}/\text{m}^3$) had a high probability of being individually associated with an increase in odds of ALS [...]

b. Abstract Conclusions, lines 25-27 – suggested revision to ‘This study found a high probability of a positive association between ALS diagnosis and EC concentrations. Further work is needed...’ I suggest removing reference to the results being inconclusive, as this seems inconsistent with conclusion of the high probability finding for EC.

We have added this to the revised manuscript (P. 2, Lines 25-27):

This study found a high probability of a positive association between ALS diagnosis and EC concentration. Further work is needed to understand the role of traffic-related air pollution on ALS pathogenesis.

2. Thank you for including reference to the Nunez et al. article. This reference should be helpful for readers to place the air pollution-ALS literature into context. If I’m not mistaken, the outcome database as well as the source of the pollution data (despite the different specific pollutants examined) are largely overlapping between your work and that of Nunez et al. For further transparency, I suggest:

a. Adding a mention at the very end of the introduction or beginning of the methods, that this study builds on (or pairs with) the work of Nunez et al.

We have added a recognition that this study builds on the work of Nunez et al. to the revised manuscript (P. 4, Lines 77-78):

This study pairs with and complements the work of Nunez et al.⁴⁰

b. When citing references to epidemiology studies that use the predicted pollutant concentrations in Denmark (line 129), please include the Nunez et al. citation #40

We have added the citation of Nunez et al. here in the revised manuscript (P. 6, Lines 129-130):

These predicted pollutant concentrations have been extensively used in previous air pollution epidemiologic studies in Denmark.^{17,40,57–59}

3. The definitions and terminology for the joint and average traffic pollution effects are much clearer than before. A couple of suggestions to further clarify:

a. On line 182, suggest ‘... i.e., total percentage change in the odds of ALS diagnosis with a simultaneous increase in each of EC, NO_x, CO).’

We have added this to the revised manuscript (PP. 8-9, Lines 182-184):

[...] (b) a joint association of the three pollutants (i.e., total percentage change in odds of ALS diagnosis with a simultaneous increase in each of EC, NO_x, CO) [...]

b. On line 183-184, suggest ‘... i.e., average percentage change in the odds of ALS diagnosis with increases in each of EC, NO_x, CO),...’

We have added this to the revised manuscript (P. 9, Lines 184-185):

[...] and (c) an average traffic association (i.e., average percentage change in odds of ALS diagnosis with increases in each of EC, NO_x, CO), [...]

4. Currently, eFigure 2 is referenced at the very end of the results section (lines 302-304). From a flow perspective, this sentence seems better placed in the second paragraph of the results, when discussing pollutant concentrations. As such, I suggest switching the order of eFigure 2 and eFigure 1.

We have moved the sentence and switched the order of the eFigures to match this in the revised manuscript (P. 13, Lines 283-285):

Maps of average concentration of included pollutants (EC, NO_x, CO, PM_{2.5}, O₃) across Denmark for a representative year (2000; middle of study period 1989-2013) is also available in eFigure 1.

5. Paragraph 3 of the results section needs a few clarifications.

a. Please reference ‘odds of ALS diagnosis’ as the outcome when introducing observed associations on lines 284-286.

We have added this to the revised manuscript (P. 13, Lines 287-288):

For 5-year average pollutant concentrations, we observed the largest overall increase in odds of ALS diagnosis for the individual SD increase in EC [...]

b. Please also specify the units for the increases. E.g., they are specifically mentioned for EC, NO_x, and CO on lines 284-289, however missing for non-EC PM_{2.5} on line 290 and 1-year EC on line 291.

We have added units for the increases to the revised manuscript (P. 13, Lines 292-296):

Non-EC PM_{2.5} was not associated with ALS diagnosis (0.7%; 95% CrI: -9.2%, 12.4% per 2.37 µg/m³; 54.1% posterior probability of positive association). 1-year EC average exposure was associated with an increase in odds of ALS diagnosis (15.4%; 95% CrI: 1.6%, 25.6% per 0.42 µg/m³; 98.9% posterior probability of positive association).

c. It also occurs to me that while the SD increases for 5-year exposures are included in Table 2, the SD increases for 1-year and 10-year exposure measures are not specified anywhere. Please consider either adding summaries for these exposures to Table 2, or perhaps include the SD units for each pollutant and averaging time in the Figure 2 title notes.

We have now added 1-year and 5-year exposure measures to Table 2 in the revised manuscript, copied below for convenience:

Table 2. Summary of 1-, 5-, and 10-year average pollutant concentrations (all in $\mu\text{g}/\text{m}^3$).

	Pollutant	Overall, N = 23,270 ^a	Case, N = 3,937 ^a	Control, N = 19,333 ^a
1-year average	EC	0.81 (0.42)	0.83 (0.44)	0.81 (0.42)
	NO _x	26 (19)	26 (20)	26 (19)
	CO	224 (97)	226 (101)	224 (96)
	non-EC PM _{2.5}	11.17 (2.32)	11.20 (2.34)	11.17 (2.31)
	O ₃	52.6 (6.1)	52.4 (6.2)	52.6 (6.1)
	Pollutant	Overall, N = 23,232 ^a	Case, N = 3,934 ^a	Control, N = 19,298 ^a
5-year average	EC	0.85 (0.42)	0.86 (0.45)	0.85 (0.42)
	NO _x	27 (20)	28 (21)	27 (20)
	CO	238 (106)	239 (112)	237 (105)
	non-EC PM _{2.5}	11.76 (2.37)	11.78 (2.41)	11.76 (2.37)
	O ₃	51.9 (6.0)	51.9 (6.1)	52.0 (6.0)
	Pollutant	Overall, N = 23,179 ^a	Case, N = 3,929 ^a	Control, N = 19,250 ^a
10-year average	EC	0.89 (0.43)	0.89 (0.46)	0.88 (0.43)
	NO _x	29 (20)	29 (22)	29 (20)
	CO	253 (115)	255 (122)	253 (113)
	non-EC PM _{2.5}	12.53 (2.55)	12.55 (2.59)	12.52 (2.55)
	O ₃	51.3 (6.0)	51.3 (6.1)	51.4 (6.0)

^aMean (SD)

6. Line 295, referring to the EC single-pollutant model in eFigure 1, should this be model F (not model D)?

We thank the Editor for pointing this out. We have corrected this in the revised manuscript (P. 14, Lines 299-301):

The 95% credible interval for EC in the single-pollutant model (eFigure 1; model F) overlapped with the credible intervals of the EC term in the multi-pollutant models (eFigure 1; models A to C, G to P).

7. Lines 315-319, the sentences here seem somewhat duplicative. Consider tightening the wording.

We have tightened up the language by merging the two referenced sentences in the revised manuscript (PP. 14-15, Lines 318-321):

The inconsistent associations for NO_x and CO in the multi- and single-pollutant models and the consistency of the EC association suggest that EC concentrations may have been more relevant than NO_x and CO for ALS diagnosis, though further study is required.

8. Regarding the discussion on BMI and smoking, the previous reviewer and I both questioned the likelihood of these as potential confounders independent of SES. It seems plausible that BMI and smoking could co-vary with pollution levels, independently of SES; this wouldn't depend on these factors being included in the air pollution modeling system. In any case, if I understand your argument correctly, may I suggest the following revision simply to clarify your discussion of this (starting on line 367): '....not currently available through the Danish Civil Registration System. These variables, while potential risk factors for ALS, are not likely confounders in this analysis as they are not expected to be associated with pollutant concentrations in a manner independent of neighborhood SES. If information on these variables were available, it could be used to further adjust for SES. ... '

We have added this to the revised manuscript (P. 17, Lines 368-373):

Information on individual-level variables, such as body mass index (BMI) and smoking status is not currently available through the Danish Civil Registration System. These variables, while potential risk factors for ALS, are not likely confounders in this analysis as they are not expected to be associated with pollutant concentrations in a manner independent of neighborhood SES. If this information were available, it could be used to further adjust for SES.⁸⁶

* * * * *

Preparing a revision

1. For estimates of causal effects, we strongly discourage the use of categorized P-values and language referring to statistical significance, including whether a confidence interval covers the null. We prefer instead interval estimation, which conveys the precision of the estimate with respect to sampling variability. We are more open to testing with respect to modeling decisions, such as for tests of interaction and for tests for trend.

We have avoided p-values throughout.

2. We do not permit acronyms unless they are generally recognized by epidemiologists (e.g. HIV is okay, but LVA is not). When in doubt, we recommend that you spell out.

We have been careful to introduce acronyms where used.

3. Please do not include uninformative precision (excessive decimal places). For example, percents should be rounded to nn%, n.n%, or 0.0n% and risk ratios should be rounded to nn, n.n, or 0.nn unless clarity of the presentation and the sample size justify more significant digits.

We have done this.

4. Please be sure to include explicit information about approval of human subjects research by an independent review board. If no such review was required, include an explicit statement about why the requirement for review was waived.

We have done this in the manuscript (P. 5, Lines 113-114):

This study was approved by the Institutional Review Board Committee at Columbia University and the Danish Data Protection Agency.

5. Do not include public health policy recommendations in Brief Reports or Original Articles that present new research findings.

We have not included any public health policy recommendations.

6. Data appearing in the abstract must also be cited in the main text, not just in tables or figures.

We have done this.

7. Resubmissions must adhere to word limits. The word limits for main text (generally the introduction, methods, results, and discussion) are 1500 words for Brief Reports (plus 150 words for its abstract), 4000 words for Original Articles (plus 250 words for its abstract), 5000 words for reviews (plus 250 words for its abstract), 2000 words for Commentaries (no abstract), 600 words for Research Letters (no abstract), and 400 words for Letters to the Editor (no abstract).

Based on Editors' suggestions, we now have an Abstract of 251 words and an Original Article of 4,154 words in the revised manuscript.

8. We advise that total word counts for Original Articles should not exceed 7500 words and for Brief Reports should not exceed 3500 words. The total word count includes main text (introduction, methods, results, and discussion), bibliography, figure legends, tables, and figures (250 words per figure, including each figure in a panel). The title page, abstract, acknowledgments, and funding information do not count in the total word count.

Our total word count of 6,784 words in the revised manuscript.

9. Figure labels: Make font size as large as possible, so as to be legible when figures are reduced for publication (typically one column [8.5cm] in width).

We have made the Figure labels large and legible.

10. Footnotes to tables and figures should use superscript lowercase letters to link content to the footnote, not symbols or numerals.

The footnotes in Tables 1 and 2 use a superscript lowercase letter.

11. Do not use parenthetical phrases like “(data not shown), (results not shown), or (available from the authors upon request).” In these circumstances, the data or results should be provided in Supplementary Digital Content.

We have avoided any use of these phrases.

12. Additional details regarding submission requirements can be found in the Instructions for Authors, which are posted at <http://edmgr.ovid.com/epid/accounts/ifaauth.htm>.

We have reviewed these details.

Preparing for resubmission

13. Prepare a response document for the Editor that responds point-by-point to the reviewers' comments (presenting each comment followed by your response). Give the page number where revised text can be found and, where practical, paste revised text directly into the reply document.

We have done this.

14. Submit versions of the manuscript with and without your changes displayed.

We have submitted clean and tracked versions of the revised manuscript.

15. Supplementary Digital Content should be submitted as a single PDF file, and you should use our convention - e.g. eFigure 1, eAppendix 2 - to label and refer to online content.

We have done this.

16. Authors should submit copies of any closely related manuscripts (published, in press, or under review).

As discussed with the Editors, we now mention the Nunez et al. paper into our revised manuscript.

17. Please revisit information about page charges and color printing charges available in the Instructions for Authors, which are posted at <http://edmgr.ovid.com/epid/accounts/ifaauth.htm> .

We acknowledge the charges on the link provided.

18. We request that the complete revised manuscript (with all tables and figures) be completed by 28 Sep 2022. If you are not able to meet this deadline, please notify the editorial office.

We have submitted before 28th September 2022.

Resubmitting via Editorial Manager

19. Log-in to Editorial Manager as an author using the credentials above.

20. Click on the "Submissions Needing Revision" link.

21. To view the previous decision letter and reviewer comments, please click the blue decision term listed under the View Decision menu.

22. If you would like to download the previous manuscript to make revisions, click on "Download Files" under the Action menu.

23. To begin the resubmission: Click "Submit Revision" under the Action menu.

24. Proof each screen to ensure the information is still correct (the Title, Authors, etc.), then click Next at the bottom of each page.

25. On the Attach Files screen, select each previous submission item that you would like to carry forward to the resubmission.

26. Upload the revised versions of the main text (with and without tracked changes), and order them with the highlighted version first.

27. Upload the point-by-point reply to review.

28. When you are finished uploading, please click Next.

29. Click "Build PDF for My Approval."

30. Click "Go to Submissions Waiting for Author's Approval."

31. Wait for the PDF to build. When it has been built, you will see the link "View Submission" in the Action menu. Click "View Submission," and open the manuscript to proof your work.

32. If you find problems with the manuscript, click "Edit Submission" from the Action menu. Make the required changes, and begin again at the file uploads.

33. Once the submission is complete and acceptable, click "Approve Submission" from the Action menu.

34. If you have difficulty with these procedures, you may send questions to timothy.lash@epidemiology-journal.com.

Thank you for the resubmission instructions. We have followed them.

1 Abstract

2 **Background:** Amyotrophic lateral sclerosis (ALS) is a fatal neurodegenerative disease. Limited
3 evidence suggests ALS diagnosis may be associated with air pollution exposure and specifically
4 traffic-related pollutants.

5
6 **Methods:** In this population-based case-control study, we used 3,939937 ALS cases from the
7 Danish National Patient Register diagnosed during 1989–2013 and matched on age, sex, year of
8 birth and vital status to 19,298333 population-based controls free of ALS at index date. We used
9 validated predictions of elemental carbon (EC), nitrogen oxides (NO_x), carbon monoxide (CO),
10 and fine particles (PM_{2.5}) to assign 1-, 5-, and 10-year average exposures pre-ALS diagnosis at
11 study participants' present and historical residential addresses. We used a Bayesian hierarchical
12 conditional logistic model, adjusting for potential confounders, to estimate individual pollutant
13 associations, well as joint and average associations for the traffic-related pollutants (EC, NO_x,
14 CO).

15
16 **Results:** For a standard deviation (SD) increase in 5-year average concentrations, EC
17 (SD=0.42µg/m³) had a high probability of being individually associated with an increase in odds
18 of ALS (11.5%; 95% credible interval[CrI]:-1.0%,25.6%; 96.3% posterior probability of a positive
19 association), with negative associations for NO_x (SD=20µg/m³) (-4.6%;95%CrI-18.1%,8.9%;
20 27.8% posterior probability of a positive association), CO (SD=106µg/m³) (-3.2%;95%CrI-
21 14.4%,10.0%; 26.7% posterior probability of a positive association) and a null association for non-
22 EC PM_{2.5} (SD=2.37µg/m³) (0.7%;95%CrI-9.2%,12.4%). We found no association between ALS
23 and joint or average traffic pollution concentrations.

24

25 **Conclusions:** ~~This study found a~~ high probability of a positive association between ALS
26 diagnosis and EC concentration, ~~though results are inconclusive~~. Further work is needed to
27 understand the role of traffic-related air pollution on ALS pathogenesis.

28

29 **Abbreviations:**

30	ALS	Amyotrophic lateral sclerosis
31	BKMR	Bayesian kernel machine regression
32	BMI	Body mass index
33	CO	Carbon monoxide
34	CrI	Credible interval
35	DEHM-UBM-AirGIS	Spatio-temporal air pollution modelling system used in study
36	EC	Elemental carbon
37	ICD	International Classification of Diseases
38	IQR	Interquartile range
39	IR	Incidence ratio
40	Non-EC PM _{2.5}	Non-elemental carbon fine particles
41	NO _x	Nitrogen oxides
42	O ₃	Ozone
43	PM _{2.5}	Fine particles
44	SD	Standard deviation
45	SES	Socioeconomic status

46 **Introduction**

47 Amyotrophic lateral sclerosis (ALS) is a devastating and fatal neurodegenerative disease,¹
48 currently without a cure.² Approximately half of patients die within three years of symptom onset.³
49 Annually, there are nearly 30,000 cases of ALS in Europe and over 200,000 worldwide.⁴ Known
50 inherited genetic variants only account for 5–10% of ALS cases.^{5,6} Environmental factors,
51 therefore, are likely important in ALS pathogenesis.⁷ However, because the disease is relatively
52 rare, it is challenging to conduct large-scale prospective studies. There is a recognized need for
53 more evidence of the environmental contributors of ALS.^{5,8}

54
55 Although air pollution is commonly studied in association with respiratory- and cardiovascular-
56 related outcomes, e.g., references^{9–14}, epidemiologic and toxicological studies also support several
57 plausible biological mechanisms in association with the nervous system and neurodegeneration,
58 e.g., references^{15–34}. Ambient air pollution, especially urban air pollution, is a ubiquitous exposure
59 that has been associated with several other neurodegenerative disorders, e.g., references^{16–21,35,36},
60 and is consistently linked to systemic inflammation,^{22–24} oxidative stress,^{25–28} and
61 neuroinflammation,^{15,29} all of which, in turn, have been reported as key pathways to ALS
62 pathogenesis, e.g., references^{30–34}.

63
64 Despite the compelling plausibility, few studies to date have evaluated the association between air
65 pollution and ALS.^{35,37–39} A study in 2021 found that traffic-related air pollutants may be driving
66 observed associations.³⁸ Another study of ALS and PM_{2.5} in Denmark examining critical windows
67 of exposure found that more recent exposure to PM_{2.5} (i.e., the previous 1 to 5 years) may be the
68 most important driver of the potential association, though the constituents of PM_{2.5} were not

analyzed, neither together nor separately.⁴⁰ No study has hitherto attempted to understand the individual, joint, and average associations of the pollutants in a single model. Air pollutants have been consistently associated with adverse health, primarily in single pollutant analyses.^{13,17,41–43} However, they are highly correlated with one another.⁴¹ It is therefore a mixture modelling challenge to infer the association of multiple air pollutants and health outcomes.⁴⁴ Using three air pollutants commonly used in health studies as traffic-related emissions tracers—elemental carbon (EC), nitrogen oxides (NO_x), and carbon monoxide (CO)—we aimed to assess whether exposure to (a) each individual air pollutant is independently associated with ALS diagnosis, and estimate their (b) joint and (c) average traffic-related emissions associations. This study pairs with and complements the work of Nunez et al.⁴⁰

Methods

Study Population and Outcome Assessment

We used data from the Danish National Patient Register during 1989-2013, through which details on demographic characteristics and certain health outcomes of all Danish residents can be linked based on a unique personal identifier.⁴⁵ The Register was established in 1977 and is comprehensive, including nationwide clinical and administrative records for all inpatient data, with outpatient data available since 1995.⁴⁶

We identified ALS cases based on their International Classification of Diseases (ICD) discharge diagnoses, i.e., ICD-8 code 348.0 (ALS) until 1993 and ICD-10 code G12.2 (motor neuron disease) thereafter, using the date of the first relevant code as the diagnosis date. This was the index date. We only included patients who were at least 20 years old when diagnosed because (i) cases

92 younger than 20 years old were at a greater chance of misclassification, since ALS has been
93 predominantly diagnosed in older adults,⁴⁷ and (ii) the very few juvenile ALS cases have been
94 explained to a much larger degree by genetic mutations (~40%).⁴⁸ In our validation study, Register
95 data for ALS ascertainment were highly reliable; working with a specialist ALS neurologist to
96 review medical records and comparing to death certificates and hospital discharges, the Danish
97 National Patient Register was found to have an overall predictive value for ALS of 82%.⁴⁹

98

99 We obtained controls through the Danish Civil Registration System, established in 1968 and
100 updated daily, which includes administrative records (e.g., date and place of birth, sex, vital status,
101 and history of civil status and addresses since 1971) on all persons living in Denmark; records are
102 kept even when a person dies or emigrates.⁵⁰ We randomly matched five controls per case by age,
103 sex, year of birth, and vital status. Controls were alive and free of diagnosed ALS at the ALS
104 diagnosis date of the matched case (index date). The control-sampling scheme followed a risk-set
105 matching pattern, so cases could have served as controls before diagnosis of ALS.⁵¹

106

107 We obtained all addresses of cases and controls from January 1st 1979 onwards from the Danish
108 Civil Registration System,⁵⁰ including the dates of moving to and from each address, before the
109 index date. We then obtained the geographical coordinates at the door of each house of the
110 residential history of the participants, with previous evidence of the high accuracy of this method
111 of geocoding of addresses in Denmark.¹⁷

112

113 This study was approved by the Institutional Review Board Committee at Columbia University
114 and the Danish Data Protection Agency.

115

116 *Exposure data*

117 We obtained predictions on monthly concentrations of elemental carbon (EC), nitrogen oxides
118 (NO_x), carbon monoxide (CO), and fine particles (PM_{2.5}) (as well as ozone, O₃, for a sensitivity
119 analysis, usually negatively correlated with other pollutants due to its chemistry⁵²), at residential
120 addresses of study participants from the validated spatio-temporal air pollution modelling system
121 DEHM-UBM-AirGIS that provides full space and time coverage over the study period, described
122 in detail elsewhere.^{53–56} In brief, DEHM-UBM-AirGIS is a human exposure modelling system for
123 traffic pollution, developed for application in Danish air pollution epidemiological studies. The
124 modelling system integrates air pollution dispersion models, digital maps, national and local
125 administrative databases, concentrations of air pollutants at regional, urban background and street
126 level, meteorological data, and a Geographic Information System (GIS). The modelling system is
127 therefore able to generate street configuration and traffic data based on digital maps and national
128 databases, which enables estimation of air quality levels at a large number of addresses in an
129 automatic and effective way. These predicted pollutant concentrations have been extensively used
130 in previous air pollution epidemiologic studies in Denmark.^{17,57–59,17,40,57–59} The models have good
131 predictive accuracy, with average monthly correlations between measured and modelled results of
132 0.79 for EC, 0.85 for NO_x, 0.91 for CO, 0.92 for O₃, and 0.83 for annual concentrations of
133 PM_{2.5}.^{53,56} Because traffic is a major source of PM_{2.5} and EC one of the main PM_{2.5} components in
134 urban environments,⁶⁰ we removed the EC concentration from the total PM_{2.5} mass concentration
135 (non-EC PM_{2.5}) by subtraction to avoid overadjustment when including both in the models
136 simultaneously; this was valid since the DEHM-UBM-AirGIS modelling system constructed
137 PM_{2.5} concentrations by adding from specific species of pollutants, one of which was EC.^{53–56}

138
139 Based on the residential history of each case or control, we calculated 1-, 5-, and 10-year average
140 exposure to each pollutant ending at one year before the index date, as diagnosis has been shown
141 previously to occur at a median of 12 months after symptoms onset.⁶¹ Specifically, each case or
142 control average value (1-, 5- or 10-year) was calculated as the mean of all concentrations recorded
143 across time at the recorded addresses within each time window. A small number of Danish
144 residents lack a complete address history (1.7%; lack of house number). To ensure we were
145 including participants with adequately complete exposure records, we set the following minimum
146 criteria for number of complete exposure record months to include cases and controls: (i) 1-year
147 averages: 9 of 12 months, at least one measurement in each season; (ii) 5-year averages (main
148 exposure): 30 of 60 months; and (iii) 10-year averages: 60 of 120 months.

149

150 *Covariate data*

151 We included a set of covariates based on as close as possible to index date to account for potential
152 confounding bias, including household socioeconomic status (SES) based on last-reported job title
153 at index date; civil status at index date, last reported place of residence at index date, and place of
154 birth. We used a five-category individual-level SES definition developed by the Danish Institute
155 of Social Sciences, based on job titles from income tax forms, which has been associated with ALS
156 diagnosis in Denmark,⁶² as well as how quickly one is identified as having ALS in the Danish Civil
157 Registration System.⁶³ Group 1 (highest status) includes corporate managers and academics; group
158 2: proprietors, managers of small businesses and teachers; group 3: technicians and nurses; group
159 4: skilled workers; and group 5: unspecialized workers, such as entry-level positions within food
160 and retail environments. We also included a group for participants whose job title was unknown

(group 9). For each married participant, we used the higher of the couple's individual SES categories, when available. We also used information on civil status (never married, married, divorced, widowed) due to the influence that a spouse may have on visiting a family physician,⁶⁴ last reported place of residence from postcode (Greater Copenhagen, big cities of Denmark, rest of Denmark, Greenland) to account for various local environmental and behavioral stressors,⁷ and place of birth (Greater Copenhagen, big cities of Denmark, rest of Denmark, Greenland, foreign, unknown) to adjust for other potential family-specific, location-specific, and early-life confounders, which may have an impact on the probability of developing ALS.⁶⁵ Ultimately, we were limited by what was available in the Danish Civil Registration System.⁶³ As part of a sensitivity analysis, we also included parish-level SES, measured by percentage of residents with greater than high-school education, in the model. In Denmark, parishes are administrative units with an average population of ~2,500 residents.

Statistical analysis

We analyzed the association between ALS diagnosis (binary) and exposure to traffic-related pollutants by applying a Bayesian formulation of the conditional logistic model, with Bayesian hierarchy on the traffic-related pollutants (EC, NO_x, CO).^{66,67} The conditional approach examines contrasts within matched strata, i.e., groupings of case and matched controls, implicitly adjusting for matching factors (age, sex, year of birth, vital status) within each matched stratum.⁶⁶ Matching by finer scale than year of birth was not possible. Bayesian inference allows for full distributional estimation of parameters of interest.⁶⁷ We employed a Bayesian hierarchical formulation because it enables estimates of (a) independent pollutant-outcome associations, (b) a joint association of the three pollutants (i.e., total percentage change in odds of ALS diagnosis with [a simultaneous](#)

increase in each of EC, NO_x, CO), and (c) an average traffic association (i.e., average percentage change in odds of ALS diagnosis ~~from~~ with increases in each of EC, NO_x, CO), while accounting for the variance-covariance structure between the highly-correlated exposures and their coefficients.⁶⁷ We included a linear term for each included pollutant and adjusted for individual- and parish-level SES, civil status, last reported place of residence, and place of birth.

Specifically, via a logit function, we modelled the log-odds of ALS diagnosis, as follows:

$$\begin{aligned} \text{logit}[\text{Pr}(Y_{ci} = 1)] = & \alpha_c + \\ & \beta_{EC}EC_{ci} + \beta_{NO_x}NO_{xci} + \beta_{CO}CO_{ci} + \\ & \beta_{PM_{2.5}}(\text{non-EC } PM_{2.5ci}) + \\ & \beta_{SES}SES_{ci} + \beta_{Civil\ Status}Civil\ Status_{ci} + \beta_{Residence}Residence_{ci} + \beta_{Birth}Birth_{ci}, \end{aligned}$$

where Y_{ci} denotes whether subject i in matched stratum c was diagnosed with ALS, i.e., c represents a case and its matched controls; α_c the matched stratum-specific intercepts (not estimated in conditional logistic models); β_{EC} , β_{NO_x} , β_{CO} , $\beta_{PM_{2.5}}$ the individual pollutant coefficients (log-odds) per standard deviation (SD) increase in concentration of EC, NO_x, CO, non-EC PM_{2.5} respectively, scaled by their respective SDs and centered at their means, with each β an individual pollutant association adjusted by other terms in the model and the rest as coefficients for subject-specific covariates. Interquartile Range (IQR) could equivalently be used to scale pollutant concentrations. If other sources of air pollution are associated with ALS, then including non-EC PM_{2.5} adjusts for PM_{2.5} from other sources,⁶⁸ as well as indicating whether pollution from other sources not explicitly quantified might also have associations with ALS. Therefore, $\beta_{PM_{2.5}}$ is interpreted as the association with air pollutants not specifically included in our analysis. In urban European environments, traffic-related pollutants typically represent on-

average 14% of $PM_{2.5}$ concentrations.⁶⁹ In a sensitivity analysis, we included O_3 in the model, as O_3 concentrations have been associated with many adverse health outcomes,⁷⁰ and were negatively correlated with traffic-related pollutants, and added $ns(SES_{\text{parish}_{c_i}})$, as a natural spline with three degrees of freedom.

In our model, β_{EC} , β_{NO_x} , and β_{CO} represent the independent individual pollutant associations with ALS diagnosis. In the same model, we estimated the joint association between these three pollutants and ALS diagnosis as:

$$Traffic_{Joint} = \sum_{p=EC,NO_x,CO} \beta_p p.$$

This sum quantifies the association (log-odds) with ALS of a one-SD increase in the three pollutants simultaneously.

Finally, we assumed that the traffic-related individual pollutant associations arise from a distribution of the average traffic association with ALS diagnosis. We placed a hierarchy on the traffic-specific individual pollutant terms in the model to account for the fact that the traffic-related pollutants, EC, NO_x , CO, originate from common sources and primarily traffic in urban environments:

$$\beta_{Traffic} = [\beta_{EC}, \beta_{NO_x}, \beta_{CO}],$$

$$\beta_{Traffic} \sim MVN(\mu, \Sigma),$$

$$\mu \sim N(\lambda, \sigma_\lambda),$$

$$\Sigma = \tau \Omega \tau,$$

228 where λ denotes the average one-SD association of traffic-related pollution with variance σ_λ . Σ ,
229 the estimated variance-covariance matrix among individual pollutant estimates, was expressed as
230 a decomposition into a positive-definite correlation matrix Ω and scale matrix τ .⁷¹

231
232 We used weakly-informative priors so that data drove parameter estimation. Hyper-priors for
233 coefficients on non-EC $\text{PM}_{2.5}$ and covariates were $N(0,10)$; for σ_λ and τ we used Half-
234 Cauchy(0,10), as recommended by Gelman, Polson and Scott as a weakly-informative prior;^{72,73}
235 Ω was defined by the weakly-informative prior $\text{LKJCorr}(1)$.⁷⁴ The exception to this was the prior
236 for λ , the average association of traffic-related pollutants, for which estimates became
237 unrealistically high (approaching infinity and not converging with further iterations) with a more
238 weakly-informative prior. We therefore used a prior of $N(0,0.1)$, which did not affect estimates of
239 other parameters. We conducted sensitivity analyses to understand the influence of priors and the
240 robustness of the results.

241
242 We present all results as percentage change in odds of ALS diagnosis per SD increase in pollutant
243 concentration (calculated via e.g., $e^{\beta_{\text{EC}}} - 1$, etc. obtained in the modelling process). Due to the
244 risk-set matching pattern of our case-control study, odds ratios are also equivalently incidence
245 ratios (IRs).⁶⁶ We ran each model with four chains with a sample size of 1,000 each, after a warm-
246 up of 1,000 samples, for 4,000 total samples. We assessed whether the models converged by
247 checking that the Gelman-Rubin potential scale reduction statistic⁷⁵ was below 1.1 for all estimated
248 model parameters. The reported 95% credible intervals (CrI) are the 2.5th to 97.5th percentiles of
249 each parameter's posterior marginal distribution. To calculate the probability that an association
250 estimate was greater than null, we used the 4,000 samples of the posterior distribution and took

251 the proportion of samples which were above the null. A 50% probability means that it is as likely
252 as not that the marginal estimate is null, a probability closer to 100% indicates that the association
253 is more likely to be truly positive, with closer to 0% indicating more likely to be truly negative.

254
255 We conducted statistical analyses using the R Statistical Software, version 4.1.1⁷⁶ and R-STAN,
256 version 2.21.2.⁶⁷ All code for analysis, results from analysis, and visualization presented in this
257 manuscript is publicly available via GitHub at
258 https://github.com/rmp15/traffic_air_pollution_als_denmark_epidemiology.

259
260 We assessed the sensitivity of our results to hyper-prior assignment; running more iterations and
261 warm-up per chain; inclusion of O₃; single traffic-related pollutant models adjusting for non-EC
262 PM_{2.5}; as well as adjusting by parish-level SES. From the parish-level SES sensitivity analysis we
263 excluded those who lived in areas without parish-level SES data, namely: (i) 819 participants for
264 1-year average exposure; (ii) 826 participants for 5-year average exposure; and (iii) 838
265 participants for 10-year average exposure.

266
267 **Results**
268 After filtering the original 4,011 cases and 20,055 controls based on completeness of exposure
269 records, we used information on 3,934 (98.1% of total) cases and 19,298 (96.2% of total) controls
270 for 5-year average exposure. We also used 3,937 cases, 19,333 controls for 1-year average exposure
271 and 3,939~~29~~ cases, 19,250 controls for 10-year average exposure. Descriptive statistics of
272 included cases and controls for 5-year average exposure can be found in Table 1. Descriptive
273 statistics of controls for 5-year exposure by socioeconomic status, civil status, residence, and place

of birth are found in eTables 1-4. For the main results, we present 5-year average exposure associations as a balance between representation of most recent exposure as well as long-term concentration.

The 5-year average traffic-related pollutant concentrations were 0.85 $\mu\text{g}/\text{m}^3$ for EC (SD=0.42 $\mu\text{g}/\text{m}^3$), 27 $\mu\text{g}/\text{m}^3$ for NO_x (SD=20 $\mu\text{g}/\text{m}^3$), and 238 $\mu\text{g}/\text{m}^3$ for CO (SD=106 $\mu\text{g}/\text{m}^3$) (Table 2). Figure 1 shows Spearman correlations between pollutants for 1-, 5-, and 10-year average exposures. Traffic-related pollutants (EC, NO_x , CO) were highly correlated in cases, controls and overall, ranging from correlations of 0.91 to 0.96. Otherwise, non-EC $\text{PM}_{2.5}$ was most highly correlated with CO, ranging from 0.67 to 0.7. O_3 was negatively correlated with other pollutants, ranging from -0.54 to -0.89. Maps of average concentration of included pollutants (EC, NO_x , CO, $\text{PM}_{2.5}$, O_3) across Denmark for a representative year (2000; middle of study period 1989-2013) are also available in eFigure 1.

Formatted: Font: Italic, Font color: Auto

For 5-year average pollutant concentrations, we observed the largest overall ~~association~~increase in odds of ALS diagnosis for the individual SD increase in EC (11.5%; 95% CrI: -1.0%, 25.6% per 0.42 $\mu\text{g}/\text{m}^3$; 96.3% posterior probability of positive association) (Figure 2). SD increases were associated with a decrease in odds of ALS diagnosis in NO_x (-4.6%; 95% CrI: -18.1%, 8.9% per 20 $\mu\text{g}/\text{m}^3$; 27.8% posterior probability of positive association) and CO (-3.2%; 95% CrI: -14.4%, 10.0% per 106 $\mu\text{g}/\text{m}^3$; 26.7% posterior probability of positive association). Non-EC $\text{PM}_{2.5}$ was not associated with ALS diagnosis (0.7%; 95% CrI: -9.2%, 12.4%); ~~% per 2.37 $\mu\text{g}/\text{m}^3$; 54.1% posterior probability of positive association).~~ 1-year EC average exposure was associated with an increase in odds of ALS diagnosis (15.4%; 95% CrI: 1.6%, 25.6%); ~~% per 0.42 $\mu\text{g}/\text{m}^3$; 98.9% posterior~~

probability of positive association). Single-pollutant models for each traffic-related pollutant adjusting for non-EC PM_{2.5} (eFigure 1; single traffic-related pollutant models D, E and F) resulted in positive associations for each of EC, NO_x, CO, with positive associations for non-EC PM_{2.5} in all but the model with EC. The 95% credible interval for EC in the single-pollutant model (eFigure 1; model ~~DE~~) overlapped with the credible intervals of the EC term in the multi-pollutant models (eFigure 1; models ~~B,A~~ to C, G to P). The joint association of traffic-related pollutants (EC, NO_x, CO) was 2.3% (95% CrI: -3.3%, 7.7%), with an 77.8% posterior probability of a positive association. The average traffic association was null (-0.1%; 95% CrI: -17.4%, 20.8%); ~~%: 45.5%~~ posterior probability of positive association). Compared to the 1- and 5-year results, the 10-year average exposure results were attenuated, as associations tended further to the null. Results from variations of the main model in the sensitivity analyses were robust to prior choices, inclusion of O₃, and inclusion of parish-level SES (eFigure 1). ~~A map of average concentration of included pollutants (EC, NO_x, CO, PM_{2.5}, O₃) across Denmark for a representative year (2000; middle of study period 1989-2013) is also available in eFigure 2.2).~~

Discussion

In the largest case-control study of ALS and traffic-related air pollution to date, we found that EC had the largest-in-magnitude independent association with ALS diagnosis, while associations with NO_x and CO were negative with credible intervals overlapping the null, and smaller in magnitude. A joint increase in concentrations of traffic-related pollutants had a high probability of being associated with an increase in odds of ALS diagnosis. Sensitivity analyses demonstrated that for single pollutant models, the association for EC was smaller than for our main multi-pollutant model, which took into account the variance-covariance structure of traffic-related pollutants.

Overall conclusions for the association between EC and ALS diagnosis were similar from the single- or multi-pollutant models. The inconsistent associations for NO_x and CO in the multi- and single-pollutant models and the consistency of the EC association suggest that EC concentrations may have been more relevant than NO_x and CO for ALS diagnosis. ~~Nevertheless, the consistency of the sign of the central estimate of EC in all models suggests that EC may be a driver of the ALS and traffic-related pollutant association,~~ though further study is required. Our results indicate that traffic-related pollutants, hazardous in many ways,^{9–21,41–43} may also be associated with ALS diagnosis. Our finding—that increases in EC, are potentially positively associated with ALS diagnosis—is plausible. A case-control study in the Netherlands from 2021 reported that ultrafine particles, another traffic emissions-related surrogate, were associated with ALS diagnosis,³⁸ while another based in Catalonia, Spain found ALS cases clustered around key road infrastructure.⁷⁷ Although we did not find an association with non-EC PM_{2.5} in our study, our results are not directly comparable to those of the other studies, as our PM_{2.5} effect estimates capture the PM_{2.5} components not accounted for by other pollutants in the analysis. A study examining critical windows of exposure of PM_{2.5} and ALS diagnosis in Denmark found that concentrations 1 to 5 years before exposure may be driving the association with ALS onset,⁴⁰ consistent with our findings that the most recent 1-year average EC concentration exhibited the largest association.

Our results indicate that EC exposure—a large part of which comes from diesel combustion and small combustion sources (such as wood stoves) in European urban centers, where prevalence of diesel cars is high⁷⁸—has a high probability of a positive association with ALS diagnosis. In our previous study of ALS and occupational exposures in Denmark we found that those working in agriculture and construction, associated with exposure to diesel engine exhausts, were at higher

relative risk than those in other employments.⁶² Truck drivers, for whom diesel exposure is common, are also at increased risk of sporadic ALS.⁷⁹ EC exposure has been associated with inflammation,⁸⁰ mitochondrial dysfunction⁸¹ and DNA damage,^{81,82} all of which are plausible pathways of neurodegeneration. These factors have also previously been identified as particular pathways to pathogenesis of ALS.³⁰⁻³⁴

We did not find a high probability of a positive association with NO_x in our analyses, in contrast with a previous study, though that study did not include EC.³⁸ NO_x is also highly correlated with EC (0.95⁹⁴ to 0.96 in our study), which is expected given that they are both combustion products commonly associated with emissions in urban environments. EC exposure was more strongly associated with 1-year than for 5-/10-year average concentrations, which may indicate that the previous year of exposure may be the most relevant exposure window relevant to traffic-related exposures and ALS; this is biologically plausible, as this critical exposure window would be at the pre-symptomatic stage of underlying ALS progression, where traffic-related pollution exposure may add to the ongoing cellular or molecular process of the disease, to the point where the body can no longer compensate and subsequently enters the clinical phase.⁸³⁻⁸⁵ We do not expect that these results are attributed to reverse causation, as we have lagged these 1-year exposures by one year already prior to diagnosis, and there was likely little substantial residential movement in the year before ALS diagnosis.⁸⁶⁸⁶ We do not expect that calendar time was a potential source of confounding, as the controls were matched on age and year of birth. The null joint association, combined with the largest associations from traffic-related pollutant in all models found with EC, further indicates that EC may be driving the association of air pollution with ALS, though further analysis will be necessary to confirm this.

366

367 Our study used one the largest number of ALS patients ever included in an environmental health
368 study. Another strength of our study is that we leveraged highly correlated traffic pollutants and
369 Bayesian hierarchical modeling and were able to estimate independent, joint, and average traffic-
370 related pollutant associations. Although we have adjusted implicitly (by matching; age, sex, year
371 of birth, vital status) and explicitly for many common covariates (SES, civil status, residence, place
372 of birth), we cannot rule out residual confounding. Information on individual-level variables, such
373 as body mass index (BMI) and smoking status is not currently available through the Danish Civil
374 Registration System. These variables, ~~however while potential risk factors for ALS,~~ are not likely
375 confounders in this analysis as they are not expected to ~~cause the predicted~~ be associated with
376 pollutant concentrations, ~~given exposure assessment in a manner independent of neighborhood~~
377 SES. If this information were available, it could be used to further adjust for SES.⁸⁶⁸⁶ To the extent
378 that the variables we included in our models to adjust for household- and neighborhood-level SES
379 are adequate, we would expect any residual SES-related confounding to be minimal. Exposure
380 measurement error is inevitable, as any modelled exposure will be inaccurate to some degree.
381 However, any error is not likely correlated with ALS diagnosis, and therefore any bias would be
382 towards null.⁸⁷ While a previous study found that ALS ascertainment from the Danish National
383 Patient Register was highly reliable,⁴⁹ outcome misclassification cannot be ruled out, nor can the
384 possibility that date of diagnosis and symptom onset were irregularly aligned. While our analysis
385 adjusted for marital status and household SES, many couples in Denmark cohabit. This would
386 not be captured by our analysis, and ALS diagnosis in relation to cohabitation status should be
387 further investigated.⁸⁸

388

Formatted: Font color: Auto

Formatted: Font color: Auto

Formatted: Font color: Auto

Formatted: Font color: Auto

Formatted: Font color: Auto

389 Future research might use larger cohort data to understand the importance of each respective
390 pollutant in a single model. Other mixture model methods, such as Bayesian Kernel Machine
391 Regression (BKMR)⁸⁹ might be useful in further exploring the robustness of joint associations in
392 a different framework, though BKMR was not appropriate for our particular research question,
393 since BKMR is currently not available for case-control study applications. The timing of exposure
394 will continue to be an important study route. ALS is projected to increase in prevalence over the
395 next few decades all over the world.⁴ Understanding ALS pathogenesis and identifying modifiable
396 risk factors is critical for preventive action.

References

1. Rowland LP, Shneider NA. Amyotrophic lateral sclerosis. *New England Journal of Medicine*. 2001;344(22):1688-1700.
2. Chio A, Logroscino G, Hardiman O, et al. Prognostic factors in ALS: A critical review. *Amyotrophic Lateral Sclerosis*. 2009;10(5-6):310-323.
3. Mitchell JD, Borasio GD. Amyotrophic lateral sclerosis. *The Lancet*. 2007;369(9578):2031-2041.
4. Arthur KC, Calvo A, Price TR, Geiger JT, Chio A, Traynor BJ. Projected increase in amyotrophic lateral sclerosis from 2015 to 2040. *Nature Communications*. 2016;7(1):1-6.
5. Al-Chalabi A, Hardiman O. The epidemiology of ALS: A conspiracy of genes, environment and time. *Nature Reviews Neurology*. 2013;9(11):617-628.
6. Hardiman O, Al-Chalabi A, Chio A, et al. Amyotrophic lateral sclerosis. *Nature reviews Disease primers*. 2017;3(1):1-19.
7. Oskarsson B, Horton DK, Mitsumoto H. Potential environmental factors in amyotrophic lateral sclerosis. *Neurologic Clinics*. 2015;33(4):877-888.
8. Longinetti E, Fang F. Epidemiology of amyotrophic lateral sclerosis: An update of recent literature. *Current Opinion In Neurology*. 2019;32(5):771.
9. Dominici F, Peng RD, Bell ML, et al. Fine particulate air pollution and hospital admission for cardiovascular and respiratory diseases. *JAMA*. 2006;295(10):1127-1134.
10. Bennett JE, Tamura-Wicks H, Parks RM, et al. Particulate matter air pollution and national and county life expectancy loss in the USA: A spatiotemporal analysis. *PLOS Medicine*. 2019;16(7):e1002856. doi:10.1371/journal.pmed.1002856
11. Schwartz J. Particulate air pollution and chronic respiratory disease. *Environmental Research*. 1993;62(1):7-13.
12. Schwartz J. The distributed lag between air pollution and daily deaths. *Epidemiology*. 2000;11(3):320-326.
13. Brook RD, Rajagopalan S, Pope III CA, et al. Particulate matter air pollution and cardiovascular disease: An update to the scientific statement from the American Heart Association. *Circulation*. 2010;121(21):2331-2378.
14. Dockery DW, Pope CA, Xu X, et al. An association between air pollution and mortality in six U.S. cities. *New England Journal of Medicine*. 1993;329(24):1753-1759. doi:10.1056/NEJM199312093292401

- 429 15. Block ML, Elder A, Auten RL, et al. The outdoor air pollution and brain health workshop.
430 *Neurotoxicology*. 2012;33(5):972-984.
- 431 16. Zanobetti A, Dominici F, Wang Y, Schwartz JD. A national case-crossover analysis of the
432 short-term effect of PM 2.5 on hospitalizations and mortality in subjects with diabetes and
433 neurological disorders. *Environmental Health*. 2014;13(1):1-11.
- 434 17. Ritz B, Lee PC, Hansen J, et al. Traffic-related air pollution and Parkinson's disease in
435 Denmark: A case-control study. *Environmental Health Perspectives*. 2016;124(3):351-356.
- 436 18. Kioumourtoglou MA, Schwartz JD, Weisskopf MG, et al. Long-term PM2.5 exposure and
437 neurological hospital admissions in the northeastern United States. *Environmental health*
438 *perspectives*. 2016;124(1):23-29.
- 439 19. Levesque S, Surace MJ, McDonald J, Block ML. Air pollution & the brain: Subchronic
440 diesel exhaust exposure causes neuroinflammation and elevates early markers of
441 neurodegenerative disease. *Journal of Neuroinflammation*. 2011;8(1):1-10.
- 442 20. Heusinkveld HJ, Wahle T, Campbell A, et al. Neurodegenerative and neurological disorders
443 by small inhaled particles. *Neurotoxicology*. 2016;56:94-106.
- 444 21. Power MC, Weisskopf MG, Alexeeff SE, Coull BA, Spiro III A, Schwartz J. Traffic-related
445 air pollution and cognitive function in a cohort of older men. *Environmental Health*
446 *Perspectives*. 2011;119(5):682-687.
- 447 22. Dubowsky SD, Suh H, Schwartz J, Coull BA, Gold DR. Diabetes, obesity, and hypertension
448 may enhance associations between air pollution and markers of systemic inflammation.
449 *Environmental Health Perspectives*. 2006;114(7):992-998.
- 450 23. Ruckerl R, Ibalá-Mulli A, Koenig W, et al. Air pollution and markers of inflammation and
451 coagulation in patients with coronary heart disease. *American Journal of Respiratory and*
452 *Critical Care Medicine*. 2006;173(4):432-441.
- 453 24. Hoffmann B, Moebus S, Dragano N, et al. Chronic residential exposure to particulate matter
454 air pollution and systemic inflammatory markers. *Environmental Health Perspectives*.
455 2009;117(8):1302-1308.
- 456 25. Kelly FJ. Oxidative stress: Its role in air pollution and adverse health effects. *Occupational*
457 *and Environmental Medicine*. 2003;60(8):612-616.
- 458 26. Chuang KJ, Chan CC, Su TC, Lee CT, Tang CS. The effect of urban air pollution on
459 inflammation, oxidative stress, coagulation, and autonomic dysfunction in young adults.
460 *American journal of respiratory and critical care medicine*. 2007;176(4):370-376.
- 461 27. Li N, Sioutas C, Cho A, et al. Ultrafine particulate pollutants induce oxidative stress and
462 mitochondrial damage. *Environmental Health Perspectives*. 2003;111(4):455-460.

- 463 28. Sørensen M, Daneshvar B, Hansen M, et al. Personal PM2.5 exposure and markers of
464 oxidative stress in blood. *Environmental health perspectives*. 2003;111(2):161-166.
- 465 29. Block ML, Calderón-Garcidueñas L. Air pollution: Mechanisms of neuroinflammation and
466 CNS disease. *Trends in neurosciences*. 2009;32(9):506-516.
- 467 30. Perry VH, Cunningham C, Holmes C. Systemic infections and inflammation affect chronic
468 neurodegeneration. *Nature Reviews Immunology*. 2007;7(2):161-167.
- 469 31. Bergeron C. Oxidative stress: Its role in the pathogenesis of amyotrophic lateral sclerosis.
470 *Journal of the neurological sciences*. 1995;129:81-84.
- 471 32. Mhatre M, Floyd RA, Hensley K. Oxidative stress and neuroinflammation in Alzheimer's
472 disease and amyotrophic lateral sclerosis: Common links and potential therapeutic targets.
473 *Journal of Alzheimer's disease*. 2004;6(2):147-157.
- 474 33. D'Amico E, Factor-Litvak P, Santella RM, Mitumoto H. Clinical perspective on oxidative
475 stress in sporadic amyotrophic lateral sclerosis. *Free radical biology and medicine*.
476 2013;65:509-527.
- 477 34. Perry VH, Nicoll JA, Holmes C. Microglia in neurodegenerative disease. *Nature Reviews*
478 *Neurology*. 2010;6(4):193-201.
- 479 35. Nunez Y, Boehme AK, Weisskopf MG, et al. Fine particle exposure and clinical aggravation
480 in neurodegenerative diseases in New York State. *Environmental Health Perspectives*.
481 2021;129(2):027003.
- 482 36. Nunez Y, Boehme AK, Li M, et al. Parkinson's disease aggravation in association with fine
483 particle components in New York State. *Environmental Research*. 2021;201:111554.
- 484 37. Malek AM, Barchowsky A, Bowser R, et al. Exposure to hazardous air pollutants and the
485 risk of amyotrophic lateral sclerosis. *Environmental Pollution*. 2015;197:181-186.
- 486 38. Yu Z, Peters S, van BL, et al. Long-Term Exposure to Ultrafine Particles and Particulate
487 Matter Constituents and the Risk of Amyotrophic Lateral Sclerosis. *Environmental Health*
488 *Perspectives*. 2021;129(9):097702. doi:10.1289/EHP9131
- 489 39. Seelen M, Toro CRA, Veldink JH, et al. Long-term air pollution exposure and amyotrophic
490 lateral sclerosis in Netherlands: A population-based case-control study. *Environmental*
491 *Health Perspectives*. 2017;125(9):097023. doi:10.1289/EHP1115
- 492 40. Nunez Y, Balalian A, Parks RM, et al. Exploring relevant time windows in the association
493 between PM2.5 exposure and Amyotrophic Lateral Sclerosis: A case-control study in
494 Denmark. *American Journal of Epidemiology*. Published online Under Revision.
- 495 41. Strak M, Weinmayr G, Rodopoulou S, et al. Long term exposure to low level air pollution
496 and mortality in eight European cohorts within the ELAPSE project: Pooled analysis. *BMJ*.
497 2021;374:n1904. doi:10.1136/bmj.n1904

498 42. Hamra GB, Laden F, Cohen AJ, Raaschou-Nielsen O, Brauer M, Loomis D. Lung cancer
499 and exposure to nitrogen dioxide and traffic: A systematic review and meta-analysis.
500 *Environmental Health Perspectives*. 2015;123(11):1107-1112.

501 43. Chen H, Kwong JC, Copes R, et al. Living near major roads and the incidence of dementia,
502 Parkinson's disease, and multiple sclerosis: A population-based cohort study. *The Lancet*.
503 2017;389(10070):718-726.

504 44. Gibson EA, Nunez Y, Abuawad A, et al. An overview of methods to address distinct
505 research questions on environmental mixtures: An application to persistent organic
506 pollutants and leukocyte telomere length. *Environmental Health*. 2019;18(1):1-16.

507 45. Frank L. When an entire country is a cohort. *Science*. 2000;287(5462):2398-2399.

508 46. Schmidt M, Schmidt SAJ, Sandegaard JL, Ehrenstein V, Pedersen L, Sørensen HT. The
509 Danish National Patient Registry: A review of content, data quality, and research potential.
510 *Clinical epidemiology*. 2015;7:449.

511 47. Trabjerg BB, Garton FC, van Rheenen W, et al. ALS in Danish registries: heritability and
512 links to psychiatric and cardiovascular disorders. *Neurology Genetics*. 2020;6(2).

513 48. Mathis S, Goizet C, Soulages A, Vallat JM, Le Masson G. Genetics of amyotrophic lateral
514 sclerosis: A review. *Journal of the Neurological Sciences*. 2019;399:217-226.

515 49. Kioumourtoglou MA, Seals RM, Himmelslev L, Gredal O, Hansen J, Weisskopf MG.
516 Comparison of diagnoses of amyotrophic lateral sclerosis by use of death certificates and
517 hospital discharge data in the Danish population. *Amyotrophic Lateral Sclerosis and*
518 *Frontotemporal Degeneration*. 2015;16(3-4):224-229.

519 50. Pedersen CB. The Danish civil registration system. *Scandinavian journal of public health*.
520 2011;39(7_suppl):22-25.

521 51. Langholz B, Goldstein L. Risk set sampling in epidemiologic cohort studies. *Statistical*
522 *Science*. Published online 1996:35-53.

523 52. Sillman S. The relation between ozone, NOx and hydrocarbons in urban and polluted rural
524 environments. *Atmospheric Environment*. 1999;33(12):1821-1845.

525 53. Khan J, Kakosimos K, Raaschou-Nielsen O, et al. Development and performance evaluation
526 of new AirGIS—a GIS based air pollution and human exposure modelling system.
527 *Atmospheric environment*. 2019;198:102-121.

528 54. Brandt J, Christensen JH, Frohn LM, Palmgren F, Berkowicz R, Zlatev Z. Operational air
529 pollution forecasts from European to local scale. *Atmospheric Environment*. 2001;35:S91-
530 S98.

531 55. Brandt J, Christensen J, Frohn L, Berkowicz R. Air pollution forecasting from regional to
532 urban street scale—implementation and validation for two cities in Denmark. *Physics and*
533 *Chemistry of the Earth, Parts A/B/C*. 2003;28(8):335-344.

534 56. Frohn LM, Ketzel M, Christensen JH, et al. Modelling ultrafine particle number
535 concentrations at address resolution in Denmark from 1979-2018—Part 1: Regional and
536 urban scale modelling and evaluation. *Atmospheric Environment*. 2021;264:118631.

537 57. Raaschou-Nielsen O, Andersen ZJ, Hvidberg M, et al. Lung cancer incidence and long-term
538 exposure to air pollution from traffic. *Environmental health perspectives*. 2011;119(6):860-
539 865.

540 58. Raaschou-Nielsen O, Sørensen M, Ketzel M, et al. Long-term exposure to traffic-related air
541 pollution and diabetes-associated mortality: A cohort study. *Diabetologia*. 2013;56(1):36-
542 46.

543 59. Sørensen M, Hoffmann B, Hvidberg M, et al. Long-term exposure to traffic-related air
544 pollution associated with blood pressure and self-reported hypertension in a Danish cohort.
545 *Environmental health perspectives*. 2012;120(3):418-424.

546 60. Seinfeld J, Pandis S. Atmospheric chemistry and physics. 1997. *New York*. Published online
547 2008.

548 61. Galvin M, Gaffney R, Corr B, Mays I, Hardiman O. From first symptoms to diagnosis of
549 amyotrophic lateral sclerosis: Perspectives of an Irish informal caregiver cohort—a
550 thematic analysis. *BMJ Open*. 2017;7(3). doi:10.1136/bmjopen-2016-014985

551 62. Dickerson AS, Hansen J, Kioumourtzoglou MA, Specht AJ, Gredal O, Weisskopf MG.
552 Study of occupation and amyotrophic lateral sclerosis in a Danish cohort. *Occup Environ*
553 *Med*. 2018;75(9):630-638. doi:10.1136/oemed-2018-105110

554 63. Roberts AL, Johnson NJ, Chen JT, Cudkowicz ME, Weisskopf MG. Race/ethnicity,
555 socioeconomic status, and ALS mortality in the United States. *Neurology*.
556 2016;87(22):2300-2308.

557 64. Bucher BT, Shi J, Pettit RJ, Ferraro J, Chapman WW, Gundlapalli A. Determination of
558 marital status of patients from structured and unstructured electronic healthcare data. In:
559 *AMIA Annual Symposium Proceedings*. Vol 2019. American Medical Informatics
560 Association; 2019:267.

561 65. Norman RE, Carpenter DO, Scott J, Brune MN, Sly PD. Environmental exposures: an
562 underrecognized contribution to noncommunicable diseases. *Reviews on environmental*
563 *health*. 2013;28(1):59-65.

564 66. Rothman KJ, Greenland S, Lash TL, others. *Modern Epidemiology*. Vol 3. Wolters Kluwer
565 Health/Lippincott Williams & Wilkins Philadelphia; 2008.

- 566 67. Gelman A, Carlin JB, Stern HS, Dunson DB, Vehtari A, Rubin DB. *Bayesian Data Analysis*,
567 *Third Edition*. CRC Press; 2013.
- 568 68. Mostofsky E, Schwartz J, Coull BA, et al. Modeling the association between particle
569 constituents of air pollution and health outcomes. *American journal of epidemiology*.
570 2012;176(4):317-326.
- 571 69. Thunis P, Degraeuwe B, Pisoni E, et al. PM2.5 source allocation in European cities: A
572 SHERPA modelling study. *Atmospheric Environment*. 2018;187:93-106.
- 573 70. Nuvolone D, Petri D, Voller F. The effects of ozone on human health. *Environmental*
574 *Science and Pollution Research*. 2018;25(9):8074-8088.
- 575 71. Martin R, Peters G, Wilkinson J. Symmetric decomposition of a positive definite matrix.
576 *Numerische Mathematik*. 1965;7(5):362-383.
- 577 72. Polson NG, Scott JG. On the half-Cauchy prior for a global scale parameter. *Bayesian*
578 *Analysis*. 2012;7(4):887-902.
- 579 73. Gelman A. Prior distributions for variance parameters in hierarchical models (comment on
580 article by Browne and Draper). *Bayesian Anal*. 2006;1(3):515-534. doi:10.1214/06-
581 BA117A
- 582 74. Lewandowski D, Kurowicka D, Joe H. Generating random correlation matrices based on
583 vines and extended onion method. *Journal of multivariate analysis*. 2009;100(9):1989-
584 2001.
- 585 75. Gelman A, Rubin DB. Inference from iterative simulation using multiple sequences.
586 *Statistical science*. 1992;7(4):457-472.
- 587 76. R Core Team. R: A language and environment for statistical computing. Published online
588 2013.
- 589 77. Povedano M, Saez M, Martinez-Matos JA, Barceló MA. Spatial assessment of the
590 association between long-term exposure to environmental factors and the occurrence of
591 amyotrophic lateral sclerosis in Catalonia, Spain: A population-based nested case-control
592 study. *Neuroepidemiology*. 2018;51(1-2):33-49.
- 593 78. von Schneidmesser E, Mar KA, Saar D. Black carbon in Europe: Targeting an air Pollutant
594 and climate forcer. Published online 2017.
- 595 79. Pamphlett R, Rikard-Bell A. Different occupations associated with amyotrophic lateral
596 sclerosis: Is diesel exhaust the link? *PloS One*. 2013;8(11):e80993.
- 597 80. Zhang R, Dai Y, Zhang X, et al. Reduced pulmonary function and increased pro-
598 inflammatory cytokines in nanoscale carbon black-exposed workers. *Part Fibre Toxicol*.
599 2014;11:73. doi:10.1186/s12989-014-0073-1

600 81. Gao X, Xu H, Shang J, et al. Ozonized carbon black induces mitochondrial dysfunction and
601 DNA damage. *Environ Toxicol.* 2017;32(3):944-955. doi:10.1002/tox.22295

602 82. Kyjovska ZO, Jacobsen NR, Saber AT, et al. DNA damage following pulmonary exposure
603 by instillation to low doses of carbon black (Printex 90) nanoparticles in mice. *Environ Mol*
604 *Mutagen.* 2015;56(1):41-49. doi:10.1002/em.21888

605 83. Benatar M, Turner MR, Wu J. Defining pre-symptomatic amyotrophic lateral sclerosis.
606 *Amyotrophic Lateral Sclerosis and Frontotemporal Degeneration.* 2019;20(5-6):303-309.

607 84. Benatar M, Wu J, McHutchison C, et al. Preventing amyotrophic lateral sclerosis: insights
608 from pre-symptomatic neurodegenerative diseases. Published online 2021.

609 85. Eisen A, Kiernan M, Mitsumoto H, Swash M. Amyotrophic lateral sclerosis: A long
610 preclinical period? *Journal of Neurology, Neurosurgery & Psychiatry.* 2014;85(11):1232-
611 1238.

612 86. Weisskopf MG, Webster TF. Trade-offs of personal vs. more proxy exposure measures in
613 environmental epidemiology. *Epidemiology (Cambridge, Mass).* 2017;28(5):635.

614 87. Carroll RJ, Ruppert D, Stefanski LA, Crainiceanu CM. *Measurement Error in Nonlinear*
615 *Models: A Modern Perspective.* CRC press; 2006.

616 88. Frisch M, Simonsen J. Marriage, cohabitation and mortality in Denmark: national cohort
617 study of 6.5 million persons followed for up to three decades (1982–2011). *International*
618 *Journal of Epidemiology.* 2013;42(2):559-578.

619 89. Bobb JF, Valeri L, Claus Henn B, et al. Bayesian kernel machine regression for estimating
620 the health effects of multi-pollutant mixtures. *Biostatistics.* 2015;16(3):493-508.

621

622 **Table 1.** Demographic characteristics of cases and controls for 5-year average exposure group.

Characteristic	Overall, N = 23,232 ^a	Case, N = 3,934 ^a	Control, N = 19,298 ^a
Average age (years)	66 (12)	66 (12)	66 (12)
Sex			
Female	10,973 (47%)	1,854 (47%)	9,119 (47%)
Male	12,259 (53%)	2,080 (53%)	10,179 (53%)
Socioeconomic status (SES)			
Group 1 (Highest)	2,337 (10%)	451 (11%)	1,886 (9.8%)
Group 2	2,839 (12%)	499 (13%)	2,340 (12%)
Group 3	4,360 (19%)	785 (20%)	3,575 (19%)
Group 4	6,598 (28%)	1,076 (27%)	5,522 (29%)
Group 5 (Lowest)	4,419 (19%)	717 (18%)	3,702 (19%)
Group 9 (Unknown)	2,679 (12%)	406 (10%)	2,273 (12%)
Place of birth			
Greater Copenhagen	4,858 (21%)	831 (21%)	4,027 (21%)
Big cities of Denmark	7,923 (34%)	1,357 (34%)	6,566 (34%)
Rest of Denmark	9,009 (39%)	1,548 (39%)	7,461 (39%)
Greenland	243 (1.0%)	53 (1.3%)	190 (1.0%)
Foreign	1,065 (4.6%)	122 (3.1%)	943 (4.9%)
Unknown	134 (0.6%)	23 (0.6%)	111 (0.6%)
Civil status			
Married	14,158 (61%)	2,411 (61%)	11,747 (61%)
Divorced	2,703 (12%)	433 (11%)	2,270 (12%)
Widowed	4,224 (18%)	726 (18%)	3,498 (18%)
Never married	2,147 (9.2%)	364 (9.3%)	1,783 (9.2%)
Last reported place of residence			
Greater Copenhagen	1,887 (8.1%)	335 (8.5%)	1,552 (8.0%)
Big cities of Denmark	9,385 (40%)	1,590 (40%)	7,795 (40%)
Rest of Denmark	11,954 (51%)	2,008 (51%)	9,946 (52%)
Greenland	6 (<0.1%)	1 (<0.1%)	5 (<0.1%)

^aMean (SD); n (%)

625 **Figure Captions**

626

627 **Figure 1.** Spearman correlation of 1-, 5-, and 10-year average pollutant concentrations.

628

629 **Figure 2.** Percentage change in odds of ALS diagnosis per 1-, 5- and 10-year average standard
630 deviation (SD) increase for each pollutant. Results are from the Bayesian hierarchical model
631 including each of EC, NO_x CO, and non-EC PM_{2.5} together, and were additionally adjusted by age,
632 sex, year of birth, vital status, socioeconomic status, civil status, last reported place of residence,
633 and place of birth.