



Aalto University
School of Science
and Technology

Transforming the array manifold Jacobian from polar to Cartesian coordinates

Robin Rajamäki

Department of Signal Processing and Acoustics
Aalto University, School of Electrical Engineering
robin.rajamaki@aalto.fi

Version 1.0, June 26, 2012

Content

1. Expressing the Fisher information matrix in Cartesian coordinates
2. Deriving the Jacobian for Cartesian coordinates
3. Expanding the Jacobian transform to the MIMO case
4. Summary

1. The FIM transform

- ▶ Let $\theta = [\phi \ \theta \ r]$ denote the polar parameters (*azimuth, elevation, distance*) and $\eta = [x \ y \ z]$ the corresponding coordinates in Cartesian space
- ▶ Assuming the FIM is expressed w.r.t. θ and we want to transform it to η , we could use the following transformation:¹

$$\mathcal{J}(\theta) = \mathcal{P}_{\eta\theta} \mathcal{J}(\eta) \mathcal{P}_{\eta\theta}^T$$

$$\Updownarrow$$

$$\mathcal{J}(\eta) = \mathcal{P}_{\eta\theta}^{-1} \mathcal{J}(\theta) \mathcal{P}_{\eta\theta}^{-T}$$

¹Andreas Richter, Estimation of Radio Channel Parameters: Models and Algorithms, 2005, p.57

1. The FIM transform

- Here the transform matrix is

$$\mathcal{P}_{\eta\theta} = \begin{bmatrix} \frac{\partial x}{\partial \phi} & \frac{\partial y}{\partial \phi} & \frac{\partial z}{\partial \phi} \\ \frac{\partial x}{\partial \theta} & \frac{\partial y}{\partial \theta} & \frac{\partial z}{\partial \theta} \\ \frac{\partial x}{\partial r} & \frac{\partial y}{\partial r} & \frac{\partial z}{\partial r} \end{bmatrix}$$

- And the parameters are

$$x = r \cdot \sin(\theta) \cos(\phi)$$

$$y = r \cdot \sin(\theta) \sin(\phi)$$

$$z = r \cdot \cos(\theta)$$

1. The FIM transform

- This leads to the final transformation matrix

$$\mathcal{P}_{\eta\theta} = \begin{bmatrix} -r \cdot \sin(\theta)\sin(\phi) & r \cdot \sin(\theta)\cos(\phi) & 0 \\ r \cdot \cos(\theta)\cos(\phi) & r \cdot \cos(\theta)\sin(\phi) & -r \cdot \sin(\theta) \\ \sin(\theta)\cos(\phi) & \sin(\theta)\sin(\phi) & \cos(\theta) \end{bmatrix}$$

- Now this can be applied to obtain the CRB in the Cartesian domain by taking the inverse of the transformed FIM

$$CRB = \mathcal{J}(\eta)^{-1} = \mathcal{P}_{\eta\theta}^T \mathcal{J}(\theta)^{-1} \mathcal{P}_{\eta\theta}$$

2. The Jacobian Transform - single-path

- ▶ Assume a transform of the array steering vector Jacobian is wanted (from θ to η), i.e.

$$\mathcal{D}(\eta) = \mathcal{D}(\theta)\mathbf{Q}$$

- ▶ \mathbf{Q} is the unknown transformation matrix
- ▶ And the Jacobians are

$$\mathcal{D}(\eta) = \begin{bmatrix} \frac{\partial \mathbf{b}(x, y, z)}{\partial x} & \frac{\partial \mathbf{b}(x, y, z)}{\partial y} & \frac{\partial \mathbf{b}(x, y, z)}{\partial z} \end{bmatrix}$$

$$\mathcal{D}(\theta) = \begin{bmatrix} \frac{\partial \mathbf{b}(\phi, \theta, r)}{\partial \phi} & \frac{\partial \mathbf{b}(\phi, \theta, r)}{\partial \theta} & \frac{\partial \mathbf{b}(\phi, \theta, r)}{\partial r} \end{bmatrix}$$

- ▶ Where \mathbf{b} is the array steering vector

2. The Jacobian Transform - single-path

- Consider one of the partial derivatives of θ

$$\frac{\partial \mathbf{b}(\phi, \theta, r)}{\partial \phi} = \frac{\partial \mathbf{b}(x(\phi, \theta, r), y(\phi, \theta, r), z(\phi, \theta, r))}{\partial \phi}$$

- Using the chain rule yields

$$\begin{aligned}\frac{\partial \mathbf{b}}{\partial \phi} &= \frac{\partial \mathbf{b}}{\partial x} \cdot \frac{\partial x}{\partial \phi} + \frac{\partial \mathbf{b}}{\partial y} \cdot \frac{\partial y}{\partial \phi} + \frac{\partial \mathbf{b}}{\partial z} \cdot \frac{\partial z}{\partial \phi} \\ &= \begin{bmatrix} \frac{\partial \mathbf{b}}{\partial x} & \frac{\partial \mathbf{b}}{\partial y} & \frac{\partial \mathbf{b}}{\partial z} \end{bmatrix} \cdot \begin{bmatrix} \frac{\partial x}{\partial \phi} & \frac{\partial y}{\partial \phi} & \frac{\partial z}{\partial \phi} \end{bmatrix}^T \\ &= \mathcal{D}(\eta) \cdot \begin{bmatrix} \frac{\partial x}{\partial \phi} & \frac{\partial y}{\partial \phi} & \frac{\partial z}{\partial \phi} \end{bmatrix}^T\end{aligned}$$

2. The Jacobian Transform - single-path

- ▶ Applying the chain rule on each of the partial derivatives of θ gives the following expression

$$\begin{aligned}\mathcal{D}(\theta) &= \mathcal{D}(\eta) \cdot \begin{bmatrix} \frac{\partial x}{\partial \phi} & \frac{\partial x}{\partial \theta} & \frac{\partial x}{\partial r} \\ \frac{\partial y}{\partial \phi} & \frac{\partial y}{\partial \theta} & \frac{\partial y}{\partial r} \\ \frac{\partial z}{\partial \phi} & \frac{\partial z}{\partial \theta} & \frac{\partial z}{\partial r} \end{bmatrix} \\ &= \mathcal{D}(\eta) \cdot \mathcal{P}_{\eta\theta}^T\end{aligned}$$

- ▶ And thus the wanted transformation is

$$\mathcal{D}(\eta) = \mathcal{D}(\theta) \cdot \mathcal{P}_{\eta\theta}^{-T}$$

- ▶ That is, the unknown matrix \mathbf{Q} is the inverse transpose of $\mathcal{P}_{\eta\theta}$, the transform matrix derived for the FIM

2. The Jacobian Transform - multi-path

- In case of K paths (or sources) the array steering matrix is

$$\mathbf{B} = [\mathbf{b}_1 \dots \mathbf{b}_K]$$

- This yields a result similar to the previously derived single-path case:

$$\mathcal{D}(\theta) = \begin{bmatrix} \frac{\partial \mathbf{B}}{\partial \phi} & \frac{\partial \mathbf{B}}{\partial \theta} & \frac{\partial \mathbf{B}}{\partial r} \end{bmatrix} = \begin{bmatrix} \frac{\partial \mathbf{B}}{\partial x} & \frac{\partial \mathbf{B}}{\partial y} & \frac{\partial \mathbf{B}}{\partial z} \end{bmatrix} \cdot \begin{bmatrix} \frac{\partial x}{\partial \phi} & \frac{\partial x}{\partial \theta} & \frac{\partial x}{\partial r} \\ \frac{\partial y}{\partial \phi} & \frac{\partial y}{\partial \theta} & \frac{\partial y}{\partial r} \\ \frac{\partial z}{\partial \phi} & \frac{\partial z}{\partial \theta} & \frac{\partial z}{\partial r} \end{bmatrix}$$

$$\Rightarrow \mathcal{D}(\eta) = \mathcal{D}(\theta) \cdot \mathcal{P}_{\eta\theta}^{-T}$$

2. The Jacobian Transform - multi-path

- If the actual values of the parameters describing a specific path are

$$\theta_k = [\phi_k \ \theta_k \ r_k], \text{ where } k = [1 \dots K],$$

the partial derivatives contained in the transform matrix, $\mathcal{P}_{\eta\theta}$, are diagonal matrices of the following form:

$$\frac{\partial \mathbf{x}}{\partial \phi} = \begin{bmatrix} \frac{\partial x(\phi_1)}{\partial \phi} & 0 & \dots & 0 \\ 0 & \frac{\partial x(\phi_2)}{\partial \phi} & 0 & 0 \\ \vdots & 0 & \ddots & 0 \\ 0 & \dots & 0 & \frac{\partial x(\phi_k)}{\partial \phi} \end{bmatrix}$$

where the diagonal contains the partial derivative with respect to the parameter in question (here ϕ) evaluated at the actual parameter values

2. The Jacobian Transform - multi-path

- ▶ Thus, in the general (multi-path) receiver only case, the transform matrix, $\mathcal{P}_{\eta\theta}$, is a $3K$ -by- $3K$ matrix consisting of 9 diagonal matrices, each corresponding to the partial derivative of a Cartesian parameter with respect to a polar parameter

3. The Jacobian transform for a MIMO system - single-path

- ▶ The previous cases assumed a single receiving array
- ▶ However, if the system is modeled with both a transmitting and a receiving array, the combined steering vector is the Kronecker product of the Tx and Rx steering vectors:

$$\mathbf{b}(\phi_R, \theta_R, r_R, \phi_T, \theta_T, r_T) = \mathbf{b}_R(\phi_R, \theta_R, r_R) \otimes \mathbf{b}_T(\phi_T, \theta_T, r_T)$$

- ▶ Here $\theta_R = [\phi_R \ \theta_R \ r_R]$ and $\theta_T = [\phi_T \ \theta_T \ r_T]$ define a single point in Cartesian space, $\eta = [x \ y \ z]$

3. The Jacobian transform for a MIMO system - single-path

- ▶ The whole system's Jacobian is defined as

$$\mathcal{D}(\theta_R, \theta_T) = \left[\frac{\partial \mathbf{b}_R \otimes \mathbf{b}_T}{\partial \phi_R} \quad \frac{\partial \mathbf{b}_R \otimes \mathbf{b}_T}{\partial \theta_R} \quad \cdots \quad \frac{\partial \mathbf{b}_R \otimes \mathbf{b}_T}{\partial \theta_T} \quad \frac{\partial \mathbf{b}_R \otimes \mathbf{b}_T}{\partial r_T} \right],$$

or more conveniently as

$$\mathcal{D}(\eta) = \left[\frac{\partial \mathbf{b}_R \otimes \mathbf{b}_T}{\partial x} \quad \frac{\partial \mathbf{b}_R \otimes \mathbf{b}_T}{\partial y} \quad \frac{\partial \mathbf{b}_R \otimes \mathbf{b}_T}{\partial z} \right]$$

- ▶ If the Rx array has M_R elements and the Tx array M_T elements, the whole system's Cartesian Jacobian is a $M_T M_R$ -by-3 matrix

3. The Jacobian transform for a MIMO system - single-path

- Using the chain rule, the Jacobian can be written as

$$\begin{aligned}\mathcal{D}(\eta) &= \left[\frac{\partial \mathbf{b}_R}{\partial x} \otimes \mathbf{b}_T + \mathbf{b}_R \otimes \frac{\partial \mathbf{b}_T}{\partial x} \quad \dots \quad \frac{\partial \mathbf{b}_R}{\partial z} \otimes \mathbf{b}_T + \mathbf{b}_R \otimes \frac{\partial \mathbf{b}_T}{\partial z} \right] \\ &= \left[\frac{\partial \mathbf{b}_R}{\partial x} \quad \frac{\partial \mathbf{b}_R}{\partial y} \quad \frac{\partial \mathbf{b}_R}{\partial z} \right] \otimes \mathbf{b}_T + \mathbf{b}_R \otimes \left[\frac{\partial \mathbf{b}_T}{\partial x} \quad \frac{\partial \mathbf{b}_T}{\partial y} \quad \frac{\partial \mathbf{b}_T}{\partial z} \right] \\ &= \mathcal{D}(\eta_R) \otimes \mathbf{b}_T + \mathbf{b}_R \otimes \mathcal{D}(\eta_T)\end{aligned}$$

- And the final transform becomes:

$$\mathcal{D}(\eta) = (\mathcal{D}(\theta_R) \cdot \mathcal{P}_{\eta\theta_R}^{-T}) \otimes \mathbf{b}_T + \mathbf{b}_R \otimes (\mathcal{D}(\theta_T) \cdot \mathcal{P}_{\eta\theta_T}^{-T})$$

3. The Jacobian transform for a MIMO system - multi-path

- ▶ In the multi-path case the combined steering matrix is the columnwise Kronecker product of the Tx and Rx steering matrices - that is, their Khatri-Rao product

$$\mathbf{B}(\phi_R, \theta_R, r_R, \phi_T, \theta_T, r_T) = \mathbf{B}_R(\phi_R, \theta_R, r_R) \diamond \mathbf{B}_T(\phi_T, \theta_T, r_T)$$

- ▶ Using previous results, the general MIMO Jacobian transform can be expressed as

$$\mathcal{D}(\eta) = \mathcal{D}(\eta_R) \diamond [\mathbf{B}_T \ \mathbf{B}_T \ \mathbf{B}_T] + [\mathbf{B}_R \ \mathbf{B}_R \ \mathbf{B}_R] \diamond \mathcal{D}(\eta_T)$$

where \diamond denotes the Khatri-Rao product and \mathbf{B}_T / \mathbf{B}_R are the Tx / Rx steering matrices

3. The Jacobian transform for a MIMO system - multi-path

► Proof:

$$\begin{aligned} \mathbf{B} &= \mathbf{B}_R \diamond \mathbf{B}_T, \\ \Rightarrow \mathcal{D}(\eta) &= \begin{bmatrix} \frac{\partial \mathbf{B}_R \diamond \mathbf{B}_T}{\partial x} & \frac{\partial \mathbf{B}_R \diamond \mathbf{B}_T}{\partial y} & \frac{\partial \mathbf{B}_R \diamond \mathbf{B}_T}{\partial z} \end{bmatrix} \\ &= \begin{bmatrix} \frac{\partial \mathbf{B}_R}{\partial x} \diamond \mathbf{B}_T & \frac{\partial \mathbf{B}_R}{\partial y} \diamond \mathbf{B}_T & \frac{\partial \mathbf{B}_R}{\partial z} \diamond \mathbf{B}_T \end{bmatrix} + \\ &\quad \begin{bmatrix} \mathbf{B}_R \diamond \frac{\partial \mathbf{B}_T}{\partial x} & \mathbf{B}_R \diamond \frac{\partial \mathbf{B}_T}{\partial y} & \mathbf{B}_R \diamond \frac{\partial \mathbf{B}_T}{\partial z} \end{bmatrix} \\ &= \begin{bmatrix} \frac{\partial \mathbf{B}_R}{\partial x} & \frac{\partial \mathbf{B}_R}{\partial y} & \frac{\partial \mathbf{B}_R}{\partial z} \end{bmatrix} \diamond [\mathbf{B}_T \ \mathbf{B}_T \ \mathbf{B}_T] + \\ &\quad [\mathbf{B}_R \ \mathbf{B}_R \ \mathbf{B}_R] \diamond \begin{bmatrix} \frac{\partial \mathbf{B}_T}{\partial x} & \frac{\partial \mathbf{B}_T}{\partial y} & \frac{\partial \mathbf{B}_T}{\partial z} \end{bmatrix} \quad \square \end{aligned}$$

4. Summary

- The transformation matrix:

$$\mathcal{P}_{\eta\theta} = \begin{bmatrix} \frac{\partial x}{\partial \phi} & \frac{\partial y}{\partial \phi} & \frac{\partial z}{\partial \phi} \\ \frac{\partial x}{\partial \theta} & \frac{\partial y}{\partial \theta} & \frac{\partial z}{\partial \theta} \\ \frac{\partial x}{\partial r} & \frac{\partial y}{\partial r} & \frac{\partial z}{\partial r} \end{bmatrix}$$

- The FIM transform:

$$\mathcal{J}(\eta) = \mathcal{P}_{\eta\theta}^{-1} \mathcal{J}(\theta) \mathcal{P}_{\eta\theta}^{-T}$$

- The Jacobian transform (receiver only):

$$\mathcal{D}(\eta) = \mathcal{D}(\theta) \cdot \mathcal{P}_{\eta\theta}^{-T}$$

- The Jacobian transform (MIMO):

$$\mathcal{D}(\eta) = \mathcal{D}(\eta_R) \diamond [\mathbf{B}_T \ \mathbf{B}_T \ \mathbf{B}_T] + [\mathbf{B}_R \ \mathbf{B}_R \ \mathbf{B}_R] \diamond \mathcal{D}(\eta_T)$$