

# Optimal minimax bounds for the Navier-Stokes equations and other infinite dimensional dissipative systems

Ricardo M. S. Rosa

Instituto de Matemática  
Universidade Federal do Rio de Janeiro

Special Session on Fluid Dynamics  
ICMC Summer Meeting on Differential Equations  
Chapter 2021 - Feb 1-3

February 2, 2021



UNIVERSIDADE FEDERAL  
DO RIO DE JANEIRO



**INSTITUTO DE MATEMÁTICA**  
Universidade Federal do Rio de Janeiro

Based on

“Optimal minimax bounds for time and ensemble averages of dissipative infinite-dimensional systems with applications to the incompressible Navier-Stokes equations”

Joint work with Roger Temam (Indiana University)

Submitted to Pure and Applied Functional Analysis

Dedicated to the memory of Ciprian Foias

Estimates are desirable for many purposes:

- ▶ existence and uniqueness of solutions;
- ▶ blow-up of solutions
- ▶ stabilization results
- ▶ estimating dimensions of invariant sets;
- ▶ localizing local or global attractors;
- ▶ assessing real quantities
  - ▶ energy, drag coefficient, mechanical stress, chemical concentration, infected population, pharmaceutical dosage, etc.

Usual techniques:

- ▶ Energy-type estimates
- ▶ Variational inequalities

Usual techniques:

- ▶ Energy-type estimates
- ▶ Variational inequalities

More recently:

- ▶ Mini-max estimates...
- ▶ ... theoretically gives optimal results!
- ▶ ... numerically gives sharp results often with less cost!

Usual techniques:

- ▶ Energy-type estimates
- ▶ Variational inequalities

More recently:

- ▶ Mini-max estimates...
- ▶ ... theoretically gives optimal results!
- ▶ ... numerically gives sharp results often with less cost!

Numerically approaches:

- ▶ Monte-Carlo estimates for classical evolution methods
- ▶ Linear programming/sums of squares for mini-max estimate

Suppose we are interested on a differential equation on a phase space  $X$ ,

$$\frac{du}{dt} = F(u),$$

and on estimating the **mean** of a quantity  $\phi : X \rightarrow \mathbb{R}$  over a **compact** and **positively-invariant** subset  $B \subset X$ :

$$\sup_{u_0 \in B} \limsup_{T \rightarrow \infty} \frac{1}{T} \int_0^T \phi(u(t)) \, dt.$$

Suppose we are interested on a differential equation on a phase space  $X$ ,

$$\frac{du}{dt} = F(u),$$

and on estimating the **mean** of a quantity  $\phi : X \rightarrow \mathbb{R}$  over a **compact** and **positively-invariant** subset  $B \subset X$ :

$$\sup_{u_0 \in B} \limsup_{T \rightarrow \infty} \frac{1}{T} \int_0^T \phi(u(t)) \, dt.$$

**In theory**, can use energy-type estimates and so-on



Suppose we are interested on a differential equation on a phase space  $X$ ,

$$\frac{du}{dt} = F(u),$$

and on estimating the **mean** of a quantity  $\phi : X \rightarrow \mathbb{R}$  over a **compact** and **positively-invariant** subset  $B \subset X$ :

$$\sup_{u_0 \in B} \limsup_{T \rightarrow \infty} \frac{1}{T} \int_0^T \phi(u(t)) \, dt.$$

**In theory**, can use energy-type estimates and so-on

**Numerically**, can use Monte-Carlo (solve multiple solutions for long time intervals, using a decent integration method, ...)

Suppose we are interested on a differential equation on a phase space  $X$ ,

$$\frac{du}{dt} = F(u),$$

and on estimating the **mean** of a quantity  $\phi : X \rightarrow \mathbb{R}$  over a **compact** and **positively-invariant** subset  $B \subset X$ :

$$\sup_{u_0 \in B} \limsup_{T \rightarrow \infty} \frac{1}{T} \int_0^T \phi(u(t)) \, dt.$$

**In theory**, can use energy-type estimates and so-on

**Numerically**, can use Monte-Carlo (solve multiple solutions for long time intervals, using a decent integration method, ...)

Or, work only on the phase space, without solving the equation...

# Auxilliary function

We look for bound  $C$  so that  $\bar{\phi} \leq C$

## Auxilliary function

We look for bound  $C$  so that  $\bar{\phi} \leq C$

For any  $\mathcal{C}^1$  function  $V : X \rightarrow \mathbb{R}$ , using that  $\dot{u} = F(u)$ ,

# Auxilliary function

We look for bound  $C$  so that  $\bar{\phi} \leq C$

For any  $\mathcal{C}^1$  function  $V : X \rightarrow \mathbb{R}$ , using that  $\dot{u} = F(u)$ ,

$$\frac{1}{T} \int_0^T F \cdot \nabla V(u(t)) \, dt$$

# Auxilliary function

We look for bound  $C$  so that  $\bar{\phi} \leq C$

For any  $\mathcal{C}^1$  function  $V : X \rightarrow \mathbb{R}$ , using that  $\dot{u} = F(u)$ ,

$$\frac{1}{T} \int_0^T F \cdot \nabla V(u(t)) \, dt = \frac{V(u(T)) - V(u(0))}{T}.$$

## Auxilliary function

We look for bound  $C$  so that  $\bar{\phi} \leq C$

For any  $\mathcal{C}^1$  function  $V : X \rightarrow \mathbb{R}$ , using that  $\dot{u} = F(u)$ ,

$$\frac{1}{T} \int_0^T F \cdot \nabla V(u(t)) \, dt = \frac{V(u(T)) - V(u(0))}{T} \rightarrow 0, \quad \text{as } T \rightarrow \infty.$$

# Auxilliary function

We look for bound  $C$  so that  $\bar{\phi} \leq C$

For any  $\mathcal{C}^1$  function  $V : X \rightarrow \mathbb{R}$ , using that  $\dot{u} = F(u)$ ,

$$\frac{1}{T} \int_0^T F \cdot \nabla V(u(t)) \, dt = \frac{V(u(T)) - V(u(0))}{T} \rightarrow 0, \quad \text{as } T \rightarrow \infty.$$



# Auxilliary function

We look for bound  $C$  so that  $\bar{\phi} \leq C$

For any  $\mathcal{C}^1$  function  $V : X \rightarrow \mathbb{R}$ , using that  $\dot{u} = F(u)$ ,

$$\frac{1}{T} \int_0^T F \cdot \nabla V(u(t)) \, dt = \frac{V(u(T)) - V(u(0))}{T} \rightarrow 0, \quad \text{as } T \rightarrow \infty.$$

Hence,  $\overline{F \cdot \nabla V} = 0$ , and get the equivalent statement

$$\overline{\phi + F \cdot \nabla V} \leq C$$

# Auxilliary function

We look for bound  $C$  so that  $\bar{\phi} \leq C$

For any  $\mathcal{C}^1$  function  $V : X \rightarrow \mathbb{R}$ , using that  $\dot{u} = F(u)$ ,

$$\frac{1}{T} \int_0^T F \cdot \nabla V(u(t)) \, dt = \frac{V(u(T)) - V(u(0))}{T} \rightarrow 0, \quad \text{as } T \rightarrow \infty.$$

Hence,  $\overline{F \cdot \nabla V} = 0$ , and get the equivalent statement

$$\overline{\phi + F \cdot \nabla V} \leq C$$

Now, if we find a **auxilliary function**  $V$  such that

$$\phi(u) + F(u) \cdot \nabla V(u) \leq C, \quad \forall u \in B,$$

then

$$\overline{\phi + F \cdot \nabla V} \leq C, \quad \text{hence } \bar{\phi} \leq C.$$

## The problem

Find  $V$  s.th.  $\phi(u) + F(u) \cdot \nabla V(u) \leq C$ ,  $\forall u \in B$ , for *best possible*  $C$ ,

can be written as a convex optimization problem

$$\sup \bar{\phi} \leq \inf_{(C, V) \in \mathbb{R} \times \mathcal{C}^1, S_{C, V}(u) \geq 0} C,$$

where  $S_{C, V}(u) = C - \phi(u) - F(u) \cdot \nabla V(u)$ .

(Optimization of linear map  $(C, V) \mapsto C$  over a convex set, since  $\mathbb{R} \times \mathcal{C}^1$  is convex and  $S_{C, V}(u)$  is linear in  $C$  and  $V$ .)

If  $\phi, F$  **polynomials**, we narrow minimization over **polynomials**  $V$  such that

$$S_{C,V}(u) = C - \phi(u) - F(u) \cdot \nabla V(u) = \text{SoS},$$

where  $\text{SoS} = \sum_i p_i(u)^2$ , for other polynomials  $p_i$ .

# Optimization with Sum of Squares (SoS)

If  $\phi$ ,  $F$  **polynomials**, we narrow minimization over **polynomials**  $V$  such that

$$S_{C,V}(u) = C - \phi(u) - F(u) \cdot \nabla V(u) = \text{SoS},$$

where  $\text{SoS} = \sum_i p_i(u)^2$ , for other polynomials  $p_i$ .

Hence, we relax the problem to ...

If  $\phi, F$  **polynomials**, we narrow minimization over **polynomials**  $V$  such that

$$S_{C,V}(u) = C - \phi(u) - F(u) \cdot \nabla V(u) = \text{SoS},$$

where  $\text{SoS} = \sum_i p_i(u)^2$ , for other polynomials  $p_i$ .

Hence, we relax the problem to ...

$$\sup \bar{\phi} \leq \inf_{(C,V) \in \mathbb{R} \times \mathcal{P}_m(X), S_{C,V}(u) = \text{SoS}} C.$$

If  $\phi, F$  **polynomials**, we narrow minimization over **polynomials**  $V$  such that

$$S_{C,V}(u) = C - \phi(u) - F(u) \cdot \nabla V(u) = \text{SoS},$$

where  $\text{SoS} = \sum_i p_i(u)^2$ , for other polynomials  $p_i$ .

Hence, we relax the problem to ...

$$\sup \bar{\phi} \leq \inf_{(C,V) \in \mathbb{R} \times \mathcal{P}_m(X), S_{C,V}(u) = \text{SoS}} C.$$

Reducing the set of possible  $V$ 's may increase the upper bound estimate, but at the advantage of turning it from NP-hard into a P-complete problem.

If  $\phi, F$  **polynomials**, we narrow minimization over **polynomials**  $V$  such that

$$S_{C,V}(u) = C - \phi(u) - F(u) \cdot \nabla V(u) = \text{SoS},$$

where  $\text{SoS} = \sum_i p_i(u)^2$ , for other polynomials  $p_i$ .

Hence, we relax the problem to ...

$$\sup \bar{\phi} \leq \inf_{(C,V) \in \mathbb{R} \times \mathcal{P}_m(X), S_{C,V}(u) = \text{SoS}} C.$$

Reducing the set of possible  $V$ 's may increase the upper bound estimate, but at the advantage of turning it from NP-hard into a P-complete problem.


Question: How large is the set of SoS?



- ▶ Hilbert's 17th problem (1900) is about decomposing positive polynomials into SoS of rational functionals.
- ▶ Hilbert had already noticed not all positive polynomials are SoSs.
- ▶ Reznik (2000) - survey of SoS and Hilbert's 17th problem
- ▶ Shor 1980s, 1990s, Choi, Lam, Reznik 1990s - SoS polynomial decomposition
- ▶ Parrilo (2000s) - several applications: Lyapunov functions, control, etc.
- ▶ Several SoS MATLAB toolbox solvers (2000s)
- ▶ Papachristodoulou, Peet (2006) - applications to PDEs
- ▶ Yu, Kashima, Imura (2008)- local stability of 2D fluid flows
- ▶ Goulart, Chernyshenko (2012) - global stability of fluid flows
- ▶ Fantuzzi, Goluskin, Doering, Goulart, Chernyshenko, Huang, Papachristodoulou (2010s) ...


# Bounds for the van der Pol limit cycle

From “Bounds for Deterministic and Stochastic Dynamical Systems using Sum-of-Squares Optimization”, by G. Fantuzzi, D. Goluskin, D. Huang, and S. I. Chernyshenko, in SIAM J. Applied Dynamical Systems, Vol. 15, No. 4, pp. 1962–1988.




**Figure 2.** Optimal upper bounds on  $\overline{\varphi} = \overline{x^2 + y^2}$  for the van der Pol oscillator computed with the upper bound problem of (2.9) for different degrees of  $V_u$ . The time average  $\overline{\varphi}$  obtained by numerical integration (N.I.) of the system is also shown.


# Auxiliary function with degree 6 for van der Pol



# Auxiliary function with degree 8 for van der Pol



# Auxiliary function with degree 10 for van der Pol



## The problem

Find  $V$  and the *best possible*  $C$  s.th.  $\phi(u) + F(u) \cdot \nabla V(u) \leq C$ ,  $\forall u \in B$ ,  
can also be written as the minimax problem

$$\sup_{u_0 \in B} \bar{\phi}(u_0) \leq \min_{V \in C^1(B)} \max_{u \in B} \{\phi(u) + F(u) \cdot \nabla V(u)\}.$$

# Optimality of the minimax formula

Tobasco-Goluskin-Doering (2018):

It turns out, the minimax formula is optimal and is achieved!

$$\max_{u_0 \in B} \bar{\phi}(u_0) = \min_{V \in \mathcal{C}^1(B)} \max_{u \in B} \{ \phi(u) + F(u) \cdot \nabla V(u) \}.$$

# Optimality of the minimax formula

Tobasco-Goluskin-Doering (2018):

It turns out, the minimax formula is optimal and is achieved!

$$\max_{u_0 \in B} \bar{\phi}(u_0) = \min_{V \in \mathcal{C}^1(B)} \max_{u \in B} \{ \phi(u) + F(u) \cdot \nabla V(u) \}.$$

Conditions:

- ▶  $X = \mathbb{R}^n$
- ▶  $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$  continuously differentiable
- ▶  $B \subset \mathbb{R}^n$  compact
- ▶  $B$  positively invariant
- ▶  $\phi : B \rightarrow \mathbb{R}$  continuous



# Optimality of the minimax formula

Tobasco-Goluskin-Doering (2018):

It turns out, the minimax formula is optimal and is achieved!

$$\max_{u_0 \in B} \bar{\phi}(u_0) = \min_{V \in C^1(B)} \max_{u \in B} \{ \phi(u) + F(u) \cdot \nabla V(u) \}.$$

Conditions:

- ▶  $X = \mathbb{R}^n$
- ▶  $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$  continuously differentiable
- ▶  $B \subset \mathbb{R}^n$  compact
- ▶  $B$  positively invariant
- ▶  $\phi : B \rightarrow \mathbb{R}$  continuous

Questions:

- ▶ What about infinite-dimensional systems? Like 2D NSE.
- ▶ What about 3DNSE?

# Proof of optimality in the finite-dimensional case

Tobasco-Goluskin-Doering (2018):

$$\max_{u_0 \in B} \bar{\phi}(u_0) =$$

# Proof of optimality in the finite-dimensional case

Tobasco-Goluskin-Doering (2018):

$$\max_{u_0 \in B} \bar{\phi}(u_0) = \max_{u_0 \in B} \limsup_{T \rightarrow \infty} \frac{1}{T} \int_0^T \phi(u(t)) \, dt \quad (\text{definition})$$

# Proof of optimality in the finite-dimensional case

Tobasco-Goluskin-Doering (2018):

$$\begin{aligned}\max_{u_0 \in B} \bar{\phi}(u_0) &= \max_{u_0 \in B} \limsup_{T \rightarrow \infty} \frac{1}{T} \int_0^T \phi(u(t)) \, dt \\ &= \max_{\mu \in \mathcal{P}_{\text{inv}}(B)} \int_B \phi \, d\mu \quad (\text{maximum at extreme, ergodic meas.})\end{aligned}$$

# Proof of optimality in the finite-dimensional case

Tobasco-Goluskin-Doering (2018):

$$\begin{aligned}\max_{u_0 \in B} \bar{\phi}(u_0) &= \max_{u_0 \in B} \limsup_{T \rightarrow \infty} \frac{1}{T} \int_0^T \phi(u(t)) \, dt \\ &= \max_{\mu \in \mathcal{P}_{\text{inv}}(B)} \int_B \phi \, d\mu \\ &= \max_{\mu \in \mathcal{P}_{\mathcal{E}\text{-inv}}(B)} \int_B \phi \, d\mu \quad (\text{Lagrangian} = \text{Eulerian invariance } \langle F \cdot \nabla V \rangle = 0)\end{aligned}$$

# Proof of optimality in the finite-dimensional case

Tobasco-Goluskin-Doering (2018):

$$\begin{aligned}\max_{u_0 \in B} \bar{\phi}(u_0) &= \max_{u_0 \in B} \limsup_{T \rightarrow \infty} \frac{1}{T} \int_0^T \phi(u(t)) \, dt \\ &= \max_{\mu \in \mathcal{P}_{\text{inv}}(B)} \int_B \phi \, d\mu \\ &= \max_{\mu \in \mathcal{P}_{\mathcal{E}\text{-inv}}(B)} \int_B \phi \, d\mu \\ &= \sup_{\mu \in \mathcal{P}(B)} \inf_{V \in \mathcal{C}^1(B)} \int_B \phi + F \cdot \nabla V \, d\mu \left( \inf_V \langle F \cdot \nabla V \rangle = \begin{cases} 0, & \mu \in \mathcal{P}_{\mathcal{E}\text{-inv}}(B) \\ -\infty, & \mu \notin \mathcal{P}_{\mathcal{E}\text{-inv}}(B) \end{cases} \right)\end{aligned}$$

# Proof of optimality in the finite-dimensional case

Tobasco-Goluskin-Doering (2018):

$$\begin{aligned}\max_{u_0 \in B} \bar{\phi}(u_0) &= \max_{u_0 \in B} \limsup_{T \rightarrow \infty} \frac{1}{T} \int_0^T \phi(u(t)) \, dt \\&= \max_{\mu \in \mathcal{P}_{\text{inv}}(B)} \int_B \phi \, d\mu \\&= \max_{\mu \in \mathcal{P}_{\mathcal{E}\text{-inv}}(B)} \int_B \phi \, d\mu \\&= \sup_{\mu \in \mathcal{P}(B)} \inf_{V \in \mathcal{C}^1(B)} \int_B \phi + F \cdot \nabla V \, d\mu \\&= \inf_{V \in \mathcal{C}^1(B)} \sup_{\mu \in \mathcal{P}(B)} \int_B \phi + F \cdot \nabla V \, d\mu \quad (\text{minimax principle})\end{aligned}$$

# Proof of optimality in the finite-dimensional case

Tobasco-Goluskin-Doering (2018):

$$\begin{aligned}\max_{u_0 \in B} \bar{\phi}(u_0) &= \max_{u_0 \in B} \limsup_{T \rightarrow \infty} \frac{1}{T} \int_0^T \phi(u(t)) \, dt \\&= \max_{\mu \in \mathcal{P}_{\text{inv}}(B)} \int_B \phi \, d\mu \\&= \max_{\mu \in \mathcal{P}_{\mathcal{E}\text{-inv}}(B)} \int_B \phi \, d\mu \\&= \sup_{\mu \in \mathcal{P}(B)} \inf_{V \in \mathcal{C}^1(B)} \int_B \phi + F \cdot \nabla V \, d\mu \\&= \inf_{V \in \mathcal{C}^1(B)} \sup_{\mu \in \mathcal{P}(B)} \int_B \phi + F \cdot \nabla V \, d\mu \\&= \inf_{V \in \mathcal{C}^1(B)} \max_{u \in B} \{ \phi(u) + F(u) \cdot \nabla V(u) \} \quad (\text{extreme at Dirac delta})\end{aligned}$$



$X$  Hausdorff space;  $W$  Banach space;  $W \subset X \subset W'$  continuous.

## Definition

A **cylindrical test functional** on  $W'$  is  $\Psi : W' \rightarrow \mathbb{R}$  of the form

$$\Psi(u) = \psi(\langle u, w_1 \rangle_{W', W}, \dots, \langle u, w_m \rangle_{W', W}), \quad \forall u \in W',$$

where  $w_1, \dots, w_m \in W$ ,  $m \in \mathbb{N}$ , and  $\psi \in \mathcal{C}_c^1(\mathbb{R}^m)$ . Space  $\mathcal{T}^{\text{cyl}}(W')$ .

## Definition

A **weak stationary statistical solution** is a Borel probability measure  $\mu$  on  $X$  such that, for any  $\mathcal{T}^{\text{cyl}}(W')$ , the map  $u \mapsto \langle F(u), \Psi'(u) \rangle_{W', W}$  is  $\mu$ -integrable and

$$\int_X \langle F(u), \Psi'(u) \rangle_{W', W} d\mu(u) = 0.$$

Space  $\mathcal{P}_{\text{wsss}}(E)$  of those carried by Borel subset  $E$ .

# Infinite-dimensional dissipative semigroup



$$\sup_{u_0 \in B} \bar{\phi}(u_0) =$$

$$\sup_{u_0 \in B} \bar{\phi}(u_0) = \sup_{u_0 \in B} \limsup_{T \rightarrow \infty} \frac{1}{T} \int_0^T \phi(u(t)) \, dt \quad (\text{definition})$$

$$\begin{aligned}\sup_{u_0 \in B} \bar{\phi}(u_0) &= \sup_{u_0 \in B} \limsup_{T \rightarrow \infty} \frac{1}{T} \int_0^T \phi(u(t)) \, dt \\ &= \max_{\mu \in \mathcal{P}_{\text{inv}}(B)} \int_B \phi \, d\mu \quad (\text{maximum at extreme, ergodic meas.})\end{aligned}$$

$$\begin{aligned}\sup_{u_0 \in B} \bar{\phi}(u_0) &= \sup_{u_0 \in B} \limsup_{T \rightarrow \infty} \frac{1}{T} \int_0^T \phi(u(t)) \, dt \\ &= \max_{\mu \in \mathcal{P}_{\text{inv}}(B)} \int_B \phi \, d\mu \\ &= \max_{\mu \in \mathcal{P}_{\text{inv}}(B \cap K)} \int_{B \cap K} \phi \, d\mu \quad (K \text{ compact attracting; } B \text{ normal})\end{aligned}$$

$$\begin{aligned}\sup_{u_0 \in B} \bar{\phi}(u_0) &= \sup_{u_0 \in B} \limsup_{T \rightarrow \infty} \frac{1}{T} \int_0^T \phi(u(t)) \, dt \\ &= \max_{\mu \in \mathcal{P}_{\text{inv}}(B)} \int_B \phi \, d\mu \\ &= \max_{\mu \in \mathcal{P}_{\text{inv}}(B \cap K)} \int_{B \cap K} \phi \, d\mu \\ &= \max_{\mu \in \mathcal{P}_{\text{wsss}}(B \cap K)} \int_{B \cap K} \phi \, d\mu \quad (\mathcal{P}_{\text{inv}}(B \cap K) \stackrel{?}{=} \mathcal{P}_{\text{wsss}}(B \cap K))\end{aligned}$$

$$\begin{aligned}
 \sup_{u_0 \in B} \bar{\phi}(u_0) &= \sup_{u_0 \in B} \limsup_{T \rightarrow \infty} \frac{1}{T} \int_0^T \phi(u(t)) \, dt \\
 &= \max_{\mu \in \mathcal{P}_{\text{inv}}(B)} \int_B \phi \, d\mu \\
 &= \max_{\mu \in \mathcal{P}_{\text{inv}}(B \cap K)} \int_{B \cap K} \phi \, d\mu \\
 &= \max_{\mu \in \mathcal{P}_{\text{wsss}}(B \cap K)} \int_{B \cap K} \phi \, d\mu \\
 &= \sup_{\mu \in \mathcal{P}(B \cap K)} \inf_{V \in \mathcal{T}_{\text{cyl}}} \int_{B \cap K} \phi + F \cdot \nabla V \, d\mu \left( \inf_V = \begin{cases} 0, & \mu \in \mathcal{P}_{\text{wsss}}(B \cap K) \\ -\infty, & \mu \notin \mathcal{P}_{\text{wsss}}(B \cap K) \end{cases} \right)
 \end{aligned}$$

$$\begin{aligned}
 \sup_{u_0 \in B} \bar{\phi}(u_0) &= \sup_{u_0 \in B} \limsup_{T \rightarrow \infty} \frac{1}{T} \int_0^T \phi(u(t)) \, dt \\
 &= \max_{\mu \in \mathcal{P}_{\text{inv}}(B)} \int_B \phi \, d\mu \\
 &= \max_{\mu \in \mathcal{P}_{\text{inv}}(B \cap K)} \int_{B \cap K} \phi \, d\mu \\
 &= \max_{\mu \in \mathcal{P}_{\text{wss}}(B \cap K)} \int_{B \cap K} \phi \, d\mu \\
 &= \sup_{\mu \in \mathcal{P}(B \cap K)} \inf_{V \in \mathcal{T}_{\text{cyl}}} \int_{B \cap K} \phi + F \cdot \nabla V \, d\mu \\
 &= \inf_{V \in \mathcal{T}_{\text{cyl}}} \sup_{\mu \in \mathcal{P}(B \cap K)} \int_{B \cap K} \phi + F \cdot \nabla V \, d\mu \quad (\text{minimax principle})
 \end{aligned}$$



$$\begin{aligned}
 \sup_{u_0 \in B} \bar{\phi}(u_0) &= \sup_{u_0 \in B} \limsup_{T \rightarrow \infty} \frac{1}{T} \int_0^T \phi(u(t)) \, dt \\
 &= \max_{\mu \in \mathcal{P}_{\text{inv}}(B)} \int_B \phi \, d\mu \\
 &= \max_{\mu \in \mathcal{P}_{\text{inv}}(B \cap K)} \int_{B \cap K} \phi \, d\mu \\
 &= \max_{\mu \in \mathcal{P}_{\text{wss}}(B \cap K)} \int_{B \cap K} \phi \, d\mu \\
 &= \sup_{\mu \in \mathcal{P}(B \cap K)} \inf_{V \in \mathcal{T}_{\text{cyl}}} \int_{B \cap K} \phi + F \cdot \nabla V \, d\mu \\
 &= \inf_{V \in \mathcal{T}_{\text{cyl}}} \sup_{\mu \in \mathcal{P}(B \cap K)} \int_{B \cap K} \phi + F \cdot \nabla V \, d\mu \\
 &= \inf_{V \in \mathcal{T}_{\text{cyl}}} \max_{u \in B \cap K} \{ \phi(u) + F(u) \cdot \nabla V(u) \} \quad (\text{extreme at Dirac delta})
 \end{aligned}$$



$$\sup_{u \in \mathcal{U}(B)} \bar{\phi}(u) =$$



$$\sup_{u \in \mathcal{U}(B)} \bar{\phi}(u) = \sup_{u \in \mathcal{U}(B)} \limsup_{T \rightarrow \infty} \frac{1}{T} \int_0^T \phi(u(t)) \, dt \quad (\text{definition})$$

$$\sup_{u \in \mathcal{U}(B)} \bar{\phi}(u) = \sup_{u \in \mathcal{U}(B)} \limsup_{T \rightarrow \infty} \frac{1}{T} \int_0^T \phi(u(t)) \, dt$$
$$\leq \max_{\mu \in \mathcal{P}_{\text{wss}}(B)} \int_{B \cap K} \phi \, d\mu \quad (\text{Bogoliubov-Krylov; } K \text{ comp. attr.; } B \text{ normal})$$

$$\begin{aligned}
 \sup_{u \in \mathcal{U}(B)} \bar{\phi}(u) &= \sup_{u \in \mathcal{U}(B)} \limsup_{T \rightarrow \infty} \frac{1}{T} \int_0^T \phi(u(t)) \, dt \\
 &\leq \max_{\mu \in \mathcal{P}_{\text{wsss}}(B)} \int_{B \cap K} \phi \, d\mu \\
 &= \sup_{\mu \in \mathcal{P}(B \cap K)} \inf_{V \in \mathcal{T}_{\text{cyl}}} \int_{B \cap K} \phi + F \cdot \nabla V \, d\mu \left( \inf_V = \begin{cases} 0, & \mu \in \mathcal{P}_{\text{wsss}}(B \cap K) \\ -\infty, & \mu \notin \mathcal{P}_{\text{wsss}}(B \cap K) \end{cases} \right)
 \end{aligned}$$

$$\begin{aligned}
 \sup_{u \in \mathcal{U}(B)} \bar{\phi}(u) &= \sup_{u \in \mathcal{U}(B)} \limsup_{T \rightarrow \infty} \frac{1}{T} \int_0^T \phi(u(t)) \, dt \\
 &\leq \max_{\mu \in \mathcal{P}_{\text{wss}}(B)} \int_{B \cap K} \phi \, d\mu \\
 &= \sup_{\mu \in \mathcal{P}(B \cap K)} \inf_{V \in \mathcal{T}_{\text{cyl}}} \int_{B \cap K} \phi + F \cdot \nabla V \, d\mu \\
 &= \inf_{V \in \mathcal{T}_{\text{cyl}}} \sup_{\mu \in \mathcal{P}(B \cap K)} \int_{B \cap K} \phi + F \cdot \nabla V \, d\mu \quad (\text{minimax principle})
 \end{aligned}$$

$$\begin{aligned}
 \sup_{u \in \mathcal{U}(B)} \bar{\phi}(u) &= \sup_{u \in \mathcal{U}(B)} \limsup_{T \rightarrow \infty} \frac{1}{T} \int_0^T \phi(u(t)) \, dt \\
 &\leq \max_{\mu \in \mathcal{P}_{\text{wss}}(B)} \int_{B \cap K} \phi \, d\mu \\
 &= \sup_{\mu \in \mathcal{P}(B \cap K)} \inf_{V \in \mathcal{T}_{\text{cyl}}} \int_{B \cap K} \phi + F \cdot \nabla V \, d\mu \\
 &= \inf_{V \in \mathcal{T}_{\text{cyl}}} \sup_{\mu \in \mathcal{P}(B \cap K)} \int_{B \cap K} \phi + F \cdot \nabla V \, d\mu \\
 &= \inf_{V \in \mathcal{T}_{\text{cyl}}} \max_{u \in B \cap K} \{ \phi(u) + F(u) \cdot \nabla V(u) \} \quad (\text{extreme at Dirac delta})
 \end{aligned}$$

## Theorem

Suppose  $K$  is a compact and metrizable subset of  $X$  and  $F$  is continuous on  $K$ . Assume the set  $\mathcal{P}_{wss}(K)$  of weak stationary statistical solutions carried by  $K$  is not empty. Let  $\phi \in \mathcal{C}(K)$ . Then,

$$\max_{\mu \in \mathcal{P}_{wss}(K)} \int_K \phi(u) d\mu(u) = \inf_{\Psi \in \mathcal{T}_{cyl}(W')} \max_{u \in K} \{ \phi(u) + \langle F(u), \Psi'(u) \rangle_{W', W} \}.$$



## Theorem

Let  $B$  be a positively invariant set for  $\{S(t)\}_{t \geq 0}$  which is closed in  $X$  and normal, and suppose that there exists a compact and metrizable subset  $K$  of  $X$  which attracts the points of  $B$ . Suppose  $F$  is continuous on  $K$ . Let  $\phi \in \mathcal{C}_b(B)$ . Then,

$$\max_{\mu \in \mathcal{P}_{\text{wss}}(K)} \int_K \phi(u) d\mu(u) = \inf_{\Psi \in \mathcal{T}_{\text{cyl}}(W')} \max_{u \in B \cap K} \{ \phi(u) + \langle F(u), \Psi'(u) \rangle_{W', W} \}.$$

Suppose, further, that  $\mathcal{P}_{\text{wss}}(K) = \mathcal{P}_{\text{inv}}(B \cap K)$ . Then

$$\begin{aligned} \max_{u_0 \in B} \limsup_{T \rightarrow \infty} \frac{1}{T} \int_0^T \phi(S(t)u_0) dt \\ = \inf_{\Psi \in \mathcal{T}_{\text{cyl}}(W')} \max_{u \in B \cap K} \{ \phi(u) + \langle F(u), \Psi'(u) \rangle_{W', W} \}. \end{aligned}$$

## Theorem

For the 3D NSE, with  $X = H_w$ ,  $W = D(A)$ , and  $K \supset \mathcal{A}_w$  compact,  $B \subset H$  bounded and positively invariant,  $\phi \in \mathcal{C}_b(B)$ ,

$$\begin{aligned}
 \sup_{u \in \mathcal{U}(B)} \limsup_{T \rightarrow \infty} \frac{1}{T} \int_0^T \phi(S(t)u_0) dt \\
 \leq \sup_{\mu \in \mathcal{P}_{fpss}(K)} \int_K \phi(u) d\mu(u) \\
 \leq \inf_{\Psi \in \mathcal{T}_{cyl}(W')} \max_{u \in B \cap K} \{ \phi(u) + \langle F(u), \Psi'(u) \rangle_{W', W} \}.
 \end{aligned}$$