

**Lecture Notes of Mathematics-I
for Integral Calculus, Improper
Integrals, Beta and Gamma
functions**

by

Prof Shiv Datt Kumar

Department of Mathematics

Motilal Nehru National Institute of Technology

Allahabad (UP), India

Pin - 211004

E-mail: sdt@mnnit.ac.in

Contents

1	Integral Calculus	v
1.1	Jacobian Matrix	v
1.1.1	Change of variables	vi
1.2	Volume of solid of revolution	viii
1.3	Double Integral	ix
1.3.1	Change of variables in double and triple integral	xi
1.4	Triple Integral	xii
2	Improper Integrals, Beta and Gamma Functions	xvii
2.1	Improper Integral	xvii
2.2	Beta Function	xx
2.2.1	Gamma Functions	xxi
2.3	Dirichlet Integral	xxiii
	Bibliography	xxv

Chapter 1

Integral Calculus

1.1 Jacobian Matrix

Let $F : \mathbb{R}^n \longrightarrow \mathbb{R}^m$ be a vector valued function given by $F(x_1, \dots, x_n) = (F_1(x_1, \dots, x_n), \dots, F_m(x_1, \dots, x_n))$. Then Jacobian matrix is the matrix of all first order partial derivatives defined as

$$J(F) = \begin{pmatrix} \frac{\partial F_1}{\partial x_1} & \cdots & \frac{\partial F_1}{\partial x_n} \\ \vdots & \vdots & \vdots \\ \frac{\partial F_m}{\partial x_1} & \cdots & \frac{\partial F_m}{\partial x_n} \end{pmatrix}$$

Jacobian matrix is important because if the function F is differentiable at point $p = (x_1, \dots, x_n)$, the Jacobian matrix defines a linear map $\mathbb{R}^n \longrightarrow \mathbb{R}^m$, which is the best linear approximation of the function F near point p . The Jacobian generalizes the gradient of a scalar valued function of several variables, which is generalization of derivative of a scalar valued function of single variable. Jacobian can be thought of as describing the amount of stretching, rotating or transforming and that transformation imposes locally.

Definition 1.1.1. *If F_1, F_2, \dots, F_n are functions of x_1, x_2, \dots, x_n , then the de-*

terminant

$$J = \begin{vmatrix} \frac{\partial F_1}{\partial x_1} & \cdots & \frac{\partial F_1}{\partial x_n} \\ \vdots & \vdots & \vdots \\ \frac{\partial F_n}{\partial x_1} & \cdots & \frac{\partial F_n}{\partial x_n} \end{vmatrix}$$

is called Jacobian of F_1, F_2, \dots, F_n with respect to x_1, x_2, \dots, x_n , In short it is written as $\frac{\partial(F_1, \dots, F_n)}{\partial(x_1, \dots, x_n)}$.

Remark: Determinant of square Jacobian matrix (called as Jacobian) gives important information about the behaviour of F near that point. If the Jacobian determinant at p is non-zero, then the continuously differentiable function F is invertible near a point $p \in \mathbb{R}^n$. This is the inverse function theorem. Further if $\text{Det}(J) > 0$, then F preserves orientation near p . If $\text{Det}(J) < 0$, then F reverses orientation. Absolute value of the Jacobian determinant gives the factor by which the function F expands or shrinks volumes near p .

Properties of Jacobian:

1. If f and g are functions of u and v and u, v are functions of x and y , then
$$\frac{\partial(f, g)}{\partial(x, y)} = \frac{\partial(f, g)}{\partial(u, v)} \frac{\partial(u, v)}{\partial(x, y)}$$
2. If J is the Jacobian of the system u, v with respect to x, y and J' is the Jacobian of x, y with respect to u and v , then $JJ' = 1$.

Example 1.1.2. If $x = r \cos \theta$ and $y = r \sin \theta$, then (i) $\frac{\partial(x, y)}{\partial(r, \theta)} = r$ and (ii)
$$\frac{\partial(r, \theta)}{\partial(x, y)} = \frac{1}{r}.$$

1.1.1 Change of variables

Suppose $z = f(x, y)$ and $x = \phi(u, v)$, $y = \psi(u, v)$. Then by chain rule

$$\frac{\partial f}{\partial u} = \frac{\partial f}{\partial x} \frac{\partial x}{\partial u} + \frac{\partial f}{\partial y} \frac{\partial y}{\partial u} \quad \text{and} \quad \frac{\partial f}{\partial v} = \frac{\partial f}{\partial x} \frac{\partial x}{\partial v} + \frac{\partial f}{\partial y} \frac{\partial y}{\partial v}.$$

Solving the system of equations by Crammer rule

$$\frac{\frac{\partial f}{\partial x}}{\frac{\partial f}{\partial u} \frac{\partial y}{\partial v} - \frac{\partial f}{\partial v} \frac{\partial y}{\partial u}} = \frac{\frac{\partial f}{\partial y}}{\frac{\partial f}{\partial v} \frac{\partial x}{\partial u} - \frac{\partial f}{\partial u} \frac{\partial x}{\partial v}} = \frac{1}{\frac{\partial x}{\partial u} \frac{\partial y}{\partial v} - \frac{\partial x}{\partial v} \frac{\partial y}{\partial u}}$$

$$\frac{\frac{\partial f}{\partial x}}{\frac{\partial(f, y)}{\partial(u, v)}} = \frac{\frac{\partial f}{\partial y}}{\frac{\partial(f, x)}{\partial(u, v)}} = \frac{1}{\frac{\partial(x, y)}{\partial(u, v)}}$$

$$\text{Determinant } J = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{vmatrix} = \frac{\partial(x, y)}{\partial(u, v)}$$

is called the Jacobian of variables of transformation. Therefore $\frac{\partial f}{\partial x} = \frac{1}{J} \frac{\partial(f, y)}{\partial(u, v)}$ and $\frac{\partial f}{\partial y} = -\frac{1}{J} \frac{\partial(f, x)}{\partial(u, v)}$.

In case of three variables, let $S = f(x, y, z)$ and $x = F(u, v, w)$, $y = G(u, v, w)$

& $z = H(u, v, w)$. Then

$$\frac{\partial f}{\partial u} = \frac{\partial f}{\partial x} \frac{\partial x}{\partial u} + \frac{\partial f}{\partial y} \frac{\partial y}{\partial u} + \frac{\partial f}{\partial z} \frac{\partial z}{\partial u}.$$

$$\frac{\partial f}{\partial v} = \frac{\partial f}{\partial x} \frac{\partial x}{\partial v} + \frac{\partial f}{\partial y} \frac{\partial y}{\partial v} + \frac{\partial f}{\partial z} \frac{\partial z}{\partial v}.$$

$$\frac{\partial f}{\partial w} = \frac{\partial f}{\partial x} \frac{\partial x}{\partial w} + \frac{\partial f}{\partial y} \frac{\partial y}{\partial w} + \frac{\partial f}{\partial z} \frac{\partial z}{\partial w}.$$

$$\frac{\partial f}{\partial x} = \frac{1}{J} \left[\frac{\partial(f, y, z)}{\partial(u, v, w)} \right], \quad \frac{\partial f}{\partial y} = -\frac{1}{J} \left[\frac{\partial(f, x, z)}{\partial(u, v, w)} \right] \text{ and } \frac{\partial f}{\partial z} = \frac{1}{J} \left[\frac{\partial(f, x, y)}{\partial(u, v, w)} \right], \text{ where}$$

$$J = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} & \frac{\partial x}{\partial w} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} & \frac{\partial y}{\partial w} \\ \frac{\partial z}{\partial u} & \frac{\partial z}{\partial v} & \frac{\partial z}{\partial w} \end{vmatrix}$$

Example 1.1.3. Let $z = f(x, y)$, $x = r \cos \theta$, $y = r \sin \theta$. Then

$$\left(\frac{\partial f}{\partial x} \right)^2 + \left(\frac{\partial f}{\partial y} \right)^2 = \left(\frac{\partial f}{\partial r} \right)^2 + \frac{1}{r^2} \left(\frac{\partial f}{\partial \theta} \right)^2.$$

$$\text{Here } J = \frac{\partial(x, y)}{\partial(r, \theta)} = r,$$

$$\frac{\partial f}{\partial x} = \frac{1}{J} \frac{\partial(f, y)}{\partial(r, \theta)} \dots \dots \dots (1) \text{ and}$$

$$\frac{\partial f}{\partial y} = -\frac{1}{J} \frac{\partial(f, x)}{\partial(r, \theta)} \dots \dots \dots (2).$$

$$\text{Also } \frac{\partial(f, y)}{\partial(r, \theta)} = r \cos \theta \frac{\partial f}{\partial r} - \sin \theta \frac{\partial f}{\partial \theta} \dots \dots \dots (3) \text{ and}$$

$$\frac{\partial(f, x)}{\partial(r, \theta)} = -r \sin \theta \frac{\partial f}{\partial r} - \cos \theta \frac{\partial f}{\partial \theta} \dots \dots \dots (4)$$

Squaring and adding (1) & (2) and using (3) & (4) we get the required.

Cartesian coordinates to cylindrical coordinates

Cylindrical coordinates (r, θ, z) are given by $x = r \cos \theta$, $y = r \sin \theta$, $z = z$ and

Jacobian of transformation is

$$J = J\left(\frac{(x, y, z)}{(r, \theta, z)}\right) = \begin{vmatrix} \frac{\partial x}{\partial r} & \frac{\partial x}{\partial \theta} & \frac{\partial x}{\partial z} \\ \frac{\partial y}{\partial r} & \frac{\partial y}{\partial \theta} & \frac{\partial y}{\partial z} \\ \frac{\partial z}{\partial r} & \frac{\partial z}{\partial \theta} & \frac{\partial z}{\partial z} \end{vmatrix}$$

$$= \begin{vmatrix} \cos \theta & -r \sin \theta & z \\ \sin \theta & r \cos \theta & 0 \\ 0 & 0 & 1 \end{vmatrix} = r$$

1.2 Volume of solid of revolution

Let AB be the portion of the curve $y = f(x)$, $f(x) > 0$, $x = a$, $x = b$. Consider the area bounded by the arc AB of the curve $y = f(x)$, x -axis and the lines $x = a$ and $x = b$. Volume of the solid generated by revolving this area about the x -axis is $V = \int_a^b \pi y^2 dx$. Similarly volume of the solid generated by revolving this area about the y -axis and lines $y = c$, $y = d$ is $V = \int_c^d \pi x^2 dy$.

Example 1.2.1. Find the volume of the solid generated by revolving the finite region bounded by the curve $y = x^2 + 1$, $y = 5$ about the line $x = 3$.

Solution: $V = \int_1^5 \pi(x_1^2 - x_2^2) dy = \int_1^5 \pi[(3 + \sqrt{y-1})^2 - (3 - \sqrt{y-1})^2] dy = \int_1^5 12\pi(\sqrt{y-1}) dy = 64\pi$.

Example 1.2.2. Consider the element $\delta x \delta y$ at $P(x, y)$ of plane area A . As this elementary area revolves about x -axis, we get a ring of volume $\pi[(y + \delta y)^2 -$

$$y^2] \delta x = 2\pi y \delta x \delta y. \quad \text{Total volume} = \int \int_A 2\pi y \, dx dy = \int \int_A 2\pi r \sin \theta \, r dr \, d\theta = \int \int_A 2\pi r^2 \sin \theta \, dr \, d\theta.$$

Example 1.2.3. Find the volume of the solid generated by revolving the finite region bounded by the curve $y = 3 - x^2$, $y = -1$ about the line $y = -1$.

Solution: $V = 2\pi$ [volume of solid generated about the $y = -1$]

$$= \int_0^2 \pi(1+y)^2 dx = \int_0^2 2\pi(1+y)^2 dx = \int_0^2 2\pi(1+3-x^2)^2 dx = \frac{512\pi}{15}.$$

Example 1.2.4. Find the volume of the solid generated by revolving the arc of cycloid $x = a(t - \sin t)$, $y = a(1 - \cos t)$, about x -axis.

Solution: $V = \int_0^{2\pi a} \pi y^2 dx = \int_0^{2\pi} \pi a^2 (1 - \cos t)^2 a(1 - \cos t) dt = 16\pi a^3 \int_0^\pi \sin^6 t \, dt = 5\pi^2 a^3.$

1.3 Double Integral

Notion of double integral is an extension of the notion of definite integral on the real line to the case of two dimensional space \mathbb{R}^2 . Let $f(x, y)$ be a continuous function in a simply connected, closed bounded region R in two variables x and y . Divide the region R into subregions (rectangles) by drawing lines $x = x_k$, $y = y_k$, $k = 1, 2, \dots, m$ parallel to coordinate axes. Let (x_i, y_i) be an arbitrary point inside the i th rectangle, whose area is ΔA_i . Let $S_n = \sum_{i=1}^n f(x_i, y_i) \Delta A_i$. When $n \rightarrow \infty$, the number of subregions increase indefinitely such that the largest of areas ΔA_i approaches zero. Then $\lim_{n \rightarrow \infty} S_n = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i, y_i) \Delta A_i$, if exists, is called the double integral of the function $f(x, y)$ over the region R and is denoted by $\int \int_R f(x, y) dx dy$.

Properties of double integrals

If $f(x, y)$ and $g(x, y)$ are integrable functions, then

1. $\int \int_R [f(x, y) \pm g(x, y)] dx dy = \int \int_R f(x, y) dx dy \pm \int \int_R g(x, y) dx dy.$
2. $\int \int_R k f(x, y) dx dy = k \int \int_R f(x, y) dx dy.$
3. $|\int \int_R k f(x, y) dx dy| \leq \int \int_R |f(x, y)| dx dy.$

4. **Mean Value Theorem:** $\int \int_R f(x, y) dx dy = f(u, v)A$, where A is the area of the region R and (u, v) is any arbitrary point in the region R . If $m \leq f(x, y) \leq M$ for all (x, y) in R , then $m A \leq \int \int_R f(x, y) dx dy \leq M A$.
5. If $0 < f(x, y) \leq g(x, y)$ for all $(x, y) \in R$, then $\int \int_R f(x, y) dx dy \leq \int \int_R g(x, y) dx dy$.
6. If $f(x, y) \geq 0$, then $\int \int_R f(x, y) dx dy \geq 0$.

Example 1.3.1. Evaluate $\int_0^1 \int_0^{\sqrt{1+x^2}} \frac{dy dx}{1+x^2+y^2}$.

Solution: $\int_0^1 \int_0^{\sqrt{1+x^2}} \frac{dy dx}{1+x^2+y^2} = \int_0^1 dx \int_0^{\sqrt{1+x^2}} \frac{dy}{1+x^2+y^2}$
 $= \int_0^1 dx \left[\frac{1}{\sqrt{1+x^2}} \tan^{-1} \frac{y}{1+x^2} \right]_0^{\sqrt{1+x^2}} = \int_0^1 dx \left[\frac{1}{\sqrt{1+x^2}} \frac{\pi}{4} \right]$
 $= \frac{\pi}{4} \int_0^1 \frac{1}{\sqrt{1+x^2}} dx = \frac{\pi}{4} [\log(x + \sqrt{x^2+1})]_0^1 = \frac{\pi}{4} [\log(\sqrt{2}+1)].$

Example 1.3.2. Evaluate $\int \int_A xy \, dx \, dy$, where A is the region bounded by the ordinate $x = 2a$, curve $x^2 = 4ay$.

Solution: Limit of integration for x are from 0 to $2a$ and for y from 0 to $\frac{x^2}{4a}$.

$$\int \int_A xy \, dx \, dy = \int_0^{2a} \int_0^{x^2/4a} xy \, dx \, dy = \int_0^{2a} x \, dx \int_0^{x^2/4a} y \, dy = a^4/3.$$

Example 1.3.3. Evaluate $\int \int_R e^{x^2} dx dy$, where $R : 2y \leq x \leq 2$ and $0 \leq y \leq 1$.

First integrate with respect to y and then with respect x .

$$\int_0^2 \left[\int_0^{\frac{x}{2}} e^{x^2} dy \right] dx = \int_0^2 [ye^{x^2}]_0^{\frac{x}{2}} dx = \frac{1}{2} \int_0^2 xe^{x^2} dx = \frac{1}{4}(e^4 - 1).$$

Applications of double integrals

1. If $f(x, y) = 1$, then $\int \int_R dx dy =$ Area of the region R .
2. If $z = f(x, y)$ is a surface, then $\int \int_R f(x, y) dx dy =$ Volume of the region beneath the surface $z = f(x, y)$ and above the $x - y$ plane.
3. If $f(x, y) = \rho(x, y)$ is the density function (mass per unit area), then
 - (i) $\int \int_R \rho(x, y) dx dy = M$ (Total mass of the region R).
 - (ii) Centre of gravity (\bar{x}, \bar{y}) is given by

$$\bar{x} = \frac{1}{M} \int \int_R x \rho(x, y) dx dy, \quad \bar{y} = \frac{1}{M} \int \int_R y \rho(x, y) dx dy.$$
 - (iii) Moment of inertia of the mass in R about x -axis, $I_x = \int \int_R y^2 \rho(x, y) dx dy$
 Moment of inertia of the mass in R about y -axis, $I_y = \int \int_R x^2 \rho(x, y) dx dy$

1.3.1 Change of variables in double and triple integral

Let R be the domain of integration $\int \int_R f(x, y) \, dx dy$ in $x - y$ plane. Let $x = \phi(u, v)$, $y = \psi(u, v)$ and R^* be the domain in the $u - v$ plane. Then

$$\int \int_R f(x, y) \, dx dy = \int \int_{R^*} F(u, v) |J| \, du dv, \text{ where}$$

$$\text{Determinant } J = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{vmatrix} = \frac{\partial(x, y)}{\partial(u, v)}.$$

If $x = \phi(u, v, w)$, $y = \psi(u, v, w)$, $z = \eta(u, v, w)$, then

$$\int \int \int_R f(x, y, z) \, dx dy dz = \int \int \int_{R^*} F(u, v, w) |J| \, du dv dw, \text{ where}$$

$$J = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} & \frac{\partial x}{\partial w} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} & \frac{\partial y}{\partial w} \\ \frac{\partial z}{\partial u} & \frac{\partial z}{\partial v} & \frac{\partial z}{\partial w} \end{vmatrix}$$

Example 1.3.4. Let R be rhombus with successive vertices at $(1, 0)$, $(2, 1)$, $(1, 2)$ and $(0, 1)$. Then find the value of the integral $\int \int_R (x - y)^2 \cos^2(x + y) dx dy$.

Solution: Equations of sides AB , BC , CD , and DA are given by $x - y = 1$, $x + y = 3$, $x - y = -1$ and $x + y = 1$ respectively. Put $x - y = u$, $y + x = v$. Then $-1 \leq u \leq 1$, $1 \leq v \leq 3$ and $x = (v - u)/2$, $y = (u + v)/2$. Jacobian of transformation is given by

$$J = \frac{\partial(x, y)}{\partial(u, v)} = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{vmatrix} = -1/2$$

$$\begin{aligned} \text{Then } I &= \int \int_R (x - y)^2 \cos^2(x + y) \, dx dy = \int_1^3 \int_{-1}^1 u^2 \cos^2 v |J| \, du dv \\ &= \frac{1}{2} \int_1^3 \int_{-1}^1 u^2 \cos^2 v \, du dv = \frac{1}{3} \int_1^3 \cos^2 v dv = \frac{1}{6} \int_1^3 (1 + \cos 2v) dv \\ &= \frac{1}{3} + \frac{\sin 6 - \sin 2}{12}. \end{aligned}$$

1.4 Triple Integral

Triple integral is an extension of the notion of double integral to three dimensional space \mathbb{R}^3 . Let $f(x, y, z)$ be a continuous function in a simply connected, closed bounded volume V . Divide the region V into small volume elements by drawing planes $x = x_k, y = y_k, z = z_k, k = 1, 2, \dots, n$ parallel to three coordinate planes. Let (x_i, y_i, z_i) be an arbitrary point inside the i th volume element $\delta V_i = \delta x_i \delta y_i \delta z_i$. Let $S_n = \sum_{i=1}^n f(x_i, y_i, z_i) \delta V_i$. When $n \rightarrow \infty$, the number of subregions increase indefinitely such that the largest of volumes δV_i approaches zero. Then $\lim_{n \rightarrow \infty} S_n = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i, y_i, z_i) \delta V_i$, if exists, is called the triple integral of the function $f(x, y, z)$ over the region V and is denoted by $\iiint_V f(x, y, z) dx dy dz$.

Let $f(x, y, z)$ be a continuous function over a regular solid V defined by $a < x < b, h_1(x) < y < h_2(x)$, and $g_1(x, y) < z < g_2(x, y)$. Then the triple integral is equal to the triple iterated integral given by

$$\iiint_V f(x, y, z) dx dy dz = \int_a^b \int_{h_1(x)}^{h_2(x)} \int_{g_1(x, y)}^{g_2(x, y)} f(x, y, z) dx dy dz.$$

Applications of triple integrals

1. If $f(x, y, z) = 1$, then $\iiint_R dx dy dz = \text{Volume of the region } R$.
2. If $f(x, y, z) = \rho(x, y, z)$ is the density of mass, then
 - (i) $\iiint_R \rho(x, y, z) dx dy dz = M$ (Total mass of the solid of region R).
 - (ii) Centre of gravity $(\bar{x}, \bar{y}, \bar{z})$ is given by

$$\bar{x} = \frac{1}{M} \iiint_R x \rho(x, y, z) dx dy dz, \bar{y} = \frac{1}{M} \iiint_R y \rho(x, y, z) dx dy dz,$$

$$\bar{z} = \frac{1}{M} \iiint_R z \rho(x, y, z) dx dy dz.$$
 - (iii) Moment of inertia of the mass in R about x -axis,

$$I_x = \iiint_R (y^2 + z^2) \rho(x, y, z) dx dy dz$$
 Moment of inertia of the solid of mass in R about y -axis,

$$I_y = \iiint_R (x^2 + z^2) \rho(x, y, z) dx dy dz$$
 and
 moment of inertia of the solid of mass in R about z -axis,

$$I_z = \iiint_R (x^2 + y^2) \rho(x, y, z) dx dy dz.$$

Example 1.4.1. Find the volume of the solid bounded by the plane $x = 0$, $y = 0$, $x + y + z = a$ and $z = 0$.

$$\text{Volume } V = \int_0^a \int_0^{a-x} \int_0^{a-x-y} dx \, dy \, dz = \int_0^a \int_0^{a-x} (a-x-y) dy \, dx = \frac{1}{2} \int_0^a (a-x)^2 dx = a^3/6.$$

Example 1.4.2. Find the volume generated by the revolution of cardioid

$r = a(1 - \cos \theta)$ about its axis.

$$\begin{aligned} \text{Volume } V &= \int_0^\pi \int_0^{a(1-\cos \theta)} 2\pi r^2 \, dr \sin \theta d\theta \\ &= \frac{2\pi a^3}{3} \int_0^\pi (1 - \cos \theta)^3 \sin \theta d\theta = \frac{\pi a^3}{6}. \end{aligned}$$

Example 1.4.3. Calculate the volume of the ellipsoid $\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1$.

Using the transformation of spherical coordinates $x = a \sin \theta \cos \phi$, $y = b \sin \theta \sin \phi$, $z = c \cos \theta$. Jacobian of transformation

$$J = \begin{vmatrix} \frac{\partial x}{\partial r} & \frac{\partial x}{\partial \theta} & \frac{\partial x}{\partial \phi} \\ \frac{\partial y}{\partial r} & \frac{\partial y}{\partial \theta} & \frac{\partial y}{\partial \phi} \\ \frac{\partial z}{\partial r} & \frac{\partial z}{\partial \theta} & \frac{\partial z}{\partial \phi} \end{vmatrix} = abc \, r^2 \sin \theta$$

$$dx dy dz = J dr \, d\theta \, d\phi = abc \, r^2 \sin \theta dr \, d\theta \, d\phi$$

$$\begin{aligned} V &= 8 \times \text{volume of sphere in positive octant} \\ &= 8 \int_0^1 \int_0^{\pi/2} \int_0^{\pi/2} abc \, r^2 \sin \theta dr d\theta \, d\phi \\ &= 8 \int_0^1 abc \, r^2 dr \int_0^{\pi/2} \sin \theta d\theta \int_0^{\pi/2} d\phi = \frac{4}{3} \pi abc. \end{aligned}$$

Example 1.4.4. Calculate the volume of the solid surrounded by the surface $(x/a)^{2/3} + (y/b)^{2/3} + (z/c)^{2/3} = 1$.

Using the transformation of coordinates $x = aX^3$, $y = bY^3$, $z = cZ^3$. Jacobian of transformation $J = \frac{\partial(x,y,z)}{\partial(X,Y,Z)} = 27abcX^2Y^2Z^2$. Required volume = $\int \int \int dx \, dy \, dz = 27abc \int \int \int X^2Y^2Z^2 dX \, dY \, dZ$

Now use the transformation $X = r \sin \theta \cos \phi$, $Y = r \sin \theta \sin \phi$, $Z = r \cos \theta$. Then Jacobian of transformation $J' = \frac{\partial(X,Y,Z)}{\partial(r,\theta,\phi)} = r^2 \sin \theta \, dr d\theta d\phi$. Thus $dX dY dZ = r^2 \sin \theta dr d\theta d\phi$ and $dx dy dz = 27abcX^2Y^2Z^2 r^2 \sin \theta \, dr d\theta d\phi$.

$V = 8 \times \text{volume in positive octant} =$

$$\begin{aligned} & 8 \int_0^1 \int_0^{\pi/2} \int_0^{\pi/2} 27abc (r \sin \theta \cos \phi)^2 (r \sin \theta \sin \phi)^2 (r \cos \theta)^2 r^2 \sin \theta dr d\theta d\phi. \\ & = 216abc \int_0^1 r^8 dr \int_0^{\pi/2} \sin^5 \theta \cos^2 \theta d\theta \int_0^{\pi/2} \sin^2 \phi \cos^2 \phi d\phi = \frac{4\pi abc}{35}. \end{aligned}$$

Example 1.4.5. Find total mass of the solid bounded by the surfaces $x^2 + y^2 = 16$, $z = 2$ and $z = 4$ if its density function is $\rho(x, y, z) = x^2 + y^2$.

Convert the integral $\int_{x=-4}^4 \int_{y=-\sqrt{16-x^2}}^{\sqrt{16-x^2}} \int_{z=2}^{z=4} \rho(x, y, z) dx dy dz$ in cylindrical coordinates $x = r \cos \theta$, $y = r \sin \theta$, $z = z$, then Jacobian $J = r$ and

$$\int_{r=0}^4 \int_{\theta=0}^{2\pi} \int_{z=2}^4 r^2 r dr d\theta dz = \int_{r=0}^4 r^3 dr \int_{\theta=0}^{2\pi} d\theta \int_{z=2}^4 dz = \left\{ \frac{r^4}{4} \right\}_0^4 \times 2\pi \times 2 = 256\pi.$$

Example 1.4.6. Show that in the first octant, paraboloid $z = 36 - 4x^2 - 9y^2$ has the volume 27π cubic units.

Solution: Projection of the given paraboloid in the $x y$ plane is the first quadrant of the ellipse $4x^2 + 9y^2 = 36$. Then region R is given by $0 \leq z \leq 36 - (4x^2 + 9y^2)$,

$0 \leq y \leq \frac{1}{3}\sqrt{36 - 4x^2}$, $0 \leq x \leq 3$. Therefore volume

$$\begin{aligned} V &= \int_0^3 \left[\int_0^{\frac{1}{3}\sqrt{36-4x^2}} (36 - 4x^2 - 9y^2) dy \right] dx \\ &= \int_0^3 \left[4(9 - x^2)y - 3y^3 \right]_0^{\frac{1}{3}\sqrt{36-4x^2}} dx = \frac{16}{9} \int_0^3 (9 - x^2)^{3/2} dx. \end{aligned}$$

Now put $x = 3 \sin \theta$. Then $V = 144 \int_0^{\pi/2} \cos^4 \theta d\theta = 144 \left(\frac{3}{4} \cdot \frac{1}{2} \cdot \frac{\pi}{2} \right) = 27\pi$ cubic unit.

Example 1.4.7. Find the volume of the solid which is contained in between the cone $z^2 = 2(x^2 + y^2)$ and the hyperboloid $z^2 = x^2 + y^2 + a^2$.

Solution: Put $x = r \cos \theta$, $y = r \sin \theta$. Then $dx dy = r dr d\theta$ and intersection of $z^2 = 2(x^2 + y^2)$ and $z^2 = x^2 + y^2 + a^2$ is $2(x^2 + y^2) = x^2 + y^2 + a^2$ i.e. $x^2 + y^2 = a^2$.

Therefore

$$\begin{aligned} V &= \int_0^{2\pi} \int_0^a \int_{\sqrt{2(x^2+y^2)}}^{\sqrt{x^2+y^2+a^2}} dz dy dx = \int_0^{2\pi} \int_0^a \left(\sqrt{x^2+y^2+a^2} - \sqrt{2(x^2+y^2)} \right) dy dx = \\ &= \int_0^{2\pi} \int_0^a (\sqrt{r^2+a^2} - \sqrt{2}r) r dr d\theta = \frac{4\pi a^3(\sqrt{2}-1)}{3}. \end{aligned}$$

Example 1.4.8. Find mass of a solid lying between spheres of radius 3 and 4 in the region $y \geq 0$, $z \geq 0$. Density at any point (x, y, z) is given by $\rho(x, y, z) = \sqrt{x^2 + y^2 + z^2}$.

Solution: Convert the integral into spherical coordinates $x = r \sin \theta \cos \phi$,

$y = r \sin \theta \sin \phi$, $z = r \cos \theta$, Region occupied by the solid is described by $3 \leq r \leq 4$, $0 \leq \theta \leq \pi$, $0 \leq \phi \leq \frac{\pi}{2}$. Also

$$\rho(r, \theta, \phi) = \rho(r \sin \theta \cos \phi, r \sin \theta \sin \phi, r \cos \theta) \\ = \sqrt{(r \sin \theta \cos \phi)^2 + (r \sin \theta \sin \phi)^2 + (r \cos \theta)^2} = r.$$

Jacobian of transformation $J = r^2 \sin \phi$.

$$\text{Hence mass} = \int_0^\pi \int_0^{\pi/2} \int_3^4 r^2 \sin \phi \, dr \, d\phi \, d\theta = \int_0^\pi d\theta \int_0^{\pi/2} \sin \phi \, d\phi \int_3^4 r^3 \, dr = \frac{175\pi}{4}$$

Exercise 1.4.1. 1. Evaluate $\iiint \sin(x+y+z) \, dx \, dy \, dz$ over portion cut off by the plane $x+y+z = \pi$.

2. By a suitable change of variables calculate the integral $\iint_R \log \frac{x-y}{x+y} \, dx \, dy$, where R is the triangular region bounded the vertices $(1, 0)$, $(4, -3)$, $(4, 1)$.
Ans: $\frac{1}{4}(49 \log 7 - \frac{75}{2} \log 5 - 27 \log 3 + 6)$.

3. Evaluate $\iint_R (x+y)^2 \, dx \, dy$, where R is the region bounded by the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$, using transformation of variables.
Ans: $\frac{\pi ab}{4}(a^2 + b^2)$.

4. Show that the area of the surface of paraboloid $\frac{x^2}{a} + \frac{y^2}{b} = 2z$ inside the cylinder $\frac{x^2}{a^2} + \frac{y^2}{b^2} = k$ is $\frac{2}{3}\pi\{(1+k)^{3/2} - 1\}ab$.

5. Find the area of the part of the cylinder $x^2 + y^2 = a^2$ which is cut by the cylinder $x^2 + z^2 = a^2$.

6. Let $f(x, y) = \frac{xy(x^2 - y^2)}{x^2 + y^2}$, $(x, y) \neq (0, 0)$, $f(0, 0) = 0$. Then show that $f_{xy}(0, 0) \neq f_{yx}(0, 0)$.

Chapter 2

Improper Integrals, Beta and Gamma Functions

2.1 Improper Integral

For the existence of Riemann integral (definite integral) $\int_a^b f(x)dx$, we require that the limit of integration a and b are finite and function $f(x)$ is bounded. In case

(i) limit of integration a or b or both become infinite (improper integral of first kind),

(ii) integrand $f(x)$ has singular points (discontinuity) i.e. $f(x)$ becomes infinite at some points in the interval $a \leq x \leq b$ (improper integral of second kind),

then the integral $\int_a^b f(x)dx$ is called improper integral. Note that improper integral are evaluated by limiting process.

Example 2.1.1. $\int_{-1}^{\infty} \frac{-1}{x^2} dx$, $\int_{-\infty}^{\infty} \frac{1}{1+x^2} dx$, $\int_0^1 \frac{dx}{x(1-x)}$ are improper integrals.

Definition 2.1.2. If a is the only point of infinite discontinuity of $f(x)$, then

$$\int_a^b f(x)dx = \lim_{\epsilon \rightarrow 0^+} \int_{a+\epsilon}^b f(x)dx \quad (\text{if finite limit exists})$$

If b is the only point of infinite discontinuity, then

$$\int_a^b f(x)dx = \lim_{\epsilon \rightarrow 0+} \int_a^{b-\epsilon} f(x)dx, \quad 0 < \epsilon < b-a$$

If end points a and b are the only points of infinite discontinuity,

$$\int_a^b f(x)dx = \lim_{\epsilon \rightarrow 0+} \lim_{\mu \rightarrow 0+} \int_{a-\epsilon}^{b-\mu} f(x)dx,$$

Improper integral is convergent if limit exists and is finite otherwise it is divergent.

If an interior point c , $a < c < b$, is the only point of infinite discontinuity, then we write

$$\int_a^b f(x)dx = \int_a^c f(x)dx + \int_c^b f(x)dx.$$

Improper integral is convergent if both the limits exist and are finite otherwise it is divergent.

Example 2.1.3. Examine the convergence of $\int_a^b \frac{dx}{(x-a)^n}$

Solution: It is a proper integral if $n \leq 0$ and improper for other values of n .

For $n \neq 1$,

$$\begin{aligned} \int_a^b \frac{dx}{(x-a)^n} &= \lim_{p \rightarrow 0+} \int_{a+p}^b \frac{dx}{(x-a)^n}, \quad 0 < p < b-a \\ &= \lim_{p \rightarrow 0+} \frac{1}{(1-n)} \left[\frac{1}{(b-a)^{n-1}} - \frac{1}{p^{n-1}} \right] = \begin{cases} \frac{1}{(1-n)(b-a)^{n-1}}, & \text{if } n < 1 \\ \infty & \text{if } n > 1, \end{cases} \end{aligned}$$

For $n = 1$

$$\begin{aligned} \int_a^b \frac{dx}{(x-a)^n} &= \lim_{p \rightarrow 0+} \int_{a+p}^b \frac{dx}{(x-a)} \\ &= \lim_{p \rightarrow 0+} \log(b-a) - \log p = \infty \end{aligned}$$

Thus integral converges only if $n < 1$.

Comparison Test I

If $f(x)$ and $g(x)$ are two functions such that $f(x) \leq g(x)$, for all $x \in [a, b]$, then

1. $\int_a^b f(x) dx$ converges if $\int_a^b g(x) dx$ converges.
2. $\int_a^b g(x) dx$ diverges if $\int_a^b f(x) dx$ diverges.

Comparison Test II

If $f(x)$ and $g(x)$ are two functions such that $\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} = l$ (nonzero & finite), then $\int_a^\infty f(x) dx$ and $\int_a^\infty g(x) dx$ converge or diverge together.

Example 2.1.4. Examine the convergence of (i) $\int_0^1 \frac{dx}{\sqrt{1-x^3}}$

(ii) $\int_0^{\pi/2} \frac{\sin x dx}{x^n}$ (iii) $\int_1^\infty \frac{dx}{x^3(e^{-x} + 1)}$

Solution: (i) Let $f(x) = \frac{1}{\sqrt{1-x^3}} = \frac{1}{\sqrt{(1-x)}\sqrt{1+x+x^2}}$.

Note that $\frac{1}{\sqrt{1+x+x^2}}$ is a bounded function and let M be its upper bound.

Then $f(x) \leq \frac{M}{(1-x)^{1/2}}$ and $\int_0^1 \frac{1}{(1-x)^{1/2}} dx$ is convergent. Therefore by comparison test $\int_0^1 \frac{dx}{\sqrt{1-x^3}}$ is convergent.

(ii) For $n \leq 1$, it is a proper integral. For $n > 1$, it is an integral and 0 is the point of infinite discontinuity. Now $\frac{\sin x}{x^n} = \frac{\sin x}{x} \frac{1}{x^{n-1}}$. Function $\frac{\sin x}{x}$ is bounded and $\frac{\sin x}{x} \leq 1$. Thus $\frac{\sin x}{x^n} \leq \frac{1}{x^{n-1}}$ and $\int_0^{\pi/2} \frac{dx}{x^{n-1}}$ converges only if $n-1 < 1$ or $n < 2$ and diverges for $n \geq 2$.

(iii) Let $f(x) = \frac{1}{x^3(e^{-x} + 1)}$ and $g(x) = \frac{1}{x^3}$. Also

$$\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} = \lim_{x \rightarrow \infty} \frac{1}{x^3(e^{-x} + 1)} \frac{x^3}{1} = \lim_{x \rightarrow \infty} \frac{1}{e^{-x} + 1} = 1.$$

Also $\int_1^\infty \frac{1}{x^3} dx$ converges. Therefore $\int_1^\infty \frac{dx}{x^3(e^{-x} + 1)}$ converges.

Example 2.1.5. Examine the convergence of (i) $\int_{-\pi/2}^{\pi/2} \tan x dx$ (ii) $\int_0^{\pi/2} \frac{\cos^n x}{x^m} dx$, if $m < 1$.

Solution:

$$(i) \int_{-\pi/2}^{\pi/2} \tan x dx = \lim_{\epsilon \rightarrow 0} \int_{-\pi/2+\epsilon}^c \tan x dx + \lim_{\eta \rightarrow 0} \int_c^{\pi/2-\eta} \tan x dx$$

$$= \lim_{\epsilon \rightarrow 0} \ln[\cos(-\pi/2 + \epsilon)] - \ln[\cos c] - \lim_{\eta \rightarrow 0} \ln[\cos(\pi/2 - \eta)] - \ln[\cos c].$$

Limits do not exist. Hence improper integral diverges.

(ii) Note that $\frac{\cos^n x}{x^m} < \frac{1}{x^m}$ for $0 < x < \pi/2$. Also $x = 0$ is the point of infinite discontinuity and $\int_0^{\pi/2} \frac{1}{x^m} dx$ is convergent if $m < 1$ by comparison test.

Absolute Convergence of Improper Integrals

If the function $f(x)$ changes sign within the interval of integration, we consider absolute convergence. The improper integral $\int_a^b f(x) dx$ is called absolutely convergent if $\int_a^b |f(x)| dx$ converges. Since $f(x) \leq |f(x)|$, $\forall x$, absolutely convergent improper integral is convergent.

Example 2.1.6. Examine the convergence of improper integral $\int_{-\infty}^{\infty} \frac{\sin x}{1+x^2} dx$.

Solution: Note that $|I| = \left| \int_{-\infty}^{\infty} \frac{\sin x}{1+x^2} dx \right| \leq \int_{-\infty}^{\infty} \left| \frac{\sin x}{1+x^2} \right| dx$

$$= \lim_{a \rightarrow -\infty} \int_a^c \left| \frac{\sin x}{1+x^2} \right| dx + \lim_{b \rightarrow \infty} \int_c^b \left| \frac{\sin x}{1+x^2} \right| dx = I_1 + I_2.$$

$$I_1 = \lim_{a \rightarrow -\infty} \int_a^c \left| \frac{\sin x}{1+x^2} \right| dx \leq \lim_{a \rightarrow -\infty} \int_a^c \left| \frac{1}{1+x^2} \right| dx = \tan^{-1}c + \frac{\pi}{2}.$$

$$I_2 = \lim_{b \rightarrow \infty} \int_c^b \left| \frac{\sin x}{1+x^2} \right| dx \leq \lim_{b \rightarrow \infty} \int_c^b \left| \frac{1}{1+x^2} \right| dx = \frac{\pi}{2} - \tan^{-1}c.$$

Thus $|I| \leq |I_1| + |I_2| \leq \pi$ and hence convergent.

2.2 Beta Function

The improper integral defined by

$$\beta(m, n) = \int_0^1 x^{m-1}(1-x)^{n-1} dx \text{ for } m > 0, n > 0 \text{ is called as beta function.}$$

Note that

$$\int_0^1 x^{m-1}(1-x)^{n-1} dx \text{ converges if } m > 0, n > 0.$$

Properties:

$$1. \beta(m, n) = \beta(n, m)$$

Put $x = 1 - y$. Then

$$\beta(m, n) = - \int_1^0 (1-y)^{m-1} y^{n-1} dy = \int_0^1 y^{n-1} (1-y)^{m-1} dy = \beta(n, m).$$

$$2. \beta(m, n) = \int_0^{\pi/2} \sin^{2m-1} \theta \cos^{2n-1} \theta d\theta, n > 0.$$

$$\text{Put } x = \sin^2 \theta, dx = 2 \sin \theta \cos \theta d\theta \text{ in } \beta(m, n) = \int_0^1 x^{m-1} (1-x)^{n-1} dx.$$

$$3. \beta(m, n) = \beta(m+1, n) + \beta(m, n+1)$$

2.2.1 Gamma Functions

The improper integral of the form $\Gamma(n) = \int_0^\infty e^{-x} x^{n-1} dx$, $n > 0$ is called gamma function.

Properties:

$$1. \Gamma(1) = \int_0^\infty e^{-x} dx = 1.$$

$$2. \text{Reduction formula } \Gamma(n+1) = n\Gamma(n)$$

$$\begin{aligned} \text{Proof: } \Gamma(n+1) &= \int_0^\infty e^{-x} x^n dx = [-x^n e^{-x}]_0^\infty + n \int_0^\infty e^{-x} x^{n-1} dx \\ &= n\Gamma(n) = n!. \end{aligned}$$

$$3. \Gamma(1/2) = \sqrt{\pi}.$$

$$\text{Proof: } \Gamma(1/2) = \int_0^\infty e^{-x} x^{1/2} dx = 2 \int_0^\infty e^{-y^2} dy \text{ (by putting } x = y^2 \text{)}.$$

$$\begin{aligned} (\Gamma(1/2))^2 &= 4 \int_0^\infty \int_0^\infty e^{-(x^2+y^2)} dx dy \\ &= 4 \int_0^{\pi/2} \int_0^\infty e^{-r^2} r dr d\theta = \frac{4\pi}{2} \int_0^\infty e^{-r^2} r dr = \pi \end{aligned}$$

$$4. \text{Relation between } \beta \text{ and } \Gamma \text{ functions}$$

$$\beta(m, n) = \frac{\Gamma(m)\Gamma(n)}{\Gamma(m+n)}$$

$$\text{Proof: } \Gamma(m) = \int_0^\infty e^{-t} t^{m-1} dt = 2 \int_0^\infty e^{-x^2} x^{2m-1} dx$$

$$\Gamma(n) = 2 \int_0^\infty e^{-y^2} y^{2n-1} dy$$

$$\begin{aligned} \Gamma(m)\Gamma(n) &= 4 \int_0^\infty e^{-x^2} x^{2m-1} dx \int_0^\infty e^{-y^2} y^{2n-1} dy \\ &= 4 \int_0^\infty \int_0^\infty e^{-(x^2+y^2)} x^{2m-1} y^{2n-1} dx dy \\ &= 4 \int_0^{\pi/2} \int_0^\infty e^{-r^2} r^{2(m+n)-1} \cos^{2m-1} \theta \sin^{2n-1} \theta dr d\theta \\ &= 2 \int_0^\infty e^{-r^2} r^{2(m+n)-1} dr \times 2 \int_0^{\pi/2} \cos^{2m-1} \theta \sin^{2n-1} \theta d\theta \\ &= \Gamma(m+n)\beta(m, n). \end{aligned}$$

$$5. \int_0^{\pi/2} \cos^m \theta \sin^n \theta d\theta$$

$$= \frac{1}{2} \beta\left(\frac{m+1}{2}, \frac{n+1}{2}\right) = \frac{\Gamma(\frac{m+1}{2})\Gamma(\frac{n+1}{2})}{2\Gamma(\frac{m+n+2}{2})}$$

Example 2.2.1. If $\int_0^\infty \frac{x^{m-1}}{1-x} dx = \frac{\pi}{\sin m\pi}$, $0 < m < 1$, then $\Gamma(m)\Gamma(1-m) = \frac{\pi}{\sin m\pi}$.

Put $\frac{x}{1+x} = y$. Then $x = \frac{y}{1-y}$ and $dx = \frac{1}{(1-y)^2} dy$. Thus

$$\begin{aligned} \int_0^\infty \frac{x^{m-1}}{1-x} dx &= \int_0^1 y^{m-1}(1-y)^{m-1} dy = \beta(m, 1-m) \\ &= \frac{\Gamma(m)\Gamma(1-m)}{\Gamma(m+1-m)} = \Gamma(m)\Gamma(1-m). \text{ Hence } \int_0^\infty \frac{x^{m-1}}{1-x} dx = \frac{\pi}{\sin m\pi}. \end{aligned}$$

Example 2.2.2. (Legendre Duplication formula)

$$\sqrt{\pi} \Gamma(2m) = 2^{2m-1} \Gamma(m) \Gamma(m + \frac{1}{2}).$$

$$\text{Since } \frac{\Gamma(m)\Gamma(n)}{\Gamma(m+n)} = \beta(m, n) = 2 \int_0^{\pi/2} \sin^{2m-1}\theta \cos^{2n-1}\theta d\theta \dots \dots \dots (1)$$

Putting $n = m$, we have

$$\begin{aligned} \frac{\Gamma(m)^2}{\Gamma(2m)} &= \beta(m, m) = 2 \int_0^{\pi/2} \sin^{2m-1}\theta \cos^{2m-1}\theta d\theta. \\ &= \frac{1}{2^{2m-2}} \int_0^{\pi/2} \sin^{2m-1} 2\theta d\theta. \text{ Put } 2\theta = \phi. \text{ Then} \\ &= \frac{1}{2^{2m-1}} \int_0^{\pi/2} \sin^{2m-1} \phi d\phi \dots \dots \dots (2). \end{aligned}$$

In (1) take $n = 1/2$, we get

$$\frac{\Gamma(m)\Gamma(1/2)}{\Gamma(m+1/2)} = 2 \int_0^{\pi/2} \sin^{2m-1}\theta d\theta \dots \dots \dots (3).$$

From (2) and (3), we get,

$$\begin{aligned} \frac{\Gamma(m)^2}{\Gamma(2m)} &= \frac{1}{2^{2m-1}} \frac{\Gamma(m)\Gamma(1/2)}{\Gamma(m+1/2)}. \text{ Since } \Gamma(1/2) = \sqrt{\pi}, \text{ we have} \\ \sqrt{\pi} \Gamma(2m) &= 2^{2m-1} \Gamma(m) \Gamma(m + \frac{1}{2}). \end{aligned}$$

Example 2.2.3. $\beta(m + \frac{1}{2}, m + \frac{1}{2}) = \frac{\pi}{m 2^{4m-1} \beta(m, m)}$.

Solution: By definition $\beta(m, n) = \int_0^1 x^{m-1}(1-x)^{n-1} dx$, $m > 0$, $n > 0$. By

substituting $x = \sin^2 \theta$, we have $\beta(m, n) = 2 \int_0^{\pi/2} \sin^{2m-1}\theta \cos^{2n-1}\theta d\theta$.

$$\text{First show that } \beta(m, m) = \frac{1}{2^{2m-1}} \beta(m, 1/2) \dots \dots \dots (1)$$

$$\text{Then } \Gamma(m+1/2)\Gamma(m) = \frac{\sqrt{\pi}}{2^{2m-1}} \Gamma(2m) \dots \dots \dots (2)$$

$$\text{Since } \frac{\Gamma(m)\Gamma(n)}{\Gamma(m+n)} = \beta(m, n) = 2 \int_0^{\pi/2} \sin^{2m-1}\theta \cos^{2n-1}\theta d\theta, \text{ we have}$$

$$\beta(m+1/2, m+1/2) = \frac{\Gamma(m+1/2)\Gamma(m+1/2)}{\Gamma(2m+1)}. \text{ Using (2) we have}$$

$$\beta(m + \frac{1}{2}, m + \frac{1}{2}) = \frac{\pi}{m 2^{4m-1} \beta(m, m)}.$$

Example 2.2.4. (i) Let $\phi(\alpha) = \int_{a(\alpha)}^{b(\alpha)} f(x, \alpha) dx$. Then by Leibniz formula

$$\frac{d\phi}{d\alpha} = \int_{a(\alpha)}^{b(\alpha)} \frac{\partial f(x, \alpha)}{\partial \alpha} dx + f(b, \alpha) \frac{db}{d\alpha} - f(a, \alpha) \frac{da}{d\alpha} \text{ show that } \frac{d\phi}{da} = \frac{\pi}{2(a+1)}.$$

$$(ii) \phi(a) = \int_0^\infty \frac{\tan^{-1}(ax)}{x(1+x^2)} dx.$$

Solution: Let $\phi(a) = \int_0^\infty \frac{\tan^{-1}(ax)}{x(1+x^2)} dx$. Then by Leibniz formula

$$\begin{aligned} \frac{d\phi}{da} &= \int_0^\infty \frac{\partial}{\partial a} \left[\frac{\tan^{-1}(ax)}{x(1+x^2)} \right] dx = \int_0^\infty \frac{dx}{(1+x^2)(1+a^2x^2)} \\ &= \frac{1}{a^2-1} \int_0^\infty \left[\frac{a^2}{a^2x^2+1} - \frac{1}{1+x^2} \right] dx = \frac{1}{a^2-1} [\{a \tan^{-1}(ax)\}_0^\infty - \{\tan^{-1} x\}_0^\infty] \\ &= \frac{\pi}{2} \left[\frac{a-1}{a^2-1} \right], \quad a > 0, a \neq 1 \\ &= \frac{\pi}{2(a+1)}. \quad \text{Integrating with respect to } a, \text{ we have} \end{aligned}$$

$$\phi(a) = \frac{\pi}{2} \ln(a+1) + c. \quad \text{Since } \phi(0) = 0, c = 0.$$

$$\text{Therefore } \phi(a) = \int_0^\infty \frac{\tan^{-1}(ax)}{x(1+x^2)} dx = \frac{\pi}{2} \ln(a+1).$$

2.3 Dirichlet Integral

Theorem 2.3.1. $\int \int_D x^{l-1} y^{m-1} dx dy = \frac{\Gamma(l)\Gamma(m)}{\Gamma(l+m+1)} h^{l+m}$, where D is domain $x \geq 0, y \geq 0, x+y \leq h$.

Proof. Put $x = Xh, y = Yh, dx dy = h^2 dX dY$. Then

$$\begin{aligned} \int \int_D x^{l-1} y^{m-1} dx dy &= \int \int_D (Xh)^{l-1} (Yh)^{m-1} dX dY \\ &= h^{l+m} \int_0^1 \int_0^{1-X} X^{l-1} Y^{m-1} dX dY \\ &= h^{l+m} \int_0^1 X^{l-1} dX \int_0^{1-X} Y^{m-1} dY \\ &= h^{l+m} \int_0^1 X^{l-1} dX \left[\frac{Y^m}{m} \right]_0^{1-X} \\ &= \frac{h^{l+m}}{m} \int_0^1 X^{l-1} (1-X)^m dX \\ &= \frac{h^{l+m}}{m} \beta(l, m+1) \\ &= \frac{h^{l+m}}{m} \frac{\Gamma(l)\Gamma(m+1)}{\Gamma(l+m+1)} \\ &= h^{l+m} \frac{\Gamma(l)\Gamma(m)}{\Gamma(l+m+1)} \quad \square \end{aligned}$$

Theorem 2.3.2. $\int \int \int_V x^{l-1} y^{m-1} z^{n-1} dx dy dz = \frac{\Gamma(l)\Gamma(m)\Gamma(n)}{\Gamma(l+m+n+1)}$, where V is region $x \geq 0, y \geq 0, z \geq 0, x+y+z \leq 1$.

Proof. Put $x+z \leq 1-x=h$. then $z \leq h-y$.

$$\begin{aligned} \int \int \int_V x^{l-1} y^{m-1} z^{n-1} dx dy dz &= \int_0^1 x^{l-1} dx \int_0^{1-x} y^{m-1} dy \int_0^{1-x-y} z^{n-1} dz \\ &= \int_0^1 x^{l-1} dx \left[\int_0^h \int_0^{h-y} y^{m-1} dy z^{n-1} dz \right] \end{aligned}$$

$$\begin{aligned}
&= \int_0^1 x^{l-1} dx [h^{m+n} \frac{\Gamma(m)\Gamma(n)}{\Gamma(m+n+1)}] \\
&= \frac{\Gamma(m)\Gamma(n)}{\Gamma(m+n+1)} \int_0^1 x^{l-1} (1-x)^{m+n} dx \\
&= \frac{\Gamma(m)\Gamma(n)}{\Gamma(m+n+1)} \beta(l, m+n+1) \\
&= \frac{\Gamma(m)\Gamma(n)}{\Gamma(m+n+1)} \frac{\Gamma(l)\Gamma(m+n+1)}{\Gamma(l+m+n+1)} \\
&= \frac{\Gamma(l)\Gamma(m)\Gamma(n)}{\Gamma(l+m+n+1)} \quad \square
\end{aligned}$$

Corollary 2.3.3. $\int \int \int_V x^{l-1} y^{m-1} z^{n-1} dx dy dz = \frac{\Gamma(l)\Gamma(m)\Gamma(n)}{\Gamma(l+m+n+1)} h^{l+m+n}$, where V is region $x \geq 0, y \geq 0, z \geq 0, x+y+z \leq h$.

Exercise 2.3.4. 1. Discuss the convergence of (i) $\int_1^2 \frac{\sqrt{x}}{\ln x} dx$ (ii) $\int_0^{\pi/2} \frac{\sin x}{x\sqrt{x}} dx$.

2. Examine the convergence of improper integrals (i) $\int_2^\infty \frac{dx}{\ln x}$
(ii) $\int_1^\infty e^{-x^2} dx$ (iii) $\int_1^\infty \frac{dx}{x^2(e^{-x}+1)}$ (iv) $\int_1^\infty \frac{dx}{x^p}$.

3. Show that for $m, n > 0$,

$$\beta(m, n) = \int_0^\infty \frac{x^{m-1}}{(1+x)^{m+n}} dx. \text{ Hint: Put } x = \frac{t}{1-t}$$

4. Show that $\int_0^\infty \frac{x^c}{c^x} dx = \frac{\Gamma(c+1)}{(\log c)^{c+1}}$.

5. Show that $\Gamma(2n) = \frac{2^{2n-1}}{\sqrt{\pi}} \Gamma(n + \frac{1}{2}) \Gamma(n)$ and hence $\Gamma(1/4)\Gamma(3/4) = \pi\sqrt{2}$.

6. Show that $\int_0^{\pi/2} \sqrt{\tan \theta} d\theta = \frac{1}{2} \Gamma(\frac{1}{4}) \Gamma(\frac{3}{4}) = \pi\sqrt{2}$. (Hint: $\int_0^{\pi/2} \sqrt{\tan \theta} dx =$
 $\int_0^{\pi/2} \sin^{1/2} \theta \cos^{-1/2} \theta d\theta = \frac{1}{2} \frac{\Gamma(1/4)\Gamma(3/4)}{\Gamma(1)} = \frac{\pi\sqrt{2}}{2} = \frac{\pi}{\sqrt{2}}$.

7. Evaluate the improper integral $\int_0^\infty \sqrt{x} e^{-x^2} dx$.

Bibliography

- [1] R K Jain and S R K Iyengar, Advance Engineering Mathematics, Narosa Publishing Corporation.
- [2] Ervin Kreszig, Advance Engineering Mathematics.
- [3] S C Malik, Savita Arora, Mathematical Analysis, New Age International Publishers, Fifth edition, 2017.

Index

Change of variables, vi, xi

Gamma function, xxi

Jacobian, v, vi

Legendre Duplication formula, xxii

Volume of solid of revolution, viii