

# Lecture : Beta and Gamma Functions

**Dr.Sanjay Mishra**  
[drsanjaymishra1@gmail.com](mailto:drsanjaymishra1@gmail.com)

Department of Mathematics  
Lovely Professional University  
Punjab, India

© 2014 All Rights Reserved

No part of this presentation may be reproduced or transmitted in any form or any means, electronic or mechanical, including photocopying or by any information storage retrieval system, without permission from author.

January 31, 2015

# Outline

## 1 Introduction

# Outline

- 1 Introduction
- 2 Gamma Functions

# Outline

- 1 Introduction
- 2 Gamma Functions
- 3 Transformation of Gamma Function

# Outline

- 1 Introduction
- 2 Gamma Functions
- 3 Transformation of Gamma Function
- 4 Example

# Outline

- 1 Introduction
- 2 Gamma Functions
- 3 Transformation of Gamma Function
- 4 Example
- 5 Exercise

# Introduction I

$$\Gamma(n) = \int_0^{\infty} x^{n-1} e^{-x} dx$$

$$(x_1, x_2, \dots, x_n) \times (y_1, y_2, \dots, y_n)$$

# Gamma Functions I

## Definition

If  $n > 0$ , the gamma function  $\Gamma$  is defined by the improper integral

$$\Gamma(n) = \int_0^{\infty} x^{n-1} e^{-x} dx$$

This integral converge for all  $n > 0$ .

It is sometimes called the *second Eulerian integral*.



# Gamma Functions II

The most useful factorial property of gamma function are as follows.

## Theorem

If  $n > 0$ , then  $\Gamma(n + 1) = n\Gamma(n)$

## Proof:

By definition of gamma function

$$\Gamma(n) = \int_0^{\infty} x^{n-1} e^{-x} dx \quad (1)$$

If  $0 < a < b$ , then we can integrate by parts

$$\begin{aligned} \int_a^b x^n e^{-x} dx &= [x(-e^{-x})]_a^b - \int_a^b nx^{n-1}(-1)e^{-x} dx \\ &= -b^n e^{-b} + a^n e^{-a} + n \int_a^b x^{n-1} e^{-x} dx \end{aligned}$$

Take the limit of this equation as  $a \rightarrow 0$  and  $b \rightarrow \infty$  to get

$$= \Gamma(n + 1) = n \int_0^{\infty} x^{n-1} e^{-x} dx = n\Gamma(n)$$

# Gamma Functions III

By repeating this result, we get

$$\begin{aligned}
 \Gamma(n+1)n\Gamma(n) &= n(n-1)\Gamma(n-1) \\
 &= n(n-1)(n-2)\Gamma(n-3) \\
 &= \dots = n(n-1)(n-2)\dots(2)(1)\Gamma(1) \\
 &= n!\Gamma(1)
 \end{aligned}$$

But

$$\Gamma(1) = \int_0^{\infty} e^{-x} dx = 1 \quad (2)$$

So

$$\Gamma(n+1) = n!$$

# Gamma Functions IV

From this result we can find the gamma function of negative integer.

## Theorem

*If  $n$  is zero of a negative integer then  $\Gamma(n) = \infty$ .*

*Proof:*

# Transformation of Gamma Function I

We will prove some important results of gamma function using the improper integral formulation.

## Theorem

If  $n > 0$ , then  $\frac{\Gamma(n)}{z^n} = \int_0^\infty e^{-zx} x^{n-1} dx$

## Proof:

We know that

$$\Gamma(n) = \int_0^\infty x^{n-1} e^{-x} dx \quad (3)$$

On putting  $x = az$  in the above equation (3), we get

$$\begin{aligned} \Gamma(n) &= \int_0^\infty e^{-az} (az)^{n-1} a dz \\ &= a^n \int_0^\infty e^{-az} z^{n-1} a dz \\ &= a^n \int_0^\infty e^{-ax} x^{n-1} dx \end{aligned} \quad (4)$$

Replacing  $a$  by  $z$  in above equation, we get

$$\frac{\Gamma(n)}{z^n} = \int_0^\infty e^{-zx} x^{n-1} dx$$

## Theorem

$$\Gamma\left(\frac{1}{2}\right) = \sqrt{\pi}$$

**Proof:** By definition of gamma function

$$\Gamma(n) = \int_0^{\infty} x^{n-1} e^{-x} dx \quad (5)$$

Put  $x^n = y$  and  $nx^{n-1}dx = dy$ , we get

$$\begin{aligned} \Gamma(n) &= \int_0^{\infty} y^{n-1/n} e^{-y^{1/n}} \frac{1}{nx^{n-1}} dy \\ &= \int_0^{\infty} y^{n-1/n} e^{-y^{1/n}} \frac{1}{ny^{n-1/n}} dy \\ \Gamma(n) &= \frac{1}{n} \int_0^{\infty} e^{-y^{1/n}} dy \end{aligned}$$

Now put  $n = \frac{1}{2}$  in above equation, we get

$$\Gamma\left(\frac{1}{2}\right) = \frac{1}{2} \int_0^{\infty} e^{-y^{1/2}} dy = 2\left(\frac{1}{2}\sqrt{\pi}\right) = \sqrt{\pi}$$

## Example

Evaluate  $\int_0^\infty \frac{x^c}{c^x} dx$  where  $c > 1$ .

### *Solution:*

Putting  $c^x = e^t$  and  $dx = \frac{dt}{\log c}$ , we obtain

$$\begin{aligned} I &= \int_0^\infty \left( \frac{t}{\log c} \right)^c e^{-t} \frac{dt}{\log c} \\ &= \frac{1}{(\log c)^{c+1}} \int_0^\infty e^{-t} t^c dt \\ &= \frac{1}{(\log c)^{c+1}} \Gamma(c+1) \end{aligned}$$

## Example

Evaluate  $\Gamma\left(-\frac{7}{2}\right)$ .

**Solution:** As we now that in  $n$  is negative but not integer, then

$$\Gamma(n) = \frac{1}{n} \Gamma(n+1) \quad (6)$$

Therefore

$$\Gamma\left(-\frac{7}{2}\right) = \frac{1}{\left(-\frac{7}{2}\right)} \Gamma\left(-\frac{7}{2} + 1\right) = \left(-\frac{2}{7}\right) \Gamma\left(-\frac{5}{2}\right)$$

Again repeating the above formula. we get

$$\begin{aligned} \Gamma\left(-\frac{7}{2}\right) &= \left(-\frac{2}{7}\right) \left(-\frac{2}{5}\right) \left(-\frac{2}{3}\right) \left(-\frac{2}{1}\right) \Gamma\left(\frac{1}{2}\right) \\ &= \frac{16}{105} \Gamma\left(\frac{1}{2}\right) \\ \Gamma\left(-\frac{7}{2}\right) &= \frac{16}{105} \sqrt{\pi} \end{aligned}$$

# Exercise I

- Evaluate  $\int_0^\infty x^{2n-1} e^{-ax^2} dx$ .
- Prove that  $\int_0^\infty e^{-y^{1/m}} dy = m\Gamma(m)$ .
- Evaluate  $\Gamma\left(-\frac{3}{2}\right)$ .



You can learn more on click from my Home Page ...



<http://mathematicsresources.wordpress.com/>

Feel free to mail me for any type of suggestion for improvement of this lecture notes: [drsanjaymishra1@gmail.com](mailto:drsanjaymishra1@gmail.com)

Thank you!

