

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/279310361>

# On the $(p,q)$ -Gamma and the $(p,q)$ -Beta functions

**Article** · June 2015

Source: arXiv

---

CITATIONS

13

---

READS

701

**1 author:**



Patrick Njionou S.

University of Yaounde I

**19** PUBLICATIONS **134** CITATIONS

SEE PROFILE

**Some of the authors of this publication are also working on these related projects:**



Fractional calculus [View project](#)

# On the $(p, q)$ -Gamma and the $(p, q)$ -Beta functions

P. Njionou Sadjang

**Abstract.** We introduce new generalizations of the Gamma and the Beta functions. Their properties are investigated and known results are obtained as particular cases.

**Mathematics Subject Classification (2010).** 33B15; 26A51; 26A48.

**Keywords.** Gamma function, Beta function,

## 1. Introduction

The  $q$ -deformed algebras [15, 16] and their generalizations ( $(p, q)$ -deformed algebras) [4, 5] attract much attention these last years. The main reason is that these topics stand for a meeting point of today's fast developing areas in mathematics and physics like the theory of quantum orthogonal polynomials and special functions, quantum groups, conformal field theories and statistics. From these works, many generalizations of special functions arise. There is a considerable list of references.

In this work, we give a new generalization of the Gamma and the Beta functions, namely, the  $(p, q)$ -Gamma and the  $(p, q)$ -Beta functions. Their main properties are stated and proved and connections with the previous work are done. It is to be noted that in [13], the authors provided another generalization of the Gamma function.

The Euler gamma function  $\Gamma(x)$  first happens in 1729 in a correspondence between Euler and Goldbach and is defined for  $x > 0$  by (see [1, 17])

$$\Gamma(x) = \int_0^\infty t^{x-1} e^{-t} dt. \quad (1)$$

Euler gave an equivalent representation of the Gamma function [1, 2, ?, 17]

$$\Gamma(x) = \lim_{n \rightarrow \infty} \frac{n! n^x}{x(x+1) \cdots (x+n)}. \quad (2)$$

An immediate consequence of these representations is

$$\Gamma(x+1) = x\Gamma(x).$$

Also, for any nonnegative integer  $n$ ,

$$\Gamma(n+1) = n!$$

follows from the above argument.

Closely connected with the Gamma function is the Beta function [1, 17] defined as

$$B(x, y) = \int_0^1 t^{x-1} (1-t)^{y-1} dt, \quad \Re x > 0, \Re y > 0. \quad (3)$$

The Beta function is related to the Gamma function in the following way

$$B(x, y) = \frac{\Gamma(x)\Gamma(y)}{\Gamma(x+y)}. \quad (4)$$

Jackson (see [10, 11, 12, ?]) defined the  $q$ -analogue of the gamma function as

$$\Gamma_q(x) = \frac{(q; q)_\infty}{(q^x; q)_\infty} (1-q)^{1-x}, \quad 0 < q < 1, \quad (5)$$

and

$$\Gamma_q(x) = \frac{(q^{-1}; q^{-1})_\infty}{(q^{-x}; q^{-1})_\infty} (q-1)^{1-x} q^{\binom{x}{2}}, \quad q > 1. \quad (6)$$

An equivalent definition of (5) is given in [9] as

$$\Gamma_q(x) = \int_0^\infty t^{x-1} E_q^{-qt} d_q t \quad (7)$$

where the  $q$ -integral is defined by (see [6, 9, ?])

$$\int_0^a f(s) d_q s = a(1-q) \sum_{n=0}^\infty q^n f(aq^n), \quad a > 0, \quad (8)$$

$$\int_a^0 f(s) d_q s = -a(1-q) \sum_{n=0}^\infty q^n f(aq^n), \quad a < 0, \quad (9)$$

$$\int_a^\infty f(s) d_q s = a(q^{-1}-1) \sum_{n=0}^\infty q^{-n} f(aq^{-n-1}), \quad a > 0, \quad (10)$$

$$\int_{-\infty}^a f(s) d_q s = -a(q^{-1}-1) \sum_{n=0}^\infty q^{-n} f(aq^{-n-1}), \quad a < 0, \quad (11)$$

and can be extended to the whole real line by using relations (8)-(11) and the following rules

$$\begin{aligned}
\int_a^b f(s) d_q s &= \int_a^0 f(s) d_q s + \int_0^b f(s) d_q s \quad \forall a, b \in \mathbb{R} \\
\int_a^\infty f(s) d_q s &= \int_a^b f(s) d_q s + \int_b^\infty f(s) d_q s \quad \forall a, b \in \mathbb{R}, \quad a < 0, \quad b > 0 \\
\int_{-\infty}^b f(s) d_q s &= \int_{-\infty}^a f(s) d_q s + \int_a^b f(s) d_q s \quad \forall a, b \in \mathbb{R}, \quad a < 0, \quad b > 0 \\
\int_{-\infty}^\infty f(s) d_q s &= \int_{-\infty}^a f(s) d_q s + \int_a^b f(s) d_q s + \int_b^\infty f(s) d_q s \quad \forall a, b \in \mathbb{R}.
\end{aligned}$$

## 2. Definitions and Miscellaneous Relations

Let us introduce the following notation (see [7],[8],[14])

$$[n]_{p,q} = \frac{p^n - q^n}{p - q}, \quad (12)$$

for any positive integer.

The twin-basic number is a natural generalization of the  $q$ -number, that is

$$\lim_{p \rightarrow 1} [n]_{p,q} = [n]_q. \quad (13)$$

The  $(p, q)$ -factorial is defined by ([8, 14])

$$[n]_{p,q}! = \prod_{k=1}^n [k]_{p,q}!, \quad n \geq 1, \quad [0]_{p,q}! = 1. \quad (14)$$

Let us introduce also the so-called  $(p, q)$ -binomial coefficient

$$\begin{bmatrix} n \\ k \end{bmatrix}_{p,q} = \frac{[n]_{p,q}!}{[k]_{p,q}! [n-k]_{p,q}!}, \quad 0 \leq k \leq n. \quad (15)$$

Note that as  $p \rightarrow 1$ , the  $(p, q)$ -binomial coefficients reduce to the  $q$ -binomial coefficients.

It is clear by definition that

$$\begin{bmatrix} n \\ k \end{bmatrix}_{p,q} = \begin{bmatrix} n \\ n-k \end{bmatrix}_{p,q}. \quad (16)$$

Let us introduce also the so-called the  $(p, q)$ -powers [14]

$$(x \ominus a)_{p,q}^n = (x - a)(px - aq) \cdots (xp^{n-1} - aq^{n-1}), \quad (17)$$

$$(x \oplus a)_{p,q}^n = (x + a)(px + aq) \cdots (xp^{n-1} + aq^{n-1}). \quad (18)$$

These definition are extended to

$$(a \ominus b)_{p,q}^\infty = \prod_{k=0}^{\infty} (ap^k - q^k b) \quad (19)$$

$$(a \oplus b)_{p,q}^\infty = \prod_{k=0}^{\infty} (ap^k + q^k b) \quad (20)$$

where the convergence is required.

**Proposition 2.1.** *The following identities are easily verified*

$$(a \ominus b)_{p,q}^n = \frac{(a \ominus b)_{p,q}^\infty}{(ap^n \ominus bq^n)_{p,q}^\infty} \quad (21)$$

$$(a \ominus b)_{p,q}^{n+k} = (a \ominus b)_{p,q}^n (ap^n \ominus bq^n)_{p,q}^k \quad (22)$$

$$(ap^n \ominus bq^n)_{p,q}^k = \frac{(a \ominus b)_{p,q}^k (ap^k \ominus bq^k)_{p,q}^n}{(a \ominus b)_{p,q}^n} \quad (23)$$

$$(ap^k \ominus bq^k)_{p,q}^{n-k} = \frac{(a \ominus b)_{p,q}^n}{(a \ominus b)_{p,q}^k} \quad (24)$$

$$(ap^{2k} \ominus bq^{2k})_{p,q}^{n-k} = \frac{(a \ominus b)_{p,q}^n (ap^n \ominus bq^n)_{p,q}^k}{(a \ominus b)_{p,q}^{2k}} \quad (25)$$

$$(a^2 \ominus b^2)_{p,q}^n = (a \ominus b)_{p,q}^n (a \oplus b)_{p,q}^n \quad (26)$$

$$(a \ominus b)_{p,q}^{2n} = (a \ominus b)_{p^2,q^2}^n (ap \ominus bq)_{p^2,q^2}^n \quad (27)$$

$$(a \ominus b)_{p,q}^{3n} = (a \ominus b)_{p^3,q^3}^n (ap \ominus bq)_{p^3,q^3}^n (ap^2 \ominus bq^2)_{p^3,q^3}^n \quad (28)$$

$$(a \ominus b)_{p,q}^{\ell n} = \prod_{j=0}^{\ell-1} (ap^j \ominus bq^j)_{p^\ell,q^\ell}^n. \quad (29)$$

### 3. The $(p, q)$ -Gamma function

**Definition 3.1.** *Let  $x$  be a complex number, we define the  $(p, q)$ -Gamma function as*

$$\Gamma_{p,q}(x) = \frac{(p \ominus q)_{p,q}^\infty}{(p^x \ominus q^x)_{p,q}^\infty} (p - q)^{1-x}, \quad 0 < q < p. \quad (30)$$

**Remark 3.2.** *Note that in (30), if we put  $p = 1$ , then  $\Gamma_{p,q}$  reduces to  $\Gamma_q$ .*

**Proposition 3.3.** *The  $(p, q)$ -Gamma function fulfils the following fundamental relation*

$$\Gamma_{p,q}(x+1) = [x]_{p,q} \Gamma_{p,q}(x). \quad (31)$$

*Proof.* By definition, we have

$$\Gamma_{p,q}(x+1) = \frac{(p \ominus q)_{p,q}^\infty}{(p^{x+1} \ominus q^{x+1})_{p,q}^\infty} (p - q)^{-x}$$

□

**Remark 3.4.** If  $n$  is a nonnegative integer, it follows from (31) that

$$\Gamma_{p,q}(n+1) = [n]_{p,q}!.$$

It can be also easily seen from the definition that

$$\Gamma_{p,q}(n+1) = \frac{(p \ominus q)_{p,q}^n}{(p-q)^n}.$$

**Proposition 3.5 ((p, q)-Legendre's multiplication formula).** The following multiplication formula applies

$$\Gamma_{p,q}(2x)\Gamma_{p^2,q^2}\left(\frac{1}{2}\right) = (p+q)^{2x-1}\Gamma_{p^2,q^2}(x)\Gamma_{p^2,q^2}\left(x+\frac{1}{2}\right). \quad (32)$$

*Proof.* From the definition, we have

$$\begin{aligned} \Gamma_{p^2,q^2}(x) &= \frac{(p^2 \ominus q^2)_{p^2,q^2}^\infty}{(p^{2x} \ominus q^{2x})_{p^2,q^2}^\infty} (p^2 - q^2)^{1-x} \\ \Gamma_{p^2,q^2}\left(x + \frac{1}{2}\right) &= \frac{(p^2 \ominus q^2)_{p^2,q^2}^\infty}{(p^{2x+1} \ominus q^{2x+1})_{p^2,q^2}^\infty} (p^2 - q^2)^{\frac{1}{2}-x} \\ \Gamma_{p^2,q^2}\left(\frac{1}{2}\right) &= \frac{(p^2 \ominus q^2)_{p^2,q^2}^\infty}{(p \ominus q)_{p^2,q^2}^\infty} (p^2 - q^2)^{\frac{1}{2}}. \end{aligned}$$

Hence,

$$\begin{aligned} \frac{\Gamma_{p^2,q^2}(x)\Gamma_{p^2,q^2}\left(x+\frac{1}{2}\right)}{\Gamma_{p^2,q^2}\left(\frac{1}{2}\right)} &= \frac{(p^2 \ominus q^2)_{p^2,q^2}^\infty (p \ominus q)_{p^2,q^2}^\infty}{(p^{2x} \ominus q^{2x})_{p^2,q^2}^\infty (p^{2x+1} \ominus q^{2x+1})_{p^2,q^2}^\infty} (p^2 - q^2)^{1-2x} \\ &= \frac{(p \ominus q)_{p,q}^\infty}{(p^{2x} \ominus q^{2x})_{p,q}^\infty} (p-q)^{1-2x} (p+q)^{1-2x} \\ &= (p+q)^{1-2x} \Gamma_{p,q}(2x). \end{aligned}$$

This proves the proposition.  $\square$

The (p, q)-Legendre's multiplication formula is generalized as follows.

**Proposition 3.6 ((p, q)-Gauss' multiplication formula).** The following multiplication formula applies

$$\Gamma_{p,q}(nx) \prod_{k=1}^{n-1} \Gamma_{p^n,q^n}\left(\frac{k}{n}\right) = ([n]_{p,q})^{nx-1} \prod_{k=0}^{n-1} \Gamma_{p^n,q^n}\left(x + \frac{k}{n}\right). \quad (33)$$

*Proof.* As for the previous proposition, we start by using the definition as follows

$$\begin{aligned} \Gamma_{p^n,q^n}\left(\frac{k}{n}\right) &= \frac{(p^n \ominus q^n)_{p^n,q^n}^\infty}{(p^k \ominus q^k)_{p^n,q^n}^\infty} (p^n - q^n)^{1-\frac{k}{n}}, \\ \Gamma_{p^n,q^n}\left(x + \frac{k}{n}\right) &= \frac{(p^n \ominus q^n)_{p^n,q^n}^\infty}{(p^{nx+k} \ominus q^{nx+k})_{p^n,q^n}^\infty} (p^n - q^n)^{1-\frac{k}{n}-x}. \end{aligned}$$

Hence, we have

$$\begin{aligned}
 \prod_{k=1}^{n-1} \Gamma_{p^n, q^n} \left( \frac{k}{n} \right) &= \frac{[(p^n \ominus q^n)_{p^n, q^n}^\infty]^{n-1}}{\prod_{k=1}^{n-1} (p^k \ominus q^k)_{p^n, q^n}^\infty} (p^n - q^n)^{\sum_{k=1}^{n-1} (1 - \frac{k}{n})} \\
 &= \frac{[(p^n \ominus q^n)_{p^n, q^n}^\infty]^n}{\prod_{k=0}^{n-1} (p.p^k \ominus q.q^k)_{p^n, q^n}^\infty} (p^n - q^n)^{\frac{n-1}{2}} \\
 &= \frac{[(p^n \ominus q^n)_{p^n, q^n}^\infty]^n}{(p \ominus q)_{p, q}^\infty} (p^n - q^n)^{\frac{n-1}{2}}
 \end{aligned}$$

and

$$\begin{aligned}
 \prod_{k=0}^{n-1} \Gamma_{p^n, q^n} \left( x + \frac{k}{n} \right) &= \frac{[(p^n \ominus q^n)_{p^n, q^n}^\infty]^n}{\prod_{k=0}^{n-1} (p^{nx+k} \ominus q^{nx+k})_{p^n, q^n}^\infty} (p^n - q^n)^{\sum_{k=0}^{n-1} (1 - \frac{k}{n} - x)} \\
 &= \frac{[(p^n \ominus q^n)_{p^n, q^n}^\infty]^n}{(p^{nx} \ominus q^{nx})_{p, q}^\infty} (p^n - q^n)^{(\frac{n-1}{2} + 1 - nx)}.
 \end{aligned}$$

It follows that

$$\begin{aligned}
 \frac{\prod_{k=0}^{n-1} \Gamma_{p^n, q^n} \left( x + \frac{k}{n} \right)}{\prod_{k=1}^{n-1} \Gamma_{p^n, q^n} \left( \frac{k}{n} \right)} &= \frac{(p \ominus q)_{p, q}^\infty}{(p^{nx} \ominus q^{nx})_{p, q}^\infty} (p^n - q^n)^{1-nx} \\
 &= \frac{(p \ominus q)_{p, q}^\infty}{(p^{nx} \ominus q^{nx})_{p, q}^\infty} (p - q)^{1-nx} \left( \frac{p^n - q^n}{p - q} \right)^{1-nx} \\
 &= ([n]_{p, q})^{1-nx} \Gamma_{p, q}(nx).
 \end{aligned}$$

The proposition is then proved.  $\square$

#### 4. The $(p, q)$ -Beta function

**Definition 4.1.** Following (4), we define the  $(p, q)$ -Beta functions as

$$B_{p, q}(x, y) = \frac{\Gamma_{p, q}(x) \Gamma_{p, q}(y)}{\Gamma_{p, q}(x + y)}. \quad (34)$$

**Proposition 4.2.** *The (p, q)-Beta function fulfils the following properties*

$$B_{p,q}(x, y + 1) = \frac{[y]_{p,q}}{[x + y]_{p,q}} B_{p,q}(x, y) \quad (35)$$

$$B_{p,q}(x + 1, y) = \frac{[x]_{p,q}}{[x + y]_{p,q}} B_{p,q}(x, y) \quad (36)$$

$$B_{p,q}(x + 1, y) = \frac{[x]_{p,q}}{[y]_{p,q}} B_{p,q}(x, y + 1) \quad (37)$$

$$B_{p,q}(x + n, y) = \frac{(p^x \ominus q^x)_{p,q}^n}{(p^{x+y} \ominus q^{x+y})_{p,q}^n} B_{p,q}(x, y). \quad (38)$$

## References

- [1] G.E. Andrews, R. Askey, R. Roy: *Special Functions*, Encyclopedia Math. Appl. **71**, Cambridge University Press, Cambridge, 2000.
- [2] T.M. Apostol, *Introduction to Analytic Number Theory*, Springer, 1976.
- [3] J.D. Bukweli-Kyemba, M.N. Hounkonnou: *Quantum deformed algebras : coherent states and special functions*, <http://arXiv:1301.0116v1>, 2013.
- [4] I. M. Burban, A. U. Klimyk: (P, Q)-Differentiation, (P, Q) integration, and (P, Q)-hypergeometric functions related to quantum groups, *Integral Transform and Special Functions*, 2:1, 15-36, (1994)
- [5] R. Chakrabarti, R. Jagannathan, *A (p, q)-oscillator realization of two-parameter quantum algebras*, J. Phys. A: Math. Gen. **24**, L711, 1991
- [6] M. Foupouagnigni: *Laguerre-Hahn orthogonal polynomials with respect to the Hahn operator, fourth-order difference equation for the rth associated and the Laguerre-Freud equations for the recurrence coefficients*, Ph.D. Thesis, Universit Nationale du Bnin, Bnin, 1998.
- [7] R. Jagannathan, K. Srinivasa Rao, *Two-parameter quantum algebras, twin-basic numbers, and associated generalized hypergeometric series*, in: *Proceedings of the International Conference on Number Theory and Mathematical Physics*, Srinivasa Ramanujan Centre, Kumbakonam, India, 20-21 December 2005
- [8] R. Jagannathan, R. Sridhar, *(p, q)-Rogers-Szegő Polynomials and the (p, q)-Oscillator*, K. Alladi et al. (eds.), *The Legacy of Alladi Ramakrishnan in the Mathematical Sciences*, 2010.
- [9] V. Kac, P. Cheung: *Quantum calculus*, Springer, (2001).
- [10] T. Kim, *On a q-analogue of the p-adic log gamma functions and related integrals*, J. Number Theory **76** (1999) 320-329.
- [11] T. Kim, *A note on the q-multiple zeta functions*, *Advan. Stud. Contemp. Math.* **8** (2004) 111-113.
- [12] T. Kim, S.H. Rim, *A note on the q-integral and q-series*, *Advanced Stud. Contemp. Math.* **2** (2000) 37-45.
- [13] V. Krasniqi, F. Merovci, *Some completely monotonic properties for the (p, q)-Gamma function*, <http://arxiv.org/abs/1407.4231v1>, 2014.
- [14] P. Njionou Sadjang, *On the fundamental theorem of (p, q)-calculus and some (p, q)-Taylor formulas*, Submitted to *Integral Transform*, 2014.



- [15] A. Odziejewicz, *Quantum algebraa and  $q$ -special functions related to coherent states maps of the disc*, Commun. Math. Phys., **192**, 183-215, 1998
- [16] C. Quesne, K. A. Penson, V. M. Tkachuk, *Maths-type  $q$ -deformed coherent states for  $q > 1$* , Phys. Lett. A 313 **29**, 2003
- [17] N. M. Temme: *Special Functions, an Introduction to Classical Functions of Mathematical Physics*, John Wiley & Sons, New York, 1996.

P. Njionou Sadjang  
Faculty of Industrial Engineering  
University of Douala  
Cameroon  
e-mail: pnjionou@yahoo.fr