Math 126	Name:
Summer 2018	
Final	
8/9/2018	
Time Limit: 110 Minutes	

This exam contains 6 questions. Total of points is 85.

For this exam, you are allowed a handwritten cheat sheet consisting of one side of an $8 ext{1/2}$ by 11 piece of paper.

The last sheet of paper is intentionally blank so that you may use it to finish your solutions if clearly marked where the first part of your solution stops.

Grade Table (for instructor use only)

Question	Points	Score
1	10	
2	15	
3	15	
4	15	
5	15	
6	15	
Total:	85	

1. (10 points) Suppose that u(x,t) solves

$$u_t - ku_{xx} = f(x,t)$$
 $0 < x < \infty, t > 0$
 $u(x,0) = \phi(x),$ $\infty < x < \infty$

where $\phi(x)$ is even and f(-x,t)=f(x,t). Show that u(-x,t)=u(x,t).

2. (15 points) Solve

$$\begin{split} \Delta u &= 0 & 0 \leq x, y \leq \pi \\ u(x,0) &= u(x,\pi) = 0 & 0 \leq x \leq \pi \\ u(0,y) &= 0, u(\pi,y) = \sin y & 0 \leq y \leq \pi \end{split}$$

3. (15 points) (a) (8 points) Use the forward difference scheme discussed in class to solve

with $\Delta t = \Delta x = 1$ until time t = 4. Express your answer in the form of a grid like we did in class and be clear to describe your scheme, as well as your notation. Also, for consistency, take u(0,0) = 1.

- (b) (2 points) Is this scheme stable?
- (c) (5 points) If the scheme is stable, then how do you know? If the scheme is not stable, then what could you do to fix it?

4. (15 points) Use Fourier transform techniques to solve

$$u_t + cu_x = 0 -\infty < x < \infty, t > 0$$

$$u(x, 0) = \phi(x).$$

(Hint: Instead of computing the inverse Fourier transform directly, you may want to simply recognize your function as the Fourier transform of something using the function and rule tables.)

5. (15 points) Find the Green's function for D=[0,1] in one dimension. (Hint: First find a function v(x) such that $v_{xx}=\delta(x)$, then adjust to fit the other conditions.

6. (15 points) Use the method of reflection to solve the wave equation on the half-plane $\{y > 0\}$ with Neumann boundary condition, i.e.

$$u_{tt} = c^{2}(u_{xx} + u_{yy}) \qquad -\infty < x < \infty, y > 0, t > 0$$

$$u_{y}(x, 0, t) = 0 \qquad -\infty < x < \infty, t > 0$$

$$u(x, y, 0) = 0, u_{t}(x, y, 0) = \psi(x, y) \qquad -\infty < x < \infty, y > 0.$$

In your answer, you may use any statements about even and/or odd solutions without justification provided they are true. Also, feel free to leave your solution in terms of the source function, provided you say what that is.