SWAOC Anaesthesia for patients undergoing Spinal Surgery

NHS Southwest Ambulatory Orthopaedic Centre

Pre-meds

- Paracetamol 1g, Ibuprofen 1.6g MR
- Pre-warming (as required). Ensure A2RB/ACEi are stopped day before & day of surgery.
- Fasting times minimised: 6hrs food; 2hrs coffee/tea with milk; water until sending time

Intra-op Care

- General Anaesthesia (ideally TIVA)
- BIS must be used if using TIVA and muscle relaxant
- Routine dual antiemetics intraop (**Dexamethasone 9.9mg**, **Ondansetron 4mg**)
- Tranexamic Acid 1g iv at induction (10mg/kg if eGFR <60)
- Fentanyl 250 mcg during closure
- Aim **1000mls only** IV fluids unless bleeding. Active **warming**. Mechanical thromboprophylaxis.
- Normotension: aim MAP within 20% normal, **Vasopressor** use rather than excess fluids

Antibiotics

Teicoplanin – dose on actual body weight			
40 – 66kg	67 – 100kg	101 – 133kg	134 – 166kg
400mg	**600mg**	**800mg**	**1g**

Gentamicin – dose on actual body weight				
40 – 53kg	54 – 80kg	81 – 106kg	107 – 133kg	134 – 166kg
160mg	**240mg**	**320mg**	**400mg**	**480mg**

Teicoplanin & Gentamicin

Single IV doses

Avoid mixing of agents

Surgical LIA

- Before Knife to Skin
- Surgical infiltration into skin and deep paravertebral muscles: 0.25% levobupivacaine with 1:200,000 adrenaline 40mls (reduced if <50 kg)

Postop

- Build up energy drink – Ensure Compact 125ml to be given to all patients in PACU.
- Apart from Haemocue check (>100g/L) or BM checks in diabetics, no routine bloods needed
- PACU: Fentanyl 25mcg IV prn or
- PACU: 2-4hrly prn: Oxynorm (age <60: 5-10mg >60yrs: 2.5-5mg)

TTAs

Paracetamol	1g	qds	for 10 days
Ibuprofen	400mg	qds	for 6 days
Codeine	30-60mg	qds	for 7 days OR
Tramadol	50-100mg	qds	for 7 days (if codeine intolerant)
Macragols	1 sachet	bd	for 10 days
Oxycodone IR	5mg	prn	5 doses, maximum 4 hourly