

However, we disagree with Stewart-Brown and colleagues' view that wellbeing interventions have been proven to reduce the prevalence of mental disorders.² To show the effectiveness of this theory, use of a wellbeing intervention would need to report a reduction in cases of people with mental disorders in the population offered the intervention. However, the evidence that Stewart-Brown and colleagues² invoke simply does not show this reduction. For example, they cited a systematic review of mindfulness interventions that assessed clinical populations of people with mental or physical disorders, and none of the studies reported a reduction in the prevalence of mental disorders.⁴

Population-based strategies to prevent mental health disorders that include wellbeing interventions are not the answer. New proposals must be judged against a background whereby clinical commissioning groups in England spend a small and decreasing proportion of their budgets on mental health,⁵ despite between only a quarter and a third of people with common mental disorders receiving any treatment.⁶

Primary prevention of mental disorders is more likely to depend on equitable distribution of wealth, reforming the criminal justice system, and ensuring equal opportunities in education and employment than on wellbeing interventions.

NM is the editor-in-chief of the annual report of the Chief Medical Officer, 2013.⁵ MHo, MHe, and SW declare no competing interests.

Matthew Hotopf, *Nisha Mehta, Max Henderson, Simon Wessely
nisha.mehta@nhs.net

Institute of Psychiatry, Psychology, and Neuroscience, King's College London, London, UK (MHo, MHe, SW); and National Health Service Tayside, Ninewells Hospital, Dundee DD1 9SY, Scotland (NM)

- 1 Mehta N, Croudace T, Davies SC. Public mental health: evidenced-based priorities. *Lancet* 2015; **385**: 1472–75.
- 2 Stewart-Brown S, Middleton J, Ashton J. Responses to the Chief Medical Officer's report 2013. *Lancet* 2015; **385**: 2576.
- 3 Rose G. The strategy of preventive medicine. Oxford: Oxford University Press, 1992.

- 4 de Vibe M, Bjørndal A, Tipton E, Hammerstrøm K, Kowalski K. Mindfulness based stress reduction (mbsr) for improving health, quality of life, and social functioning in adults. *Campbell Syst Rev* 2012; **3**. DOI:10.4073/csr.2012.3.
- 5 Thornicroft G, Docherty M. Mind the gaps—treatment, funding, access and service provision. Annual report of the Chief Medical Officer 2013. Public mental health priorities: investing in the evidence. London: UK Department of Health, 2014.
- 6 Ormel J, Petukhova M, Chatterji S, et al. Disability and treatment of specific mental and physical disorders across the world. *Br J Psychiatry* 2008; **192**: 368–75.

Methodology of the SEYLE trial on suicide prevention in schools

We congratulate Danuta Wasserman and colleagues for conducting the first multi-country, randomised controlled trial investigating school-based interventions for the prevention of suicidal behaviour in adolescents (April 18, p1536).¹ However, a few issues must be addressed.

Exclusion of pupils with a history of suicide attempt(s) omits a particularly vulnerable group because past suicide attempts have been shown to be the strongest predictors of future attempts.^{2,3} Efficacy of intervention could be tested in this group, and it is not clear if any specific help was offered to the pupils who had previously attempted suicide. It is not clear why only pupils aged 14–16 years were studied, especially as adolescents (13–19 years old) have greater risk of suicide,⁴ and such restrictive inclusion could limit the generalisability of the study.

Furthermore, no mention is made of pupils (if any) found to have substantial psychopathology, or taking psychotropic medications, during the study. No comment was made on substance use, which is another risk factor associated with adolescent suicide⁵ and could have been included as a covariate during analysis.

Finally, more than half the potential study population (14 267 of

27 099 pupils, or their parents) declined to participate in the study, a major issue in school-based studies.⁶ Even a brief exploration into the reasons for non-participation could help readers understand potential obstacles to these interventions.

We declare no competing interests.

***Ananya Mahapatra, Rishi Gupta**
nnyaa09@gmail.com

Department of Psychiatry and National Drug-Dependence Treatment Centre, and All India Institute of Medical Sciences, New Delhi, India

- 1 Wasserman D, Hoven CW, Wasserman C, et al. School-based suicide prevention programmes: the SEYLE cluster-randomised, controlled trial. *Lancet* 2015; **385**: 1536–44.
- 2 Borowski IW, Ireland M, Resnick MD. Adolescent suicide attempts: risks and protectors. *Pediatrics* 2001; **107**: 485–93.
- 3 Borges G, Angst J, Nock MK, Ruscio AM, Walters EE, Kessler RC. Risk factors for twelve-month suicide attempts in the National Comorbidity Survey Replication (NCS-R). *Psychol Med* 2006; **36**: 1747–57.
- 4 WHO. Preventing suicide: a global imperative. Geneva: World Health Organization, 2014. http://www.who.int/mental_health/suicide-prevention/world_report_2014/en/ (accessed Nov 24, 2014).
- 5 Brent DA, Perper JA, Moritz G, et al. Psychiatric risk factors for adolescent suicide: a case-control study. *J Am Acad Child Adolesc Psychiatry* 1993; **32**: 521–29.
- 6 Robinson J, Cox G, Malone A, et al. A systematic review of school-based interventions aimed at preventing, treating, and responding to suicide-related behavior in young people. *Crisis* 2013; **34**: 164–82.

Over the past decade, UK Child and Adolescent Mental Health Services (CAMHS) have faced a rising tide of ever-increasing demand, coupled with diminishing funding. Cost-effective interventions to target suicide attempts in adolescents are needed urgently. In this bleak context, the SEYLE trial¹ could chart a course to safer waters.

The advantage of preventive intervention targeted to high-risk groups has long been debated against broader population-level approaches.² Results from the SEYLE trial support these wider-ranging methods. Although the outcome seems promising, patients were excluded from the trial if there was evidence of recent suicidal ideation. This exclusion reduces the general



Chalmers/Chris Ryan/Getty Images

application of an otherwise admirably pragmatic trial.

Furthermore, the study highlights the possible danger of extrapolating conclusions from self-report questionnaires. Results from the two trial groups show an illogical reduction in the number of “yes” responses to the question, “Have you ever made an attempt to take your own life?” at 12 months post intervention compared with the same participants at 3 months.

The Health and Social Care reforms of 2012 incentivised CAMHS to move toward reaction instead of prevention. Strategies to manage severe morbidity are prioritised over protective community measures. After proof of cost-effectiveness, integration of the Youth Aware of Mental Health tool into schools might help to redress this imbalance. However, further investigation into the value for recently suicidal young people and clarification of the reliability of SEYLE’s outcome measures must be addressed before the ship sets sail.

We declare no competing interests.

**Georgina Corbet Burcher,
Rob McCutcheon
georgina.corbetburcher@nhs.net*

Royal Free London NHS Foundation Trust, London, UK (GCB); and Institute of Psychiatry, Psychology and Neuroscience, London, UK (RM)

- 1 Wasserman D, Hoven CW, Wasserman C, et al. School-based suicide prevention programmes: the SEYLE cluster-randomised, controlled trial. *Lancet* 2015; **385**: 1536–44.
- 2 Rose G. Mental disorder and the strategies of prevention. *Psychol Med* 1993; **23**: 553–55.

Authors’ reply

We thank Georgina Corbet Burcher, Rob McCutcheon, Ananya Mahapatra, and Rishi Gupta for their comments on our study.¹ The objective of the SEYLE study was to evaluate the primary preventive effects of school-based interventions, therefore only new cases of attempted suicide and severe suicidal ideation were analysed. We agree that the effect of treatment on pre-existing attempts and ideation are of interest. However, evaluation of treatment requires different methods

than evaluation of primary preventive measures as used in the SEYLE study. In compliance with ethical treatment principles, individuals with pre-existing severe suicidal ideation or recent suicide attempts were immediately identified and treated according to the local availability of treatment options. We believe that a collaborative approach between primary prevention and treatment in public mental health and clinical practices is needed to reduce suicide rates.²

We are aware of the limitations of self-report for the measurement of suicidal ideation and attempts. However, prevention is primarily a population-based science and as such, researchers have to rely on self-reporting. Clinical interviews or observer-rated scales are not feasible with large samples, in this case more than 11 000 adolescents. Because self-report was used in the entire SEYLE population (interventions and control), the effect of any potential over-reporting or under-reporting is assumed to be randomly distributed.

Participation rates were mostly related to the complex procedures required by the ethics committees in different countries. Outside the strict experimental requirements of a randomised controlled trial, we believe participation rates would be much higher. For example, according to the follow-up questionnaires with instructors from all SEYLE countries, adolescents were very interested and engaged in the YAM programme.³ The experimental design significantly affected the selection of the age group as well. It was necessary to ensure that recruited pupils were of comparable age between different school systems, and that each participant could be followed up in the same school for at least an entire year.

More than 20 scientific articles have been published based on the SEYLE data, and additional information about health risk behaviours and psychiatric symptoms can therefore be found elsewhere.^{4–6}

We declare no competing interests.

**Danuta Wasserman, Vladimir Carli,
Christina W Hoven, Marco Sarchiapone,
Camilla Wasserman
danuta.wasserman@ki.se*

National Centre for Suicide Research and Prevention of Mental Ill-Health, Karolinska Institutet, Stockholm, Sweden (DW, VC); Division of Child and Adolescent Psychiatry and Department of Psychiatry, New York State Psychiatric Institute, Columbia University, New York, NY, USA (CWH, CW); Department of Epidemiology, Mailman School of Public Health, Columbia University, New York, NY, USA (CWH); Department of Medicine and Health Science, University of Molise, Campobasso, Italy (CW, MS); and National Institute for Health, Migration and Poverty, Rome, Italy (MS)

- 1 Wasserman D, Hoven CW, Wasserman C, et al. School-based suicide prevention programmes: the SEYLE cluster-randomised, controlled trial. *Lancet* 2015; **385**: 1536–44.
- 2 Wasserman D, Durkee T. Strategies in suicide prevention. In: Wasserman D, Wasserman C, eds. *Oxford textbook of suicidology and suicide prevention: a global perspective*. Oxford, UK: Oxford University Press, 2009: 381–84.
- 3 Wasserman C, Hoven CW, Wasserman D, et al. Suicide prevention for youth—a mental health awareness program: lessons learned from the Saving and Empowering Young Lives in Europe (SEYLE) intervention study. *BMC Public Health* 2012; **12**: 776.
- 4 Balazs J, Miklosi M, Keresztesny A, et al. Adolescent subthreshold-depression and anxiety: psychopathology, functional impairment and increased suicide risk. *J Child Psychol Psychiatry* 2013; **54**: 670–77.
- 5 Carli V, Hoven CW, Wasserman C, et al. A newly identified group of adolescents at “invisible” risk for psychopathology and suicidal behavior: findings from the SEYLE study. *World Psychiatry* 2014; **13**: 78–86.
- 6 Kelleher I, Corcoran P, Keeley H, et al. Psychotic symptoms and population risk for suicide attempt: a prospective cohort study. *JAMA Psychiatry* 2013; **70**: 940–48.

Caring for patients with end-stage kidney disease

The renal issue of *The Lancet* emphasises worldwide inequalities in access to renal replacement therapy.^{1,2} Innovative work in Tanzania supported by the International Society of Nephrology³ was particularly interesting. In Malawi, two hospitals provide 15 haemodialysis stations, treating about 60 patients with end-stage kidney disease and with