




STAT 241: Statistics (Advanced Level)

Prof. Mu Zhu • Winter 2014 • University of Waterloo
TR 11:30AM–12:50PM in MC 4007

Transcribed, $\text{\LaTeX}$ ed, and edited by Rob Zimmerman

Note: These are personal lecture notes, not official course materials. They may contain errors (which by default you should attribute to me, Rob Zimmerman, rather than the course instructor). The permanent home of these — and all of my other lecture notes — is <https://rob-zimmerman.github.io/coursenotes>.

Contents

1	Probability Review and Point Estimation	3
	Important Probability Distributions	4
	Maximum Likelihood Estimation	8
	Method-of-Moments Estimator (MOM)	10
	Constrained Optimization and Applied Examples	13
	Properties of Estimators: Bias, Efficiency, and MSE	17
	Sampling Distributions and Order Statistics	18

2	Asymptotic Theory	23
	Asymptotics and Moment Generating Functions	23
	The Weak Law of Large Numbers	26
3	Interval Estimation and Hypothesis Testing	28
	Confidence Intervals and Pivotal Quantities	28
	Fisher Information and the Cramér-Rao Bound	30
	Sample Size Calculations	33
	Hypothesis Testing	34
	Likelihood Ratio Tests and the Neyman-Pearson Lemma	35
	Power Analyses	36
	Composite Hypotheses	38
	Interval Estimation versus Hypothesis Testing	40
	Generalized Likelihood Ratio Tests and Wilks' Theorem	40
	P-values	43
4	Classical Inference for Normal Models	45
	Student's t -Distribution and the One-Sample t -Test	45
	Two-Sample t -Test	51
5	Regression and Categorical Data Analysis	53
	Simple Linear Regression	53
	Fitted Values and Prediction Intervals	58
	Standardized Regression	60
	Pearson's Goodness-of-Fit Test	61
	Applications of the Multinomial Model	64
6	Study Design and Sampling	66
	Stratified Sampling and Paired Analysis	66
	Paired Analysis	67
	Graphical Methods	69
	Empirical Studies	69
7	Causation and Bayesian Statistics	70
	Causation versus Association	70
	Bayesian Statistics	71
	Conjugate Priors and Jeffrey's Rule	73
	Empirical Bayes	76
	Posterior Odds	79
	Appendix: Lecture and Tutorial Index	81

Lecture 1 — Tuesday, January 07, 2014

1. Probability Review and Point Estimation

In probability, we begin with $X \sim f(x; \theta)$; that is, a random variable with a known distribution. For example, we may ask for $\mathbb{E}[X]$, $\text{Var}(X)$, and $\mathbb{P}(a \leq X \leq b)$. These all depend on θ , which is assumed known. **Statistics** is, in some sense, the inverse of probability. We *observe* $X_1, X_2, \dots, X_n \sim f(x; \theta)$, and we want to estimate the unknown parameter θ from the data.

There are several levels of **statistical inference**, which we study in the first half of the course. First, **point estimation** seeks a function $\hat{\theta}(X_1, X_2, \dots, X_n)$ called an estimator which is supposed to approximate the unknown value θ . We then ask how to construct such a function and how to compare competing estimators. Second, **interval estimation** aims to find lower and upper bounds $\ell(X_1, X_2, \dots, X_n)$ and $u(X_1, X_2, \dots, X_n)$ such that

$$\mathbb{P}(\ell(X_1, \dots, X_n) < \theta < u(X_1, \dots, X_n)) = 1 - \alpha$$

for some small $\alpha > 0$. Third, **hypothesis testing** asks the following: given $H_0 : \theta = \theta_0$ and $H_A : \theta = \theta_A$, which hypothesis is more plausible given the observations $X_1, \dots, X_n$? These three topics comprise the first half of the course.

To start the second half of the course, several important **applied models** will be studied. First, the classical **normal model** $X_1, X_2, \dots, X_n \stackrel{\text{iid}}{\sim} N(\mu, \sigma^2)$ with σ^2 known; this is a simplified canonical model. Second, we revisit the same normal model with σ^2 unknown and investigate what changes. Third, we study the **two-sample case**: $X_1, X_2, \dots, X_n \stackrel{\text{iid}}{\sim} N(\mu_A, \sigma^2)$ and $Y_1, Y_2, \dots, Y_m \stackrel{\text{iid}}{\sim} N(\mu_B, \sigma^2)$. Fourth, we study **simple linear regression** where $Y_i \stackrel{\text{indep}}{\sim} N(\alpha + \beta X_i, \sigma^2)$ for $i = 1, \dots, n$. Fifth, we study the **multinomial distribution** $(X_1, X_2, \dots, X_k) \stackrel{\text{vector}}{\sim} \text{Multinomial}(n; p_1, p_2, \dots, p_k)$ with $p_1 + \dots + p_k = 1$.

We end with **Bayesian statistics**. The starting point of Bayesian statistics is to treat θ as a random variable. We write $\pi(\theta)$ for the prior distribution, which represents our beliefs about the true value prior to collecting data. The posterior distribution is given by

$$\pi(\theta \mid X_1, \dots, X_n) = \frac{f(X_1, \dots, X_n \mid \theta) \cdot \pi(\theta)}{\int f(X_1, \dots, X_n \mid \theta) \cdot \pi(\theta) \, d\theta},$$

which is a direct consequence of Bayes' theorem:

$$\mathbb{P}(B_i | A) = \frac{\mathbb{P}(A | B_i) \cdot \mathbb{P}(B_i)}{\sum_{j=1}^m \mathbb{P}(A | B_j) \cdot \mathbb{P}(B_j)}.$$

In practice, the integral in the denominator of the posterior distribution can be difficult to compute, especially when $\theta \in \mathbb{R}^n$; the distribution is then studied indirectly using more advanced methods. For this reason, Bayesian methods are not a major focus of this course.

Important Probability Distributions

The most basic distribution is the **Bernoulli distribution**. If $X \sim \text{Bernoulli}(p)$, then X takes the value 1 with probability p and the value 0 with probability $1 - p$. The probability mass function is $f(x) = p^x(1 - p)^{1-x}$ for $x = 0, 1$. We have $\mathbb{E}[X] = p$ and $\text{Var}(X) = p(1 - p)$.

The **binomial distribution** arises as the number of successes in n independent Bernoulli trials. If $X \sim \text{Bin}(n, p)$, then X can take on the values $0, 1, \dots, n$. The probability mass function is

$$f(x) = \binom{n}{x} p^x (1 - p)^{n-x}.$$

The expectation is $\mathbb{E}[X] = \mathbb{E}[X_1] + \dots + \mathbb{E}[X_n] = np$, and the variance is $\text{Var}(X) = np(1 - p)$.

Lecture 2 — Thursday, January 09, 2014

Let us study the **multinomial distribution**. Consider a vector of random variables

$$(X_1, \dots, X_k) \sim \text{Multinomial}(n; p_1, \dots, p_k).$$

The distribution function is

$$f_{X_1, \dots, X_k}(x_1, x_2, \dots, x_k) = \frac{n!}{x_1! x_2! \dots x_k!} p_1^{x_1} p_2^{x_2} \dots p_k^{x_k}$$

subject to

$$\sum_{i=1}^k p_i = 1 \quad \text{and} \quad x_1, \dots, x_k \in \{0, 1, \dots, n\} \quad \text{and} \quad \sum_{i=1}^k x_i = n.$$

This is the joint distribution of the random vector $(X_1, \dots, X_k)$. As a special case, consider $k = 2$ (the trinomial case), which involves only two free random variables with distribution function

$$f_{X_1, X_2}(x_1, x_2) = \frac{n!}{x_1! x_2! (n - x_1 - x_2)!} p_1^{x_1} p_2^{x_2} (1 - p_1 - p_2)^{n - x_1 - x_2}.$$

Two natural questions arise. First, what are the marginal distributions of X_1 and X_2 ? Marginalizing out X_2 gives

$$\begin{aligned} f_{X_1}(x_1) &= \sum_{x_2=0}^{n-x_1} f(x_1, x_2) = \sum_{x_2=0}^{n-x_1} \frac{n!}{x_1! x_2! (n - x_1 - x_2)!} p_1^{x_1} p_2^{x_2} (1 - p_1 - p_2)^{n - x_1 - x_2} \\ &= \frac{n!}{(n - x_1)! x_1!} p_1^{x_1} \sum_{x_2=0}^{n-x_1} \frac{(n - x_1)!}{x_2! (n - x_1 - x_2)!} p_2^{x_2} (1 - p_1 - p_2)^{n - x_1 - x_2} \\ &= \binom{n}{x_1} p_1^{x_1} (1 - p_1)^{n - x_1} \text{ by the binomial theorem.} \end{aligned}$$

Thus

$$X_1 \sim \text{Binomial}(n, p_1).$$

By symmetry, $X_2 \sim \text{Binomial}(n, p_2)$.

Second, what is the conditional distribution of $X_2 \mid X_1 = x_1$? We have

$$\begin{aligned} f_{X_2|X_1}(x_2 \mid x_1) &= \frac{f(x_1, x_2)}{f_{X_1}(x_1)} = \frac{(n - x_1)!}{x_2! (n - x_1 - x_2)!} \frac{p_2^{x_2} (1 - p_1 - p_2)^{n - x_1 - x_2}}{(1 - p_1)^{n - x_1}} \\ &= \binom{n - x_1}{x_2} \left(\frac{p_2}{1 - p_1} \right)^{x_2} \left(1 - \frac{p_2}{1 - p_1} \right)^{n - x_1 - x_2}. \end{aligned}$$

Thus,

$$X_2 \mid X_1 = x_1 \sim \text{Binomial} \left(n - x_1, \frac{p_2}{1 - p_1} \right).$$

More generally,

$$(X_1, X_2, \dots, X_{k-1}) \mid X_k = x_k \sim \text{Multinomial} \left(n - x_k, \frac{p_1}{1 - p_k}, \frac{p_2}{1 - p_k}, \dots, \frac{p_{k-1}}{1 - p_k} \right).$$

Next, recall the **Poisson distribution**. A random variable $X \sim \text{Poisson}(\lambda)$ takes on values

$X = 0, 1, 2, \dots$ with probability mass function

$$f_X(x) = \frac{e^{-\lambda} \lambda^x}{x!}, \quad x \in \{0, 1, 2, \dots\}.$$

This distribution is commonly used as a model for rare events. To motivate this, start from the binomial model $f_X = \binom{n}{x} p^x (1-p)^{n-x}$, and consider the limit as $n \rightarrow \infty$, $p \rightarrow 0$, such that $np = \lambda$; that is, the number of trials n increases, but the success probability p of each trial decreases in such a way that the product np remains constant. Substituting $p = \frac{\lambda}{n}$, we get

$$\begin{aligned} f_X(x) &= \binom{n}{x} \left(\frac{\lambda}{n}\right)^x \left(1 - \frac{\lambda}{n}\right)^{n-x} = \frac{n!}{x!(n-x)!} \left(\frac{\lambda}{n}\right)^x \left(1 - \frac{\lambda}{n}\right)^{n-x} \\ &= \frac{n(n-1)\dots(n-x+1)}{n \cdot n \dots n} \frac{\lambda^x}{x!} \left(1 - \frac{\lambda}{n}\right)^{-x} \left(1 - \frac{\lambda}{n}\right)^n \\ &\xrightarrow{n \rightarrow \infty} \frac{\lambda^x}{x!} e^{-\lambda}, \end{aligned}$$

using the exponential limit. Thus, the Poisson approximation to the binomial distribution is most appropriate when p is small and n is large.

Now we can compute the expectation:

$$\mathbb{E}[X] = \sum_{x=0}^{\infty} x \cdot f(x) = \sum_{x=0}^{\infty} x \cdot \frac{e^{-\lambda} \lambda^x}{x!} = 0 + \sum_{x=1}^{\infty} \frac{e^{-\lambda} \lambda^{(x-1)}}{(x-1)!} \cdot \lambda = e^{-\lambda} \cdot \lambda \sum_{y=0}^{\infty} \frac{\lambda^y}{y!} = \lambda = np$$

(with $y = x - 1$).

The **exponential distribution** is very important. A random variable $X \sim \text{Exp}(\lambda)$ has density function

$$f_X(x) = \lambda e^{-\lambda x}$$

for $x > 0$. This distribution is often used as a model of failure time.

Let us construct a **Poisson process**. Consider a stochastic process $X(t)$ which satisfies three properties:

1. $X(0) = 0$.
2. $X(t) - X(s) \sim \text{Poisson}(\lambda(t-s))$, which models the number of events between times s and t .
3. Increments over disjoint time intervals, such as $X(t_2) - X(t_1)$ and $X(t_4) - X(t_3)$, are independent.

Let T be the waiting time from one event to the next. Then $\mathbb{P}(T > t)$ is the probability of no event on a time interval of length t , so

$$\mathbb{P}(T > t) = \frac{e^{-\lambda t}(\lambda t)^0}{0!} = e^{-\lambda t}.$$

Hence

$$F_T(t) = \mathbb{P}(T \leq t) = 1 - e^{-\lambda t},$$

and

$$f_T(t) = \frac{d}{dt}F_T(t) = \lambda e^{-\lambda t}.$$

Therefore $T \sim \text{Exp}(\lambda)$, so the exponential distribution models waiting times between events.

Finally, recall the **normal distribution**. A random variable $X \sim N(\mu, \sigma^2)$ has density function

$$f_X(x) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(x-\mu)^2}{2\sigma^2}}.$$

We have $\mathbb{E}[X] = \mu$ and $\text{Var}(X) = \sigma^2$. Recall that the **standard normal** model is $Z \sim N(0, 1)$. Some rules of thumb are useful:

$$\mathbb{P}(\mu - \sigma < X < \mu + \sigma) \approx 68\% \quad \text{and} \quad \mathbb{P}(\mu - 2\sigma < X < \mu + 2\sigma) \approx 95\%.$$

Example 2.1 Consider the following application. Suppose that $X \sim N(\mu_A, \sigma_A^2)$ if the observation is of type A , and $X \sim N(\mu_B, \sigma_B^2)$ if it is of type B . Without loss of generality, assume $\mu_A > \mu_B$. Given an observation X from either type, we would like to infer which type was more likely to have generated X . A reasonable **decision rule** is to claim type A if $X > c$ for some $c \in \mathbb{R}$. We need to optimize c to minimize the error. The probability of an erroneous classification is given by

$$\mathbb{P}(\text{error}) = \mathbb{P}(X > c \mid B) \cdot \mathbb{P}(B) + \mathbb{P}(X \leq c \mid A) \cdot \mathbb{P}(A),$$

by the law of total probability. Assume $\mathbb{P}(A) = \mathbb{P}(B) = \frac{1}{2}$. Then

$$\mathbb{P}(\text{error}) = \frac{1}{2} (1 - F_B(c) + F_A(c)).$$

Differentiating with respect to c , we get

$$\frac{d}{dc} \mathbb{P}(\text{error}) = 0 \Rightarrow f_B(c) = f_A(c).$$

If $\sigma_A = \sigma_B$ (that is, we have a location shift), we see that

$$c = \frac{\mu_A + \mu_B}{2}.$$

Lecture 3 — Tuesday, January 14, 2014

Maximum Likelihood Estimation

The first step in statistical inference is to obtain data and use it to estimate the parameters of the distribution.

Assume we have already selected a model (i.e., the distributional family from which the data arises). Suppose $X_1, X_2, \dots, X_n \stackrel{\text{iid}}{\sim} f(x; \theta)$. We want to estimate the unknown parameter θ using the likelihood function. Recall that the joint density or mass function of $X_1, X_2, \dots, X_n$ can be written as $\prod_{i=1}^n f(X_i; \theta)$ due to independence. The **likelihood function** is defined as

$$L(\theta) = \prod_{i=1}^n f(X_i; \theta).$$

The **maximum likelihood estimate (MLE)** is

$$\hat{\theta}_{\text{MLE}} = \operatorname{argmax} L(\theta).$$

That is, the MLE is the value of θ that maximizes the likelihood function $L(\theta)$.

In practice, we often maximize $l(\theta) = \log(L(\theta))$, called the **log-likelihood function**, using the natural logarithm. This transformation changes the scale but not the maximizer. Differentiating a large product is cumbersome, while the log-likelihood converts products into sums which are much easier to handle.

Example 3.1 Suppose $X_1, X_2, \dots, X_n \stackrel{\text{iid}}{\sim} \text{Bernoulli}(p)$. Then the likelihood function is

$$L(p) = \prod_{i=1}^n p^{X_i} (1-p)^{1-X_i} = p^{\sum_{i=1}^n X_i} (1-p)^{n - \sum_{i=1}^n X_i}.$$

Let us maximize this. We have

$$l(p) = \sum_{i=1}^n X_i \log(p) + (n - \sum_{i=1}^n X_i) \log(1 - p).$$

Differentiating yields

$$\frac{d}{dp} l(p) = \frac{\sum_{i=1}^n X_i}{p} - \frac{n - \sum_{i=1}^n X_i}{1 - p} = 0 \implies \sum_{i=1}^n X_i = np \implies \hat{p}_{\text{MLE}} = \frac{\sum_{i=1}^n X_i}{n} = \bar{X}.$$

Strictly speaking, we should check that

$$\frac{d^2}{dp^2} l(p) = -\frac{\sum_{i=1}^n X_i}{p^2} - \frac{n - \sum_{i=1}^n X_i}{(1 - p)^2} < 0,$$

which verifies that this is indeed a maximum. We usually omit this second-derivative check in routine calculations.

Example 3.2 Suppose $X_1, X_2, \dots, X_n \stackrel{\text{iid}}{\sim} \text{Exp}(\lambda)$. Then the likelihood function is

$$L(\lambda) = \prod_{i=1}^n \lambda e^{-\lambda X_i} = \lambda^n e^{-\lambda \sum_{i=1}^n X_i},$$

and

$$l(\lambda) = n \log(\lambda) - \lambda \sum_{i=1}^n X_i.$$

We have

$$l'(\lambda) = \frac{n}{\lambda} - \sum_{i=1}^n X_i = 0 \implies \hat{\lambda}_{\text{MLE}} = \frac{n}{\sum_{i=1}^n X_i} = \frac{1}{\bar{X}},$$

and it is easy to perform a second-derivative test to verify that $\hat{\lambda}_{\text{MLE}}$ is a global maximum.

This makes intuitive sense because

$$\mathbb{E}[X_i] = \int_0^\infty x \lambda e^{-\lambda x} dx = \frac{1}{\lambda}$$

for all i , and it is reasonable to estimate λ by setting $\bar{X} = \frac{1}{\lambda}$.

Method-of-Moments Estimator (MOM)

Suppose $X_1, X_2, \dots, X_n \stackrel{\text{iid}}{\sim} f(x; \theta)$. We can calculate

$$\mathbb{E}[X_i] = \int x \cdot f(x; \theta) dx = g_1(\theta), \quad \mathbb{E}[X_i^2] = \int x^2 \cdot f(x; \theta) dx = g_2(\theta),$$

and so on. These are the **moments** of X_i . The **method of moments (MOM)** consists of substituting $\mathbb{E}[X_i^m]$ with

$$\frac{1}{n} \sum_{i=1}^n X_i^m$$

to obtain equations

$$\frac{1}{n} \sum_{i=1}^n X_i = g_1(\theta), \quad \frac{1}{n} \sum_{i=1}^n X_i^2 = g_2(\theta),$$

and so on, which we then solve for θ . We can choose whichever moments are convenient. This method has some arbitrariness, so it is not as elegant as maximum likelihood estimation. Sometimes the two methods yield the same estimator, and sometimes they do not.

Example 3.3 Suppose $X_1, X_2, \dots, X_n \stackrel{\text{iid}}{\sim} N(\mu, \sigma^2)$. Let us apply both methods to estimate (μ, σ^2) . For the maximum likelihood approach, we have

$$L(\mu, \sigma^2) = \prod_{i=1}^n \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(X_i - \mu)^2}{2\sigma^2}} = \left(\frac{1}{\sqrt{2\pi}}\right)^n \left(\frac{1}{\sigma}\right)^n e^{-\frac{\sum_{i=1}^n (X_i - \mu)^2}{2\sigma^2}}.$$

Therefore

$$l(\mu, \sigma^2) = \text{constant} - n \log(\sigma) - \frac{\sum_{i=1}^n (X_i - \mu)^2}{2\sigma^2}.$$

Differentiating with respect to μ , we get

$$\frac{\partial}{\partial \mu} l(\mu, \sigma^2) = \frac{\sum_{i=1}^n 2(X_i - \mu)}{2\sigma^2} = 0 \implies \sum_{i=1}^n (X_i - \mu) = 0 \implies \sum_{i=1}^n X_i - n\mu = 0$$

which implies that

$$\hat{\mu}_{\text{MLE}} = \frac{\sum_{i=1}^n X_i}{n} = \bar{X}.$$

Differentiating with respect to σ , we have

$$\frac{\partial}{\partial \sigma} l(\mu, \sigma^2) = -\frac{n}{\sigma} - (-2) \frac{\sum_{i=1}^n (X_i - \mu)^2}{2\sigma^3} = 0.$$

(Note: it is fair to assume $\sigma \neq 0$.) This implies

$$\hat{\sigma}_{\text{MLE}}^2 = \frac{\sum_{i=1}^n (X_i - \mu)^2}{n} = \frac{\sum_{i=1}^n (X_i - \bar{X})^2}{n}.$$

Now for the method of moments. We have

$$\mathbb{E}[X_i] = \mu \quad \text{and} \quad \mathbb{E}[X_i^2] = \text{Var}(X_i) + \mathbb{E}[X_i]^2 = \sigma^2 + \mu^2.$$

This gives $\hat{\mu}_{\text{MOM}} = \bar{X} = \hat{\mu}_{\text{MLE}}$ and

$$\frac{1}{n} \sum_{i=1}^n X_i^2 = \sigma^2 + \mu^2,$$

which implies

$$\hat{\sigma}_{\text{MOM}}^2 = \frac{1}{n} \sum_{i=1}^n X_i^2 - \bar{X}^2.$$

Officially we are done. But we can verify that the two estimates of σ^2 agree:

$$\begin{aligned}
 \hat{\sigma}_{\text{MOM}}^2 &= \frac{\sum_{i=1}^n X_i^2 - n\bar{X}^2}{n} \\
 &= \frac{1}{n} \sum_{i=1}^n X_i^2 + \bar{X}^2 - \frac{2\bar{X}}{n} \sum_{i=1}^n X_i \\
 &= \frac{1}{n} \sum_{i=1}^n (X_i^2 + \bar{X}^2 - 2X_i\bar{X}) \\
 &= \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2 \\
 &= \hat{\sigma}_{\text{MLE}}^2.
 \end{aligned}$$

The MLE and the MOM estimator are not the same in general.

Example 3.4 Suppose $X_1, X_2, \dots, X_n \stackrel{\text{iid}}{\sim} \text{Unif}(0, \theta)$. For the method of moments, we have

$$\mathbb{E}[X_i] = \int_0^\theta x \cdot \frac{1}{\theta} dx = \frac{\theta}{2}$$

for all i , which implies $\bar{X} = \frac{\theta}{2}$, and hence $\hat{\theta}_{\text{MOM}} = 2\bar{X}$. Now for maximum likelihood, we have

$$L(\theta) = \prod_{i=1}^n f(X_i; \theta) = \left(\frac{1}{\theta}\right)^n.$$

We do not need the logarithm here, since the likelihood has a simple form. There is an implicit constraint: $X_1, X_2, \dots, X_n \in (0, \theta)$, so $\theta \geq \max\{X_1, \dots, X_n\}$. Since the likelihood is decreasing in θ , we obtain $\hat{\theta}_{\text{MLE}} = \max\{X_1, \dots, X_n\}$. This is a constrained maximization problem. The estimator is sensible, although it always underestimates the true parameter; we will later see why it is still preferable.

The key point is that both procedures produce estimators that are functions of the data $X_1, \dots, X_n$.

Lecture 4 — Thursday, January 16, 2014

Constrained Optimization and Applied Examples

When we perform maximum likelihood estimation, we are sometimes constrained by the setup or parameter space of the statistical model.

Example 4.1 Suppose we observe n independent trials, where each trial falls into exactly one of K categories with probabilities $p_1, p_2, \dots, p_K$. Let X_k denote the number of observations in category k . Then

$$(X_1, X_2, \dots, X_K) \sim \text{Multinomial}(n; p_1, p_2, \dots, p_K),$$

with constraints $p_k \geq 0$ for $k = 1, \dots, K$ and

$$\sum_{k=1}^K p_k = 1.$$

Let us determine the MLE of $(p_1, \dots, p_K)$. For the count vector $(X_1, \dots, X_K)$, satisfying

$$\sum_{k=1}^K X_k = n,$$

the likelihood is

$$L(p_1, \dots, p_K) = \frac{n!}{X_1! \cdots X_K!} \prod_{k=1}^K p_k^{X_k},$$

so the log-likelihood is

$$l(p_1, \dots, p_K) = \text{constant} + \sum_{k=1}^K X_k \log(p_k).$$

We cannot maximize this by setting the partial derivatives of l equal to zero independently, because the parameters must satisfy

$$\sum_{k=1}^K p_k = 1.$$

Instead, we introduce a Lagrange multiplier and consider

$$\mathcal{L}(p_1, \dots, p_K, \lambda) = \sum_{k=1}^K X_k \log(p_k) + \lambda \left(\sum_{k=1}^K p_k - 1 \right).$$

Differentiating gives

$$\frac{\partial \mathcal{L}}{\partial p_k} = \frac{X_k}{p_k} + \lambda = 0, \quad k = 1, \dots, K,$$

and

$$\frac{\partial \mathcal{L}}{\partial \lambda} = \sum_{k=1}^K p_k - 1 = 0.$$

Hence

$$p_k = -\frac{X_k}{\lambda}, \quad k = 1, \dots, K.$$

Summing over k , we obtain

$$1 = \sum_{k=1}^K p_k = -\frac{1}{\lambda} \sum_{k=1}^K X_k = -\frac{n}{\lambda},$$

so $\lambda = -n$. Substituting this back yields

$$\hat{p}_k^{\text{MLE}} = \frac{X_k}{n}, \quad k = 1, \dots, K.$$

This is intuitive: the MLE of each category probability is the observed sample proportion, and it also agrees with $\hat{p}_k^{\text{MOM}}$.

Example 4.2 (Hardy–Weinberg model) Suppose we have genotypes AA, Aa, and aa with probabilities $(1 - \theta)^2$, $2\theta(1 - \theta)$, and θ^2 , respectively; this is a specifically parametrized multinomial model. We want to estimate θ . Suppose we have data X_1, X_2, X_3 . Then

$$L(\theta) = \frac{n!}{X_1! X_2! X_3!} [(1 - \theta)^2]^{X_1} [2\theta(1 - \theta)]^{X_2} [\theta^2]^{X_3} = (\text{const.}) \cdot (1 - \theta)^{2X_1 + X_2} \cdot \theta^{X_2 + 2X_3}.$$

Therefore

$$l(\theta) = (\text{const.}) + (2X_1 + X_2) \log(1 - \theta) + (X_2 + 2X_3) \log(\theta),$$

and so

$$\frac{d}{d\theta} l(\theta) = 0$$

implies

$$\hat{\theta}_{\text{MLE}} = \frac{X_2 + 2X_3}{2n}.$$

For example, suppose we have data

$$X_1 = 37, X_2 = 46, X_3 = 17.$$

$$\text{Then } \hat{\theta}_{\text{MLE}} = \frac{46+2 \cdot 17}{2(37+46+17)} = 0.4.$$

We can check the Hardy-Weinberg model by comparing actual observations with model predictions. The model predicts that among 100 people (properly chosen), we would have

$$n(1 - \hat{\theta})^2 = 100(1 - 0.4)^2 = 36$$

individuals with genotype AA, 48 with Aa, and 16 with aa. These are quite close to the observed values. The difference is tolerable, and we could say that the Hardy-Weinberg model is reasonable for this data. Note that $p_1 + p_2 + p_3 = (1 - \theta + \theta)^2 = 1$.

Example 4.3 Suppose $Y_i \stackrel{\text{indep}}{\sim} \text{Poisson}(\theta x_i)$ for $i = 1, \dots, n$. These variables are independent but not identically distributed; however, they still belong to the same distributional family. The values x_i are **covariates** (or **predictors**). We have

$$L(\theta) = \prod_{i=1}^n \frac{e^{-(\theta x_i)} (\theta x_i)^{Y_i}}{Y_i!},$$

and

$$l(\theta) = \sum_{i=1}^n (-\theta x_i + Y_i \log(\theta x_i)) + \text{constant} = -\theta \sum_{i=1}^n x_i + \log(\theta) \sum_{i=1}^n Y_i + \text{constant}.$$

Differentiating,

$$\frac{d}{d\theta} l(\theta) = 0$$

yields

$$-\sum_{i=1}^n x_i + \frac{1}{\theta} \sum_{i=1}^n Y_i = 0,$$

which implies

$$\hat{\theta}_{\text{MLE}} = \frac{\sum_{i=1}^n Y_i}{\sum_{i=1}^n x_i}.$$

Example 4.4 Suppose $X_i \stackrel{\text{indep}}{\sim} \text{Poisson}(V_i\theta)$ for $i = 1, \dots, n$ (for example, X_i could be the count of bacteria from the i th laboratory sample of volume V_i). Let

$$Y_i = \begin{cases} 1, & X_i > 0 \\ 0, & X_i = 0 \end{cases}.$$

We want to estimate θ using only the data $(V_1, Y_1), \dots, (V_n, Y_n)$, thereby bypassing direct measurements of $X_1, X_2, \dots$. This is a realistic measurement scenario.

Clearly $Y_i \sim \text{Bernoulli}(p_i)$, where

$$p_i = \mathbb{P}(X_i > 0) = 1 - \mathbb{P}(X_i = 0) = 1 - \frac{e^{-(V_i\theta)}(V_i\theta)^0}{0!} = 1 - e^{-V_i\theta}.$$

Therefore

$$L(\theta) = \prod_{i=1}^n (1 - e^{-V_i\theta})^{Y_i} (e^{-V_i\theta})^{1-Y_i},$$

and

$$l(\theta) = \sum_{i=1}^n \left[Y_i \log(1 - e^{-V_i\theta}) + (1 - Y_i)(-V_i\theta) \right].$$

Differentiating yields

$$l'(\theta) = \sum_{i=1}^n \left[Y_i \frac{V_i e^{-V_i\theta}}{1 - e^{-V_i\theta}} - (1 - Y_i)V_i \right],$$

but there does not exist a closed-form solution to $l'(\theta) = 0$.

When $l'(\theta) = 0$ cannot be solved in closed form, we must resort to numerical optimization — for example, a root-finding method such as **Newton's method**. To solve $f(x) = 0$, we start from an

initial guess x_0 and iterate

$$x_1 = x_0 - \frac{f(x_0)}{f'(x_0)}.$$

In our example, this becomes

$$\theta_1 = \theta_0 - \frac{l'(\theta_0)}{l''(\theta_0)}.$$

A variation of Newton's method, the **Fisher scoring algorithm**, replaces $l''(\theta)$ with $\mathbb{E}[l''(\theta)]$. This approach is often more stable for maximum likelihood estimation problems.

Lecture 5 — Tuesday, January 21, 2014

Properties of Estimators: Bias, Efficiency, and MSE

We now ask a central question: how do we assess the quality of an estimator?

To illustrate, let $Y \sim \text{Binomial}(n, p)$. *Before* we flip a coin n times, Y is a random variable; *after* observing the flips, it becomes data. It is important to keep this connection between data and distribution in mind.

Recall [Example 3.4](#), where the method of moments and maximum likelihood give different estimators. At this point, it is not immediately clear which estimator should be preferred. The claim is that MLE performs better on average, especially as the sample size grows, and this can be justified analytically.

The estimator $\hat{\theta} = \hat{\theta}(X_1, \dots, X_n)$ is a function of the random variables $X_1, \dots, X_n$ and hence is a random variable itself. The distribution of $\hat{\theta}$ is called the **sampling distribution**.

Definition 5.1 A estimator $\hat{\theta}$ is **unbiased** if it's correctly centered; that is, if

$$\mathbb{E}[\hat{\theta}] = \theta.$$

Definition 5.2 Suppose $\hat{\theta}_1$ and $\hat{\theta}_2$ are both unbiased estimators of θ . We say $\hat{\theta}_1$ is more **efficient** than $\hat{\theta}_2$ if $\text{Var}(\hat{\theta}_1) < \text{Var}(\hat{\theta}_2)$.

Definition 5.3 The **mean-squared error (MSE)** of an estimator $\hat{\theta}$ is defined as

$$\text{MSE}(\hat{\theta}) = \mathbb{E} \left[(\hat{\theta} - \theta)^2 \right].$$

Theorem 5.4 $\text{MSE}(\hat{\theta}) = \text{Bias}(\hat{\theta})^2 + \text{Var}(\hat{\theta})$.

Proof. A direct calculation gives

$$\begin{aligned} \text{MSE}(\hat{\theta}) &= \mathbb{E} \left[(\hat{\theta} - \theta)^2 \right] \\ &= \mathbb{E} \left[(\hat{\theta} - \mathbb{E}[\hat{\theta}] + \mathbb{E}[\hat{\theta}] - \theta)^2 \right] \\ &= \mathbb{E} \left[(\hat{\theta} - \mathbb{E}[\hat{\theta}])^2 \right] + 2(\mathbb{E}[\hat{\theta}] - \theta)\mathbb{E}[\hat{\theta} - \mathbb{E}[\hat{\theta}]] + (\mathbb{E}[\hat{\theta}] - \theta)^2 \\ &= \text{Var}(\hat{\theta}) + \text{Bias}(\hat{\theta})^2. \end{aligned}$$

The middle term vanishes because $\mathbb{E}[\hat{\theta} - \mathbb{E}[\hat{\theta}]] = 0$. □

Lecture 6 — Thursday, January 23, 2014

Sampling Distributions and Order Statistics

The estimator $\hat{\theta} = g(X_1, X_2, \dots, X_n)$ is a random variable, and we want to find its distribution. Recall that we often had $\hat{\theta} = \bar{X}$ or $\hat{\theta} = X_{\max}$. In the former case, to understand the sampling distribution of $\hat{\theta}$ it usually suffices to know the distribution of $X_1 + \dots + X_n$. Let us start with a simple example.

Example 6.1 Suppose $X, Y \stackrel{\text{iid}}{\sim} \text{Unif}(0, 1)$. We want the distribution of $Z = X + Y$. We have

$$F_Z(z) = \mathbb{P}(Z \leq z) = \mathbb{P}(X + Y \leq z) = \iint_{x+y \leq z} f_{X,Y}(x, y) \, dx \, dy = \iint_{x+y \leq z} f_X(x) f_Y(y) \, dx \, dy$$

$$= \iint_{x+y \leq z} 1 \cdot 1 \, dx \, dy = \text{area} \cdot \text{height} = \text{area} = \begin{cases} z^2/2, & 0 \leq z \leq 1 \\ 1 - \frac{(2-z)^2}{2}, & z \geq 1 \end{cases}.$$

We can now find the density:

$$f_Z(z) = \frac{d}{dz} F_Z(z) = \begin{cases} z, & z \leq 1 \\ 2 - z, & z \geq 1 \end{cases}.$$

The following diagrams illustrate the geometric calculation of the cdf $F_Z(z) = \mathbb{P}(X + Y \leq z)$.




Figure 1: Geometric regions used to compute $F_Z(z) = \mathbb{P}(X + Y \leq z)$ for the cases $z \leq 1$ and $z > 1$.

The resulting density function $f_Z(z)$ is that of a **triangular distribution**:




Figure 2: Density of $Z = X + Y$ when X and Y are independent $\text{Unif}(0, 1)$ random variables.

Adding two uniform random variables gives a joined piecewise linear density. Adding three gives a piecewise quadratic density. Adding four, five, and so on gives piecewise cubic, quartic, and higher-order polynomial densities, which join smoothly. This behavior is consistent with the central limit theorem. More generally, finding the distribution of $g(X_1, \dots, X_n)$ typically requires n -dimensional integration, which is difficult in practice. This is why asymptotic approximations, such as the central limit theorem, are often used.

Example 6.2 Suppose $X_1, \dots, X_n \stackrel{\text{iid}}{\sim} f_X(x)$ where f_X is a density, and let $Y = \max\{X_1, \dots, X_n\}$. What is the density $f_Y(y)$? We have

$$F_Y(y) = \mathbb{P}(Y \leq y) = \mathbb{P}(X_1, \dots, X_n \leq y) = \mathbb{P}(X_1 \leq y) \cdot \mathbb{P}(X_2 \leq y) \cdots \mathbb{P}(X_n \leq y) = [F_X(y)]^n.$$

Differentiating, we obtain

$$f_Y(y) = \frac{d}{dy} F_Y(y) = n \cdot [F_X(y)]^{n-1} \cdot f_X(y).$$

This formula is quite general.

Example 6.3 Suppose $X_1, \dots, X_n \stackrel{\text{iid}}{\sim} \text{Unif}(0, \theta)$. We know from the method of moments that $\hat{\theta}_{\text{MOM}} = 2\bar{X}$, and from maximum likelihood estimation that $\hat{\theta}_{\text{MLE}} =$

$\max\{X_1, \dots, X_n\} = Y$. We have $f_X(y) = \frac{1}{\theta}$ for $y \in (0, \theta)$ and

$$F_X(y) = \mathbb{P}(X \leq y) = \frac{y}{\theta}$$

for $y \in (0, \theta)$. From the general formula, the sampling distribution of $\hat{\theta}_{\text{MLE}}$ is $f_Y(y) = n \left(\frac{y}{\theta}\right)^{n-1} \cdot \frac{1}{\theta}$. We can also calculate the bias using

$$\mathbb{E}[Y] = \int_0^\theta y \cdot f_Y(y) \, dy = \int_0^\theta y \cdot n \left(\frac{y}{\theta}\right)^{n-1} \cdot \frac{1}{\theta} \, dy = \frac{n}{\theta^n} \cdot \left[\frac{y^{n+1}}{n+1} \right]_0^\theta = \frac{n}{n+1} \theta \neq \theta,$$

so

$$\text{Bias}(\hat{\theta}_{\text{MLE}}) = \frac{\theta}{n}.$$

The estimator $\hat{\theta}_{\text{MLE}}$ always underestimates θ , but it approaches θ as $n \rightarrow \infty$. For the variance of $\hat{\theta}_{\text{MLE}}$, recall that $\text{Var}(Y) = \mathbb{E}[Y^2] - \mathbb{E}[Y]^2$. We compute

$$\mathbb{E}[Y^2] = \int_0^\theta y^2 \cdot n \left(\frac{y}{\theta}\right)^{n-1} \cdot \frac{1}{\theta} \, dy = \frac{n}{\theta^n} \left[\frac{y^{n+2}}{n+2} \right]_0^\theta = \frac{n}{n+2} \theta^2.$$

Thus,

$$\begin{aligned} \text{MSE}(\hat{\theta}_{\text{MLE}}) &= \text{Bias}^2(\hat{\theta}_{\text{MLE}}) + \text{Var}(\hat{\theta}_{\text{MLE}}) \\ &= \left[\frac{n}{n+1} \theta - \theta \right]^2 + \left[\frac{n}{n+2} \theta^2 - \left(\frac{n}{n+1} \theta \right)^2 \right] \\ &= \frac{2\theta^2}{(n+1)(n+2)} \\ &= O\left(\frac{1}{n^2}\right). \end{aligned}$$

Now let us examine $\hat{\theta}_{\text{MOM}} = 2\bar{X}$. For the bias:

$$\mathbb{E}[2\bar{X}] = 2\mathbb{E}[\bar{X}] = 2\mathbb{E}[X_i] = 2 \cdot \frac{\theta}{2} = \theta,$$

so it is unbiased. For the variance:

$$\text{Var}(2\bar{X}) = 4 \cdot \text{Var}(\bar{X}) = 4 \cdot \frac{\text{Var}(X_i)}{n} = \frac{4}{n} \left(\frac{(b-a)^2}{12} \right) = \frac{4}{n} \left(\frac{(\theta-0)^2}{12} \right) = \frac{\theta^2}{3n} = O\left(\frac{1}{n}\right).$$

Thus, the MSE shrinks faster for the MLE than for the MOM estimator, as expected.

The next example is just for fun.

Example 6.4 Suppose $X_1, \dots, X_n \stackrel{\text{iid}}{\sim} f_X(x)$ with continuous cumulative distribution function $F_X(x)$. Now consider an additional observation X_{n+1} . What is $\mathbb{P}(X_{n+1} \geq \max\{X_1, \dots, X_n\})$?

Let $Y = \max\{X_1, \dots, X_n\}$. We have $f_X(x)$ and $f_Y(y) = n \cdot F_X(y)^{n-1} \cdot f_X(y)$. Then, using Fubini's theorem,

$$\begin{aligned}
 \mathbb{P}(X \geq Y) &= \iint_{x \geq y} f_{X,Y}(x, y) \, dx \, dy \\
 &= \iint_{x \geq y} f_X(x) \cdot n F_X(y)^{n-1} f_X(y) \, dx \, dy \\
 &= \int_x f_X(x) \left[\int_{-\infty}^x n \cdot F_X(y)^{n-1} \cdot f_X(y) \, dy \right] dx \\
 &= \int_x f_X(x) [F_X(y)^n]_{-\infty}^x dx \\
 &= \int_{-\infty}^{\infty} f_X(x) \cdot F_X(x)^n dx \\
 &= \left[\frac{F_X(x)^{n+1}}{n+1} \right]_{-\infty}^{\infty} \\
 &= \frac{1}{n+1}.
 \end{aligned}$$

Further Insight (2026) There is no need for any integration here. Since $X_1, \dots, X_{n+1}$ are independent and identically distributed and their common distribution is continuous, the probability of obtaining any ordering (or equivalently, any permutation) of them is identical to that of obtaining any other ordering. There are $n!$ permutations of $X_1, \dots, X_n$, and $(n+1)!$ if X_{n+1} is added in. With X_{n+1} fixed and $X_1, \dots, X_n$ free, the desired probability is

$$\frac{n!}{(n+1)!} = \frac{1}{n+1}.$$

Lecture 7 — Tuesday, January 28, 2014

2. Asymptotic Theory

Asymptotics and Moment Generating Functions

Previously, we explored distribution theory through the direct technique. This approach involves determining the density function f_X and cumulative distribution function F_X , then performing integration and differentiation as needed. However, this method often proves intractable for complex problems.

Definition 7.1 We say that X_n **converges in distribution** to X and write $X_n \xrightarrow{D} X$ if $F_{X_n}(x) \rightarrow F_X(x)$ for all continuity points x of F as $n \rightarrow \infty$.

Definition 7.2 We say that X_n **converges in probability** to $c \in \mathbb{R}$ and write $X_n \xrightarrow{P} c$ if for all $\varepsilon > 0$, we have $\mathbb{P}(|X_n - c| \geq \varepsilon) \rightarrow 0$ as $n \rightarrow \infty$.

Definition 7.3 The **moment generating function (mgf)** of a random variable X is

$$\text{mgf}_X(t) := \mathbb{E}[e^{tX}].$$

We can verify that

$$\left. \frac{d}{dt} \text{mgf}_X(t) \right|_{t=0} = \mathbb{E} \left[\left. \frac{d}{dt} e^{tX} \right|_{t=0} \right] = \mathbb{E}[e^{tX} \cdot X]_{t=0} = \mathbb{E}[X],$$

under appropriate regularity conditions. This gives the first moment. Similarly, the second moment is given by

$$\left. \frac{d^2}{dt^2} \text{mgf}_X(t) \right|_{t=0} = \mathbb{E}[X^2].$$

A common sufficient condition for $X_n \xrightarrow{D} X$ is that $\text{mgf}_{X_n}(t) \rightarrow \text{mgf}_X(t)$ on a neighborhood of 0, which we will soon use in our heuristic proof of the central limit theorem.

Proposition 7.4 Suppose $Y = X_1 + \cdots + X_n$, where the random variables X_i are independent. Then

$$\text{mgf}_Y(t) = \mathbb{E}[e^{tY}] = \mathbb{E}[e^{t(X_1 + \cdots + X_n)}] = \mathbb{E}[e^{tX_1} e^{tX_2} \cdots e^{tX_n}] = \prod_{i=1}^n \mathbb{E}[e^{tX_i}] = \prod_{i=1}^n \text{mgf}_{X_i}(t).$$

Proposition 7.4 is powerful because it makes no assumption on the distribution of each X_i . The X_i may have completely different distributions, yet we can still compute the moment generating function of their sum through simple multiplication.

Proposition 7.5 If $Y = aX + b$, then

$$\text{mgf}_Y(t) = \mathbb{E}[e^{t(aX+b)}] = \mathbb{E}[e^{atX} \cdot e^{bt}] = e^{bt} \cdot \text{mgf}_X(at).$$

Example 7.6 Let $Z \sim N(0, 1)$ with $f_Z(z) = \frac{1}{\sqrt{2\pi}} e^{-z^2/2}$, the standard bell curve. The mfg of Z is

$$\begin{aligned} \text{mgf}_Z(t) &= \mathbb{E}[e^{tZ}] \\ &= \int_{-\infty}^{\infty} e^{tz} \frac{1}{\sqrt{2\pi}} e^{-z^2/2} dz \\ &= \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}} e^{-(z^2 - 2tz + t^2)/2} e^{t^2/2} dz \\ &= e^{t^2/2} \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}} e^{-(z-t)^2/2} dz \\ &= e^{t^2/2} \end{aligned}$$

because the last integral is the density function of the $N(t, 1)$ distribution, and hence integrates to 1.

Example 7.7 Suppose $X \sim N(\mu, \sigma^2)$. Let $\frac{X-\mu}{\sigma} =: Z \sim N(0, 1)$, so $X = \sigma Z + \mu$. By Proposition 7.5,

$$\text{mgf}_X(t) = e^{\mu t} e^{\sigma^2 t^2/2} = e^{\mu t + \frac{\sigma^2 t^2}{2}}.$$

Example 7.8 Suppose $X_1, \dots, X_n$ are independent and identically distributed as $N(\mu, \sigma^2)$. Let $S_n := X_1 + \dots + X_n$ and $\bar{X}_n = \frac{S_n}{n}$. Then by [Proposition 7.4](#), we have

$$\text{mgf}_{S_n}(t) = \prod_{i=1}^n \text{mgf}_{X_i}(t) = e^{n\mu t + \frac{n\sigma^2 t^2}{2}},$$

which shows that $S_n \sim N(n\mu, n\sigma^2)$. Similarly,

$$\text{mgf}_{\bar{X}}(t) = \text{mgf}_{S_n}\left(\frac{t}{n}\right) = e^{\mu t + \frac{\sigma^2 t^2}{2n}},$$

which implies that

$$\bar{X} \sim N\left(\mu, \frac{\sigma^2}{n}\right).$$

Theorem 7.9 (Central limit theorem) Let $X_1, \dots, X_n$ be independent and identically distributed random variables from some distribution with $\mathbb{E}[X_i] = \mu$ and $\text{Var}(X_i) = \sigma^2 < \infty$. Define $S_n = X_1 + \dots + X_n$. Then

$$\frac{S_n - n\mu}{\sigma\sqrt{n}} \xrightarrow{D} N(0, 1).$$

Although the central limit theorem is stated for S_n , the analogous result for $\bar{X}_n$ follows immediately:

$$\frac{\bar{X}_n - \mu}{\sigma/\sqrt{n}} \rightarrow N(0, 1).$$

The central limit theorem therefore resolves the asymptotic distribution of the sample mean $\bar{X}$ and related estimators $\bar{\theta} = g(X_1, \dots, X_n)$.

Heuristic proof of the central limit theorem. We can write the left-hand side as

$$\frac{1}{\sqrt{n}} \left[\frac{X_1 - \mu}{\sigma} + \dots + \frac{X_n - \mu}{\sigma} \right] =: \frac{1}{\sqrt{n}} [X_1^* + \dots + X_n^*].$$

Note that $\mathbb{E}[X_i^*] = 0$ and $\text{Var}(X_i^*) = 1$. Therefore it suffices to prove $\frac{S_n}{\sqrt{n}} \xrightarrow{D} N(0, 1)$ under the assumption that $\mu = 0$ and $\sigma^2 = 1$. Suppose $m(t)$ is the moment-generating function of X_i for any i , so that $\text{mgf}_{S_n}(t) = [m(t)]^n$. Then $\text{mgf}_{\frac{S_n}{\sqrt{n}}}(t) = [m\left(\frac{t}{\sqrt{n}}\right)]^n$. As $n \rightarrow \infty$, we have $\frac{t}{\sqrt{n}} \rightarrow 0$.

We expand $m(\frac{t}{\sqrt{n}})$ about 0 using a second-order Taylor expansion:

$$m\left(\frac{t}{\sqrt{n}}\right) \approx m(0) + m'(0)\frac{t}{\sqrt{n}} + m''(0)\frac{t^2}{2n}.$$

Observe that $m'(0) = \mathbb{E}[X_i^*] = 0$,

$$m''(0) = \mathbb{E}[X^2] = \text{Var}(X_i^*) + \mathbb{E}^2[X_i^*] = 1 + 0,$$

and

$$m(0) = \mathbb{E}\left[e^{tX^*}\right]\Big|_{t=0} = 1.$$

Therefore (heuristically, though not entirely rigorously), we have

$$\left[m\left(\frac{t}{\sqrt{n}}\right)\right]^n \approx \left[1 + \frac{t^2}{2n}\right]^n \rightarrow e^{t^2/2},$$

which is the moment generating function of $N(0, 1)$. Since convergence of moment generating functions implies convergence in distribution, we are done. $\square$

The Weak Law of Large Numbers

We now return to the notion of convergence in probability.

Proposition 7.10 (Chebyshev's inequality) Let X be a random variable with $\mathbb{E}[X] = \mu$ and $\text{Var}(X) = \sigma^2 < \infty$. Then for all $\varepsilon > 0$, we have

$$\mathbb{P}(|X - \mu| \geq \varepsilon) \leq \frac{\sigma^2}{\varepsilon^2}.$$

Proof. We have

$$\begin{aligned} \sigma^2 = \text{Var}(X) &= \int_{-\infty}^{\infty} (x - \mu)^2 f(x) \, dx \\ &= \int_{-\infty}^{\mu - \varepsilon} (x - \mu)^2 f(x) \, dx + \int_{\mu - \varepsilon}^{\mu + \varepsilon} (x - \mu)^2 f(x) \, dx + \int_{\mu + \varepsilon}^{\infty} (x - \mu)^2 f(x) \, dx \\ &\geq \int_{-\infty}^{\mu - \varepsilon} \varepsilon^2 f(x) \, dx + \int_{\mu + \varepsilon}^{\infty} \varepsilon^2 f(x) \, dx \end{aligned}$$

$$= \varepsilon^2 \mathbb{P}(|X - \mu| \geq \varepsilon).$$

Rearranging yields the claimed inequality. $\square$

Theorem 7.11 (Weak law of large numbers) Let $X_1, \dots, X_n$ be independent and identically distributed random variables from some distribution with $\mathbb{E}[X_i] = \mu$ and $\text{Var}(X_i) = \sigma^2 < \infty$. Then $\bar{X}_n \xrightarrow{P} \mu$.

Proof. Applying Chebyshev's inequality to $\bar{X}$, we have for all $\varepsilon > 0$ that

$$\mathbb{P}(|\bar{X} - \mu| \geq \varepsilon) \leq \frac{\text{Var}(\bar{X})}{\varepsilon^2} = \frac{\sigma^2}{n\varepsilon^2} \rightarrow 0$$

as $n \rightarrow \infty$. Since $\mu = \mathbb{E}[X_i] = \mathbb{E}[\bar{X}]$, we conclude that $\bar{X} \xrightarrow{P} \mu$. $\square$

How does the central limit theorem ([Theorem 7.9](#)) compare to the weak law of large numbers ([Theorem 7.11](#))? First, recall that technical considerations notwithstanding, a random variable is constant if and only if its variance is zero.

Now, without loss of generality, assume $\mathbb{E}[X_i] = 0$ and $\text{Var}(X_i) = 1$. [Theorem 7.11](#) states that

$$\bar{X}_n \xrightarrow{P} 0,$$

while [Theorem 7.9](#) states that

$$\frac{\bar{X} - 0}{1/\sqrt{n}} \xrightarrow{D} N(0, 1),$$

which is equivalent to $\sqrt{n}\bar{X}_n \xrightarrow{D} N(0, 1)$.

Note that $\text{Var}(\bar{X}_n) = \frac{1}{n} \rightarrow 0$, so $\bar{X}_n$ essentially becomes non-random as $n \rightarrow \infty$. In contrast,

$$\text{Var}(\sqrt{n}\bar{X}_n) = (\sqrt{n})^2 \cdot \text{Var}(\bar{X}_n) = n \cdot \frac{1}{n} = 1,$$

which is constant for all n . Therefore $\sqrt{n}\bar{X}_n$ remains a **proper** random variable for all sample sizes.

Lecture 8 — Thursday, January 30, 2014

3. Interval Estimation and Hypothesis Testing

Confidence Intervals and Pivotal Quantities

In the canonical model, we assume $X_1, \dots, X_n \stackrel{iid}{\sim} N(\mu, \sigma^2)$ with known σ^2 (an unrealistic but instructive starting point). From [Example 3.3](#), $\bar{X}$ is both the MLE and MOM for μ , with $\mathbb{E}[\bar{X}] = \mu$ and $\text{Var}(\bar{X}) = \frac{\sigma^2}{n}$. Standardizing gives

$$\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \sim N(0, 1).$$

Hence

$$\mathbb{P}\left(-1.96 \leq \frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \leq 1.96\right) \approx 95\%,$$

which is equivalent to

$$\mathbb{P}\left(\bar{X} - \frac{1.96\sigma}{\sqrt{n}} \leq \mu \leq \bar{X} + \frac{1.96\sigma}{\sqrt{n}}\right) \approx 95\%.$$

Thus a 95% **confidence interval** for μ is

$$\left(\bar{X} - \frac{1.96\sigma}{\sqrt{n}}, \bar{X} + \frac{1.96\sigma}{\sqrt{n}}\right).$$

A **pivotal quantity** q is a function that (1) depends only on the data $(X_1, \dots, X_n)$, the parameter of interest, and known constants, and (2) has a fully specified distribution. For example,

$$\frac{\bar{X} - \mu}{\sigma/\sqrt{n}}$$

is pivotal because $\bar{X}$ comes from the data, σ is known, and the distribution is $N(0, 1)$.

Knowing the distribution of a pivotal quantity allows us to write

$$\mathbb{P}(L_\alpha \leq q(X_1, \dots, X_n; \theta) \leq U_\alpha) = 1 - \alpha$$

for a given small α , since knowing the distribution of $q(X_1, \dots, X_n; \theta)$ means that we can determine the endpoints L_α and U_α . In general, the endpoints are not unique, since the only requirement is that they cover $1 - \alpha$ of the total mass of the distribution.

For the canonical case, the $(1 - \alpha) \cdot 100\%$ confidence interval for μ is

$$\left(\bar{X} - \zeta_\alpha \frac{\sigma}{\sqrt{n}}, \bar{X} + \zeta_\alpha \frac{\sigma}{\sqrt{n}} \right)$$

where ζ_α satisfies $\mathbb{P}(N(0, 1) \leq \zeta_\alpha) = 1 - \alpha$.

Example 8.1 Consider the density function $f_X(x) = \frac{2(\theta-x)}{\theta^2}$ for $x \in (0, \theta)$. First, we verify that $Y = q(X; \theta) = X/\theta$ is a pivotal quantity. Condition (1) is obviously satisfied. For condition (2), we apply some distribution theory. The cdf of Y is

$$F_Y(y) = \mathbb{P}(Y \leq y) = \mathbb{P}\left(\frac{X}{\theta} \leq y\right) = \mathbb{P}(X \leq \theta y) = \int_0^{\theta y} \frac{2(\theta-x)}{\theta^2} dx = \left[\frac{2x}{\theta} - \frac{x^2}{\theta^2} \right]_0^{\theta y} = 2y - y^2.$$

This is completely known and functionally independent of the parameter θ , which confirms (2).

Next, we construct a confidence interval for θ based on X . We have

$$f_Y(y) = \frac{d}{dy} F_Y(y) = 2 - 2y.$$




Figure 3: Equal-tail cutoff points L_α and U_α for the pivotal density $f_Y(y) = 2 - 2y$ on $0 \leq y \leq 1$.

For an **equal-tail confidence interval**, we want L_α, U_α such that $F_Y(L_\alpha) = \alpha/2$ and

$1 - F_Y(U_\alpha) = \alpha/2$. Solving $2y - y^2 = A$ gives

$$y = \frac{2 \pm \sqrt{4 - 4A}}{2} = 1 \pm \sqrt{1 - A}.$$

Since $y \leq 1$, we take $y = 1 - \sqrt{1 - A}$. Therefore

$$L_\alpha = 1 - \sqrt{1 - \alpha/2}, \quad U_\alpha = 1 - \sqrt{\alpha/2}.$$

Now we use the pivotal quantity:

$$\mathbb{P}\left(1 - \sqrt{1 - \alpha/2} \leq \frac{X}{\theta} \leq 1 - \sqrt{\alpha/2}\right) = 1 - \alpha,$$

which implies

$$\mathbb{P}\left(\frac{X}{1 - \sqrt{\alpha/2}} \leq \theta \leq \frac{X}{1 - \sqrt{1 - \alpha/2}}\right) = 1 - \alpha.$$

Therefore, a $(1 - \alpha) \cdot 100\%$ confidence interval for θ is

$$\left(\frac{X}{1 - \sqrt{\alpha/2}}, \frac{X}{1 - \sqrt{1 - \alpha/2}}\right).$$

For a concrete example, suppose we want a 28% confidence interval with $X = 1$ and $\alpha = 0.72$. Then a 28% confidence interval based on $X = 1$ is

$$\left(\frac{1}{0.4}, \frac{1}{0.2}\right) = (2.5, 5).$$

Observe that if $\alpha = 0$, we obtain a 100% confidence interval; but substituting this value yields a confidence interval of (X, ∞) , which is useless in practice.

Fisher Information and the Cramér-Rao Bound

We now present an important fact about maximum likelihood estimators.

Theorem 8.2 (Asymptotic normality of the MLE) Let $\hat{\theta}_n$ denote the maximum likelihood estimator of θ based on an independent and identically distributed sample of size n , where θ represents the unknown parameter. Under suitable regularity conditions, we have

$$\sqrt{n \cdot I(\theta)}(\hat{\theta}_n - \theta) \xrightarrow{D} N(0, 1),$$

where

$$I(\theta) = -\mathbb{E} \left[\frac{\partial^2}{\partial \theta^2} \log(f(X; \theta)) \right]$$

is the **Fisher information**.

This is a highly abstract result, so we illustrate it with a simple example.

Example 8.3 Suppose $X \sim N(\mu, \sigma^2)$ with density

$$f(x) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(x-\mu)^2}{2\sigma^2}}.$$

Then

$$\log(f(x)) = \log \frac{1}{\sqrt{2\pi}} - \log \sigma - \frac{(x-\mu)^2}{2\sigma^2},$$

and

$$\frac{\partial}{\partial \mu} \log(f(x)) = \frac{x-\mu}{\sigma^2}, \quad \frac{\partial^2}{\partial \mu^2} \log(f(x)) = -\frac{1}{\sigma^2}.$$

[Theorem 8.2](#) says that $\hat{\theta}_n = \bar{X}$, $\theta = \mu$, and

$$I(\theta) = \frac{1}{\sigma^2}.$$

Therefore,

$$\sqrt{n \cdot \frac{1}{\sigma^2}}(\bar{X} - \mu) \xrightarrow{D} N(0, 1).$$

Observe that

$$\sqrt{n \cdot \frac{1}{\sigma^2}}(\bar{X} - \mu) = \frac{\bar{X} - \mu}{\sigma/\sqrt{n}},$$

so the convergence is in fact exact. The distribution does not merely converge; it is already $N(0, 1)$ for all n .

We note several important consequences of [Theorem 8.2](#). First, for large n ,

$$\mathbb{E} \left[\sqrt{n \cdot I(\theta)} (\hat{\theta}_n - \theta) \right] \approx 0,$$

which implies

$$\mathbb{E} [\hat{\theta}_n] \approx \theta.$$

Hence maximum likelihood estimators are asymptotically unbiased. In fact, [Theorem 8.2](#) remains valid if we replace $I(\theta)$ with $I(\hat{\theta}_n)$, but this is a deep (and modern) result.

Second, for large n ,

$$\text{Var}[\sqrt{n \cdot I(\theta)} (\hat{\theta}_n - \theta)] \approx 1,$$

which implies

$$n \cdot I(\theta) \cdot \text{Var}(\hat{\theta}_n) \approx 1,$$

and therefore

$$\text{Var}(\hat{\theta}_n) \approx \frac{1}{n \cdot I(\theta)}.$$

This quantity is called the **asymptotic variance**.

Third,

$$\sqrt{n \cdot I(\theta)} (\hat{\theta}_n - \theta)$$

is an approximate pivotal quantity. This allows us to construct an approximate confidence interval for θ of the form

$$\hat{\theta}_n \pm \zeta_\alpha \frac{1}{\sqrt{n \cdot I(\theta)}},$$

where ζ_α is determined from the $N(0, 1)$ distribution.

Finally, the **Cramér-Rao bound** states that if $\hat{\theta}_n$ is unbiased for θ , then

$$\text{Var}(\hat{\theta}_n) \geq \frac{1}{n \cdot I(\theta)}.$$

That is, the maximum likelihood estimator achieves this lower bound asymptotically, which explains why maximum likelihood estimation is so widely used in practice.

Lecture 9 — Tuesday, February 04, 2014

Sample Size Calculations

Example 9.1 Suppose $X \sim \text{Binomial}(n, p)$. For instance, X might represent the number of “yes” answers to a certain yes-or-no survey question. We want to estimate p with a high degree of accuracy. How large should n be? More precisely, what value of n ensures that $\mathbb{P}(|\hat{p} - p| < 0.01) = 99\%$?

We know that $\hat{p} = \frac{X}{n}$ is both the maximum likelihood estimator and the method of moments estimator for p . We require

$$\mathbb{P}\left(\left|\frac{X}{n} - p\right| < 0.01\right) = 99\%,$$

which implies

$$\mathbb{P}(-0.01n < X - np < 0.01n) = 99\%.$$

We can apply [Theorem 7.9](#) to X , yielding

$$\mathbb{P}\left(\frac{-0.01n}{\sqrt{np(1-p)}} < \frac{X - np}{\sqrt{np(1-p)}} < \frac{0.01n}{\sqrt{np(1-p)}}\right) = 99\%.$$

[Theorem 7.9](#) tells us that the middle term converges in distribution to $N(0, 1)$.




Figure 4: Central 99% area under the standard normal curve used in the sample-size calculation.

From standard normal distribution tables or statistical software, we find that we require

$$\frac{0.01n}{\sqrt{np(1-p)}} \geq 2.576.$$

This expression still depends on p , but we can use a conservative bound in practice. Since $p(1-p)$ is a quadratic function with maximum value $1/4$ at $p = 1/2$, we substitute this value to obtain

$$0.01\sqrt{n} = (2.576)(0.5),$$

which yields

$$n \approx 16,590.$$

This is surprisingly large. As a point of comparison, political polls typically aim for 97% confidence, which requires a much smaller sample size.

Hypothesis Testing

Consider a situation where we have two **hypotheses** for a parameter: $H_0 : \theta = \theta_0$ (the **null hypothesis**) and $H_A : \theta = \theta_A$ (the **alternative hypothesis**).

Example 9.2 Suppose $X \sim \text{Binomial}(n, p)$ with hypotheses $H_0 : p = 0.5$ and $H_A : p = 0.9$. Intuitively, we would reject H_0 if $X \geq c$ for some threshold c (to be determined), and accept H_0 if $X < c$. For concreteness, suppose $n = 10$ and consider $c = 7$.

Before proceeding, we need to establish some definitions.

A **type I error** occurs when we reject H_0 while H_0 is actually true. A **type II error** occurs when we accept H_0 while H_A is actually true. The **significance level** is defined as $\mathbb{P}(\text{type I error})$, and the **power** is defined as $1 - \mathbb{P}(\text{type II error})$.

Example 9.3 Returning to [Example 9.2](#), the significance level equals

$$\mathbb{P}(\text{type I error}) = \mathbb{P}(X \geq 7 \mid H_0) = \sum_{x=7}^{10} \binom{10}{x} 0.5^x (1-0.5)^{10-x} = 0.172.$$

Similarly, the power equals

$$1 - \mathbb{P}(\text{type II error}) = 1 - \mathbb{P}(X < 7 \mid H_A) = 1 - \sum_{x=0}^6 \binom{10}{x} (0.9)^x (1-0.9)^{10-x} = 1 - 0.013 = 0.987.$$

Observe that a hypothesis test is fully characterized by its **rejection region** R , the subset of the sample space for which an observation of data in R would lead us to reject H_0 in favour of H_A .

Likelihood Ratio Tests and the Neyman-Pearson Lemma

The **likelihood ratio test** provides a systematic approach to hypothesis testing. We reject the null hypothesis H_0 if

$$\Lambda = \frac{L(\theta_A)}{L(\theta_0)} > c_\alpha$$

for a suitable critical value c_α . We choose c_α so that the significance level is a predetermined value α (often $\alpha = 0.05$). Note that the **likelihood ratio** Λ is also written as $\frac{f_A(x)}{f_0(x)}$.

Theorem 9.4 (Neyman-Pearson lemma) Among all tests with significance level α , the likelihood ratio test has the highest power.

Proof. Let R_1 be the rejection region of the likelihood ratio test:

$$R_1 = \left\{ (X_1, \dots, X_n) \mid \frac{f_A(x)}{f_0(x)} > c_\alpha \right\}.$$

Let R_2 be another rejection region with the same significance level, so

$$\mathbb{P}(R_2 \mid H_0) = \mathbb{P}(R_1 \mid H_0) = \alpha,$$

equivalently,

$$\int_{R_2} f_0(x) \, dx = \int_{R_1} f_0(x) \, dx.$$

Define $C_1 = R_1 \setminus R_2$ and $C_2 = R_2 \setminus R_1$. Then

$$\int_{C_2} f_0(x) \, dx = \int_{C_1} f_0(x) \, dx.$$

To prove optimality, it suffices to show

$$\int_{C_1} f_A(x) \, dx \geq \int_{C_2} f_A(x) \, dx,$$

since this is equivalent to $\mathbb{P}(R_1 \mid H_A) \geq \mathbb{P}(R_2 \mid H_A)$. On $C_1 \subseteq R_1$, we have $f_A \geq c_\alpha f_0$. On $C_2 \subseteq R_1^c$, we have $f_A \leq c_\alpha f_0$. Therefore,

$$\int_{C_1} f_A(x) \, dx \geq \int_{C_1} c_\alpha f_0(x) \, dx = \int_{C_2} c_\alpha f_0(x) \, dx \geq \int_{C_2} f_A(x) \, dx.$$

Hence the likelihood ratio test is most powerful among level- α tests. □

Example 9.5 Suppose $X_1, \dots, X_n$ are independent and identically distributed as $N(\mu, \sigma^2)$ with σ^2 known, and consider $H_0 : \mu = \mu_0$ versus $H_A : \mu = \mu_A$. The likelihood ratio is

$$\Lambda = \frac{\prod_{i=1}^n \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(X_i - \mu_A)^2}{2\sigma^2}}}{\prod_{i=1}^n \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(X_i - \mu_0)^2}{2\sigma^2}}}.$$

Rejecting for large Λ is equivalent to requiring

$$\begin{aligned} -\sum_{i=1}^n (X_i - \mu_A)^2 + \sum_{i=1}^n (X_i - \mu_0)^2 &\text{ to be large} \\ \iff 2(\mu_A - \mu_0) \sum_{i=1}^n X_i &\text{ to be large} \\ \iff 2(\mu_A - \mu_0)n\bar{X} &\text{ to be large.} \end{aligned}$$

Therefore, if $\mu_A > \mu_0$, we reject for large $\bar{X}$; if $\mu_A < \mu_0$, we reject for small $\bar{X}$.

Lecture 10 — Thursday, February 06, 2014

Power Analyses

Recall from [Example 9.5](#) that, in the canonical example $N(\mu, \sigma^2)$ with σ^2 known, the likelihood ratio test prescribes the following:

1. If $\mu_A > \mu_0$, reject H_0 when $\bar{X}$ is large.
2. If $\mu_A < \mu_0$, reject H_0 when $\bar{X}$ is small.

We choose thresholds for “large” and “small” to make the significance level equal to α . Without loss of generality, assume $\mu_A > \mu_0$. We wish to choose c_α such that

$$\mathbb{P}(\bar{X} > c_\alpha \mid H_0) = \alpha,$$

where H_0 is the hypothesis that $\mu = \mu_0$. Under H_0 , we have

$$\bar{X} \sim N(\mu_0, \sigma^2/n), \quad \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}} \sim N(0, 1).$$

Therefore,

$$\mathbb{P}\left(\frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}} > \frac{c_\alpha - \mu_0}{\sigma/\sqrt{n}}\right) = \alpha.$$

For instance, if we want $\alpha = 0.05$, then we reject H_0 when

$$\frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}} > 1.645,$$

or equivalently, when

$$\bar{X} > \mu_0 + 1.645 \frac{\sigma}{\sqrt{n}}.$$

The power of the likelihood ratio test equals

$$\mathbb{P}\left(\frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}} > c_\alpha \mid H_A\right) = \mathbb{P}\left(\frac{\bar{X} - \mu_A}{\sigma/\sqrt{n}} > \frac{\mu_0 - \mu_A}{\sigma/\sqrt{n}} + c_\alpha\right).$$

By convention, we denote $\Phi(\cdot)$ as the cumulative distribution function of $N(0, 1)$. Define $\Delta := \mu_A - \mu_0 > 0$. Then the power equals

$$1 - \Phi\left(\frac{\mu_0 - \mu_A}{\sigma/\sqrt{n}} + c_\alpha\right) = 1 - \Phi\left(\frac{-\Delta}{\sigma/\sqrt{n}} + c_\alpha\right) = 1 - \Phi\left(c_\alpha - \frac{\Delta}{\sigma/\sqrt{n}}\right).$$

Observe that if Δ increases, then power increases (because Φ is increasing). If σ increases, power decreases. The ratio

$$\frac{\Delta}{\sigma}$$

is commonly called the **signal-to-noise ratio**. If n increases, power increases; if c_α increases, power decreases. This illustrates the fundamental tradeoff between type I and type II errors: both cannot be reduced simultaneously. This is illustrated in [Figure 5](#), where $A = \mathbb{P}(\text{type I error})$ is the significance level, $B + C = \mathbb{P}(\text{type II error})$, and $A + D$ is the power.




Figure 5: Null and alternative sampling distributions showing the significance level, the type II error, and the power of the test.

Composite Hypotheses

So far we have considered hypotheses of the form $H_0 : \mu = \mu_0$ and $H_A : \mu = \mu_A$. These are called **simple hypotheses**. We now consider the **composite** alternative

$$\begin{cases} H_0 : \mu = \mu_0 \\ H_A : \mu \neq \mu_0 \end{cases}$$

Here, H_A simply states that the null hypothesis is wrong but does not specify a particular alternative value.

For composite hypotheses, we can construct a **generalized likelihood ratio test**. Let

$$\theta_A := \{\text{parameters under } H_A\} = \{\theta \mid \theta \text{ is a parameter under } H_A\},$$

and let

$$\theta_0 := \{\theta \mid \theta \text{ is a parameter under } H_0\}.$$

Then the generalized likelihood ratio statistic is

$$\Lambda = \frac{\sup_{\theta \in \theta_A} L(\theta)}{\sup_{\theta \in \theta_0} L(\theta)}.$$

As before, we reject H_0 if $\Lambda > c_\alpha$.

Example 10.1 Returning to [Example 9.5](#), we have $\Lambda = \frac{L(\hat{\mu}_{\text{MLE}})}{L(\mu_0)}$. We know that $\hat{\mu}_{\text{MLE}} = \bar{X}$. Therefore

$$\begin{aligned}
 \Lambda &= \frac{\frac{1}{\sqrt{2\pi}\sigma} e^{-\sum_{i=1}^n (X_i - \bar{X})^2 / 2\sigma^2}}{\frac{1}{\sqrt{2\pi}\sigma} e^{-\sum_{i=1}^n (X_i - \mu_0)^2 / 2\sigma^2}} \\
 &= \exp \left(\frac{\sum_{i=1}^n (X_i - \mu_0)^2 - \sum_{i=1}^n (X_i - \bar{X})^2}{2\sigma^2} \right) \\
 &= \exp \left(\frac{\sum_{i=1}^n (X_i^2 - 2\mu_0 X_i + \mu_0^2) - \left(\sum_{i=1}^n X_i^2 - 2\bar{X} \sum_{i=1}^n X_i + n\bar{X}^2 \right)}{2\sigma^2} \right) \\
 &= \exp \left(\frac{n(\bar{X}^2 - 2\mu_0 \bar{X} + \mu_0^2)}{2\sigma^2} \right) \\
 &= \exp \left(\frac{n(\bar{X} - \mu_0)^2}{2\sigma^2} \right).
 \end{aligned}$$

We reject H_0 if Λ is large, which is equivalent to rejecting when $2 \log \Lambda$ is large. This occurs when

$$\frac{n(\bar{X} - \mu_0)^2}{\sigma^2}$$

is large, or equivalently when

$$\frac{|\bar{X} - \mu_0|}{\sigma/\sqrt{n}}$$

is large. Thus we reject H_0 if

$$\bar{X} > \mu_0 + c_\alpha \frac{\sigma}{\sqrt{n}} \quad \text{or} \quad \bar{X} < \mu_0 - c_\alpha \frac{\sigma}{\sqrt{n}}.$$

If $\mu_A > \mu_0$ (or $\mu_A < \mu_0$), we have a **one-sided test**. If $\mu_A \neq \mu_0$, we have a **two-sided test**. For $\alpha = 0.05$, we have $c_\alpha = 1.645$ for a one-sided test and $c_\alpha = 1.96$ for a two-sided test.

For a one-sided test, we reject if

$$\frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}} > 1.645,$$

whereas for a two-sided test, we reject if

$$\frac{|\bar{X} - \mu_0|}{\sigma/\sqrt{n}} > 1.96.$$

Interval Estimation versus Hypothesis Testing

For the canonical case where $X_1, \dots, X_n$ are independent and identically distributed as $N(\mu, \sigma^2)$ with σ^2 known, our confidence interval for μ is

$$\left(\bar{X} - c_\alpha \frac{\sigma}{\sqrt{n}}, \bar{X} + c_\alpha \frac{\sigma}{\sqrt{n}} \right).$$

Our two-sided rejection region for testing $\mu = \mu_0$ consists of

$$\bar{X} > \mu_0 + c_\alpha \frac{\sigma}{\sqrt{n}} \quad \text{or} \quad \bar{X} < \mu_0 - c_\alpha \frac{\sigma}{\sqrt{n}},$$

which is equivalent to

$$\mu_0 < \bar{X} - c_\alpha \frac{\sigma}{\sqrt{n}} \quad \text{or} \quad \mu_0 > \bar{X} + c_\alpha \frac{\sigma}{\sqrt{n}}.$$

Hence we reject $H_0 : \mu = \mu_0$ if and only if μ_0 is not in the confidence interval. Equivalently, the complement of our rejection region is precisely our confidence interval. Therefore interval estimation and hypothesis testing are fundamentally equivalent in this situation.

Generalized Likelihood Ratio Tests and Wilks' Theorem

It turns out that the quantity $2 \log \Lambda$ is very important. To motivate the role, recall that if $Z \sim N(0, 1)$, then $Z^2 \sim \chi_{(1)}^2$; and if $Z_1, \dots, Z_n \stackrel{iid}{\sim} N(0, 1)$, then

$$\sum_{i=1}^n Z_i^2 \sim \chi_{(n)}^2.$$

Theorem 10.2 (Wilks' theorem) We have

$$2 \log \Lambda \xrightarrow{D} \chi^2_{(df)},$$

where $df := \dim(H_A) - \dim(H_0)$. Here, $\dim$ represents the number of parameters estimated under H_0 and H_A , respectively.

Example 10.3 For $N(\mu, \sigma^2)$ with σ^2 known, we have

$$2 \log \Lambda = \frac{(\bar{X} - \mu_0)^2}{\sigma^2/n} \sim (N(0, 1))^2 = \chi^2_{(1)}.$$

We have $\dim(H_A) = 1$ and $\dim(H_0) = 0$.

In this case, $2 \log \Lambda \sim \chi^2_{(1)}$ exactly (not just asymptotically). We would reject H_0 if $2 \log \Lambda > c_\alpha$, where c_α is obtained from a $\chi^2_{(1)}$ table. For instance, when $\alpha = 0.05$, we get $c_\alpha = 3.84$. Observe that $3.84 = (1.96)^2$, which is not a coincidence.

Lecture 11 — Tuesday, February 11, 2014

Recall from [Theorem 10.2](#) that $2 \log \Lambda \xrightarrow{D} \chi^2_{(df)}$. This makes

$$2\ell(\hat{\theta}_{\text{MLE}}) - 2\ell(\theta)$$

an approximate pivotal quantity. For a composite two-sided test (where $H_0 : \theta = \theta_0$ and $H_A : \theta \neq \theta_0$), the set

$$\{\theta \mid 2\ell(\hat{\theta}_{\text{MLE}}) - 2\ell(\theta) < c_\alpha\}$$

therefore forms an approximate $(1 - \alpha) \cdot 100\%$ confidence set.

Example 11.1 Suppose $X_1, \dots, X_m$ are independent and identically distributed as $\text{Poisson}(\lambda_1)$, and $Y_1, \dots, Y_n$ are independent and identically distributed as $\text{Poisson}(\lambda_2)$. Consider $H_0 : \lambda_1 = \lambda_2$ versus $H_A : \lambda_1 \neq \lambda_2$.

The likelihood ratio is

$$\Lambda = \frac{\sup_{\lambda_1, \lambda_2} L(\lambda_1, \lambda_2)}{\sup_{\lambda_1 = \lambda_2} L(\lambda_1, \lambda_2)}.$$

For the Poisson distribution,

$$L(\lambda) = \prod_{i=1}^n \frac{e^{-\lambda} \lambda^{Z_i}}{Z_i!} \propto e^{-n\lambda} \lambda^{\sum_{i=1}^n Z_i},$$

where $Z_1, \dots, Z_n \sim \text{Poisson}(\lambda)$.

Then

$$\ell(\lambda) = -n\lambda + \sum_{i=1}^n Z_i \log(\lambda),$$

so

$$\ell'(\lambda) = -n + \frac{\sum_{i=1}^n Z_i}{\lambda} = 0,$$

and therefore

$$\hat{\lambda}_{\text{MLE}} = \bar{Z}.$$

Under H_A , we have $\hat{\lambda}_1 = \bar{X}$ and $\hat{\lambda}_2 = \bar{Y}$. Under H_0 , we obtain the **pooled estimate**

$$\hat{\lambda}_1 = \hat{\lambda}_2 = \frac{\sum_{i=1}^m X_i + \sum_{j=1}^n Y_j}{m+n}.$$

Letting $\hat{\lambda}$ be the common value of $\hat{\lambda}_1$ and $\hat{\lambda}_2$ above, the likelihood ratio test statistic is

$$\Lambda = \frac{\prod_{i=1}^m \frac{e^{-\bar{X}} (\bar{X})^{X_i}}{X_i!} \prod_{j=1}^n \frac{e^{-\bar{Y}} (\bar{Y})^{Y_j}}{Y_j!}}{\prod_{i=1}^m \frac{e^{-\hat{\lambda}} (\hat{\lambda})^{X_i}}{X_i!} \prod_{j=1}^n \frac{e^{-\hat{\lambda}} (\hat{\lambda})^{Y_j}}{Y_j!}} = \frac{e^{-m\bar{X}} (\bar{X})^{m\bar{X}} e^{-n\bar{Y}} (\bar{Y})^{n\bar{Y}}}{e^{-(m+n)\hat{\lambda}} (\hat{\lambda})^{(m+n)\hat{\lambda}}} = \frac{(\bar{X})^{m\bar{X}} (\bar{Y})^{n\bar{Y}}}{\left(\frac{m\bar{X} + n\bar{Y}}{m+n}\right)^{m\bar{X} + n\bar{Y}}}$$

so

$$\log \Lambda = m\bar{X} \log \left(\frac{(m+n)\bar{X}}{m\bar{X} + n\bar{Y}} \right) + n\bar{Y} \log \left(\frac{(m+n)\bar{Y}}{m\bar{X} + n\bar{Y}} \right).$$

and we reject H_0 if $2 \log \Lambda > 3.84$ at a significance level of 0.05.

If we assume $n = m$, we can simplify $\log \Lambda$ to

$$\log \Lambda = n \left[\bar{x} \log \frac{2\bar{x}}{\bar{x} + \bar{y}} + \bar{y} \log \frac{2\bar{y}}{\bar{x} + \bar{y}} \right].$$

Let us show that $\log \Lambda \geq 0$ with equality if and only if $\bar{x} = \bar{y}$ (agree with our intuition). This result does not change if we divide by $n(\bar{x} + \bar{y})$, so we do so for convenience.

We can write

$$p = \frac{\bar{x}}{\bar{x} + \bar{y}}, \quad 1 - p = \frac{\bar{y}}{\bar{x} + \bar{y}}.$$

Then

$$g(p) := \frac{\log \Lambda}{n(\bar{x} + \bar{y})} = p \log(2p) + (1 - p) \log(2(1 - p)).$$

Therefore

$$g'(p) = \log(2p) + 1 - \log(2(1 - p)) - 1 = \log p - \log(1 - p).$$

Thus

$$g'(p) = 0 \implies p = 1 - p \implies p = \frac{1}{2},$$

and $p = 1/2$ means the same thing as $\bar{x} = \bar{y}$. Moreover,

$$g''(p) = \frac{1}{p} + \frac{1}{1 - p} > 0,$$

and hence $g(p)$ is minimized at $p = \frac{1}{2}$, with

$$g\left(\frac{1}{2}\right) = 0.$$

Note that we still need the asymptotic distribution of $2 \log \Lambda$ to decide *how different* $\bar{x}$ and $\bar{y}$ should be to reject H_0 .

P-values

In a hypothesis test, choosing a significance level gives a rejection rule, but it does not indicate how strongly the observed data contradict H_0 . To measure this strength of evidence, we compare the observed test statistic to its null distribution.

Let T be a test statistic, and suppose that larger values of T provide stronger evidence against H_0 . After observing T_{obs} , define the **p -value**

$$p\text{-value} := \mathbb{P}(T \geq T_{\text{obs}} \mid H_0).$$

In words, this is the probability under H_0 of obtaining a value of the test statistic at least as extreme as the one observed. Equivalently, it is the smallest significance level at which H_0 would be rejected.

For the likelihood ratio statistic Λ used above,

$$p\text{-value} = \mathbb{P}(\Lambda \geq \Lambda_{\text{obs}} \mid H_0).$$

Example 11.2 Suppose $X_1, X_2, \dots, X_5$ are independent and identically distributed as $N(\mu, 5)$, so $\bar{X} \sim N(\mu, 1)$. We test $H_0 : \mu = 0$ against $H_A : \mu \neq 0$, and reject when $|\bar{X}| > 1.96$ at significance level 0.05. There is clearly more evidence against H_0 if $|\bar{X}_{\text{obs}}| = 3$ than if $|\bar{X}_{\text{obs}}| = 2$. Therefore, the p -value here is

$$p\text{-value} = \mathbb{P}(|\bar{X}| > |\bar{X}_{\text{obs}}| \mid H_0).$$




Figure 6: Two-sided p -value region $\mathbb{P}(|Z| > 2)$ under the standard normal distribution.




Figure 7: Two-sided p -value region $\mathbb{P}(|Z| > 3)$ under the standard normal distribution.

Lecture 12 — Thursday, February 13, 2014

4. Classical Inference for Normal Models

Student's t -Distribution and the One-Sample t -Test

So far, we have been studying the canonical case where $X_1, \dots, X_n$ are independent and identically distributed as $N(\mu, \sigma^2)$ with σ^2 known. In this setting,

$$\frac{\bar{X} - \mu}{\sigma/\sqrt{n}}$$

is a pivotal quantity. We now consider a more realistic scenario where σ^2 is unknown. In this case, the pivotal quantity for the known-variance normal model is no longer valid.

Let

$$S^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2$$

be the **sample variance**. Then the **one-sample t -test** has test statistic

$$T = \frac{\bar{X} - \mu}{S/\sqrt{n}} \sim t_{(n-1)}.$$

What is this distribution?

Definition 12.1 Suppose $Z \sim N(0, 1)$ and $U \sim \chi_{(n)}^2$ are independent random variables. Then

$$T = \frac{Z}{\sqrt{U/n}}$$

is distributed according to $t_{(n)}$, the **(Student's) t -distribution** with n degrees of freedom.

Theorem 12.2 Let $X_1, \dots, X_n$ be independent and identically distributed as $N(\mu, \sigma^2)$. Then the following hold:

1.

$$\frac{\sum_{i=1}^n (X_i - \bar{X})^2}{\sigma^2} \sim \chi_{(n-1)}^2.$$

2.

$$\bar{X} \quad \text{and} \quad \sum_{i=1}^n (X_i - \bar{X})^2$$

are independent, which we denote as

$$\bar{X} \perp \sum_{i=1}^n (X_i - \bar{X})^2.$$

Given [Theorem 12.2](#), we can write

$$T = \frac{\frac{\bar{X} - \mu}{\sigma/\sqrt{n}}}{S/\sigma}.$$

The numerator has distribution $N(0, 1)$. The denominator equals

$$\frac{S}{\sigma} = \sqrt{\frac{S^2}{\sigma^2}} = \sqrt{\frac{\sum_{i=1}^n (X_i - \bar{X})^2}{(n-1)\sigma^2}} \sim \sqrt{\frac{\chi_{(n-1)}^2}{n-1}},$$

by (1) of [Theorem 12.2](#). Furthermore, the numerator and denominator are independent by (2). Hence

$$T \sim t_{(n-1)}.$$

Heuristic proof of [Theorem 12.2](#). We omit the formal proof of [Theorem 12.2](#) but provide a

heuristic justification. For (1), observe that

$$\sum_{i=1}^n (X_i - \bar{X})^2 = \sum_{i=1}^n (X_i^2 - 2X_i\bar{X} + \bar{X}^2) = \sum_{i=1}^n X_i^2 - 2\bar{X} \sum_{i=1}^n X_i + n\bar{X}^2 = \sum_{i=1}^n X_i^2 - n\bar{X}^2.$$

Thus

$$\frac{\sum_{i=1}^n (X_i - \bar{X})^2}{\sigma^2} = \frac{\sum_{i=1}^n X_i^2 - n\bar{X}^2}{\sigma^2}.$$

If we (heuristically) assume $\mu = 0$, then

$$\frac{X_i}{\sigma} \sim N(0, 1), \quad \frac{\bar{X}}{\sigma/\sqrt{n}} \sim N(0, 1),$$

and

$$\left(\frac{X_i}{\sigma}\right)^2 \sim \chi_{(1)}^2, \quad \left(\frac{\bar{X}}{\sigma/\sqrt{n}}\right)^2 \sim \chi_{(1)}^2.$$

Therefore,

$$\frac{\sum_{i=1}^n X_i^2 - n\bar{X}^2}{\sigma^2} = \sum_{i=1}^n \left(\frac{X_i}{\sigma}\right)^2 - \frac{n\bar{X}^2}{\sigma^2}$$

behaves like $\chi_{(n)}^2 - \chi_{(1)}^2 \sim \chi_{(n-1)}^2$. For (2), observe that

$$\begin{aligned} \text{Cov}(\bar{X}, X_i - \bar{X}) &= \text{Cov}(\bar{X}, X_i) - \text{Cov}(\bar{X}, \bar{X}) \\ &= \text{Cov}\left(\frac{X_1 + \cdots + X_n}{n}, X_i\right) - \text{Var}(\bar{X}) \\ &= \frac{1}{n} \text{Cov}(X_i, X_i) - \frac{\sigma^2}{n} \\ &= \frac{\sigma^2}{n} - \frac{\sigma^2}{n} = 0. \end{aligned}$$

When random variables are jointly normal, zero covariance implies independence (which is not hard to show). Therefore $\bar{X} \perp (X_i - \bar{X})$. $\square$

Let us summarize what we know about the unknown-variance normal model.

1. Recall the identity

$$\sum_{i=1}^n (X_i - \bar{X})^2 = \sum_{i=1}^n X_i^2 - n\bar{X}^2.$$

We have already shown that

$$\begin{aligned} \mathbb{E} \left[\sum_{i=1}^n (X_i - \bar{X})^2 \right] &= \sum_{i=1}^n \mathbb{E} [X_i^2] - n\mathbb{E} [\bar{X}^2] \\ &= n \left[\text{Var}(X_i) + \mathbb{E} [X_i]^2 \right] - n \left[\text{Var}(\bar{X}) + \mathbb{E} [\bar{X}]^2 \right] \\ &= n(\sigma^2 + \mu^2) - n \left(\frac{\sigma^2}{n} + \mu^2 \right) \\ &= (n-1)\sigma^2. \end{aligned}$$

Therefore, an unbiased estimator of σ^2 is

$$\frac{\sum_{i=1}^n (X_i - \bar{X})^2}{n-1},$$

whereas

$$\hat{\sigma}_{\text{MLE}}^2 = \frac{\sum_{i=1}^n (X_i - \bar{X})^2}{n}$$

is slightly biased.

2. The quantity

$$\frac{\bar{X} - \mu}{S/\sqrt{n}}$$

is a new pivotal quantity. Consequently,

$$\left(\bar{X} - c_\alpha \frac{S}{\sqrt{n}}, \bar{X} + c_\alpha \frac{S}{\sqrt{n}} \right)$$

is a confidence interval for μ , where c_α is obtained from a $t_{(n-1)}$ table. For $\alpha = 0.05$, we have $c_\alpha \approx 2$ when n is relatively large, and $c_\alpha \rightarrow 1.96$ as $n \rightarrow \infty$.

3. The one-sample t -test rejects $H_0 : \mu = \mu_0$ if

$$\frac{|\bar{X} - \mu_0|}{S/\sqrt{n}} > c_\alpha.$$

4. The one-sample t -test is equivalent to the likelihood ratio test. Under H_A ,

$$\hat{\mu}_{\text{MLE}} = \bar{X} \quad \text{and} \quad \hat{\sigma}_{\text{MLE}}^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2.$$

Under H_0 ,

$$\tilde{\mu}_{\text{MLE}} = \mu_0 \quad \text{and} \quad \tilde{\sigma}_{\text{MLE}}^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \mu_0)^2 = \hat{\sigma}_0^2.$$

Thus

$$\begin{aligned} \Lambda &= \frac{\prod_{i=1}^n \frac{1}{\sqrt{2\pi}\hat{\sigma}_A} \exp\left\{-\frac{(X_i - \bar{X})^2}{2\hat{\sigma}_A^2}\right\}}{\prod_{i=1}^n \frac{1}{\sqrt{2\pi}\hat{\sigma}_0} \exp\left\{-\frac{(X_i - \mu_0)^2}{2\hat{\sigma}_0^2}\right\}} \\ &= \left(\frac{\hat{\sigma}_0^2}{\hat{\sigma}_A^2}\right)^{n/2} \frac{\exp\left\{-\frac{\sum_{i=1}^n (X_i - \bar{X})^2}{2\hat{\sigma}_A^2}\right\}}{\exp\left\{-\frac{\sum_{i=1}^n (X_i - \mu_0)^2}{2\hat{\sigma}_0^2}\right\}} \\ &= \left(\frac{\hat{\sigma}_0^2}{\hat{\sigma}_A^2}\right)^{n/2}. \end{aligned}$$

Therefore,

$$2 \log \Lambda = 2 \log \left(\frac{\hat{\sigma}_0^2}{\hat{\sigma}_A^2}\right)^{n/2} = n \log \left(\frac{\hat{\sigma}_0^2}{\hat{\sigma}_A^2}\right) = n \log \left[\frac{\sum_{i=1}^n (X_i - \mu_0)^2}{\sum_{i=1}^n (X_i - \bar{X})^2} \right].$$

Also,

$$\begin{aligned}
 \sum_{i=1}^n (X_i - \mu_0)^2 &= \sum_{i=1}^n [(X_i - \bar{X}) + (\bar{X} - \mu_0)]^2 \\
 &= \sum_{i=1}^n (X_i - \bar{X})^2 + \sum_{i=1}^n (\bar{X} - \mu_0)^2 + 2 \sum_{i=1}^n (X_i - \bar{X})(\bar{X} - \mu_0) \\
 &= \sum_{i=1}^n (X_i - \bar{X})^2 + n(\bar{X} - \mu_0)^2,
 \end{aligned}$$

since $\sum_{i=1}^n (X_i - \bar{X}) = 0$ and $\sum_{i=1}^n (\bar{X} - \mu_0)^2 = n(\bar{X} - \mu_0)^2$. Hence

$$2 \log \Lambda = n \log \left[1 + \frac{n(\bar{X} - \mu_0)^2}{\sum_{i=1}^n (X_i - \bar{X})^2} \right].$$

Now recall that

$$T = \frac{\bar{X} - \mu_0}{S/\sqrt{n}} = \frac{\sqrt{n}(\bar{X} - \mu_0)}{\sqrt{\frac{\sum_{i=1}^n (X_i - \bar{X})^2}{n-1}}},$$

so

$$T^2 = \frac{n(\bar{X} - \mu_0)^2}{\frac{\sum_{i=1}^n (X_i - \bar{X})^2}{n-1}}.$$

Hence

$$2 \log \Lambda = n \log \left[1 + \frac{T^2}{n-1} \right].$$

It follows that $2 \log \Lambda$ is too large if and only if T^2 is too large, which holds if and only if $|T|$ is too large. Therefore, the likelihood ratio test for this model is equivalent to the t -test.

We also have

$$2 \log \Lambda \xrightarrow{D} \chi_{(1)}^2.$$

Here $\dim(H_A) = 2$ (that is, μ and σ^2) and $\dim(H_0) = 1$ (that is, σ^2). Finally,

$$n \log \left[1 + \frac{T^2}{n-1} \right] \approx \log \left(1 + \frac{T^2}{n-1} \right)^n \rightarrow \log e^{T^2} = T^2 \xrightarrow{D} \chi_{(1)}^2.$$

Lecture 13 — Tuesday, February 25, 2014

Two-Sample t -Test

Suppose

$$X_1, X_2, \dots, X_n \stackrel{\text{iid}}{\sim} N(\mu_A, \sigma^2) \quad \text{and} \quad Y_1, Y_2, \dots, Y_m \stackrel{\text{iid}}{\sim} N(\mu_B, \sigma^2),$$

where σ^2 is unknown. We are interested in testing whether these samples come from the same population:

$$H_0 : \mu_A = \mu_B \quad \text{versus} \quad H_A : \mu_A \neq \mu_B.$$

In practice, the null hypothesis often receives preferential treatment (analogous to the presumption of innocence). Note that we can equivalently write $H_0 : \mu_A - \mu_B = 0$.

We use an approach similar to the one-sample t -test. Recall that for the one-sample t -test, we first compute the maximum likelihood estimators of μ and σ^2 . We then replace the maximum likelihood estimator of σ^2 with the unbiased estimator

$$S^2 = \frac{\sum_{i=1}^n (X_i - \bar{X})^2}{n-1}.$$

Finally, we observe that

$$\frac{|\bar{X} - \mu|}{S/\sqrt{n}} \sim t_{n-1}.$$

Since

$$\text{Var}(\bar{X}) = \frac{\sigma^2}{n},$$

the standard deviation of $\bar{X}$ is $\frac{\sigma}{\sqrt{n}}$. Replacing σ with S yields the **standard error**

$$\frac{S}{\sqrt{n}}.$$

This final step amounts to using a test statistic of the form

$$\frac{\text{MLE}}{\text{standard error of maximum likelihood estimator}} \sim t_{(\text{degrees of freedom})}.$$

For the **two-sample t -test**, we proceed analogously. First, we construct the likelihood function:

$$L(\mu_A, \mu_B, \sigma^2) = \prod_{i=1}^n \frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{(X_i - \mu_A)^2}{2\sigma^2}\right) \cdot \prod_{j=1}^m \frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{(Y_j - \mu_B)^2}{2\sigma^2}\right).$$

The log-likelihood is

$$\ell(\mu_A, \mu_B, \sigma^2) = -n \log \sigma - \frac{\sum_{i=1}^n (X_i - \mu_A)^2}{2\sigma^2} - m \log \sigma - \frac{\sum_{j=1}^m (Y_j - \mu_B)^2}{2\sigma^2} + \text{constant}.$$

Setting $\frac{\partial \ell}{\partial \mu_A} = 0$ yields

$$\sum_{i=1}^n (X_i - \mu_A) = 0 \implies \hat{\mu}_A = \bar{X}.$$

Similarly,

$$\frac{\partial \ell}{\partial \mu_B} = 0 \implies \sum_{j=1}^m (Y_j - \mu_B) = 0 \implies \hat{\mu}_B = \bar{Y}.$$

Finally,

$$\frac{\partial \ell}{\partial \sigma} = \frac{-n-m}{\sigma} + \frac{\sum_{i=1}^n (X_i - \mu_A)^2}{\sigma^3} + \frac{\sum_{j=1}^m (Y_j - \mu_B)^2}{\sigma^3} = 0,$$

which gives

$$\hat{\sigma}_{\text{MLE}}^2 = \frac{\sum_{i=1}^n (X_i - \bar{X})^2 + \sum_{j=1}^m (Y_j - \bar{Y})^2}{n+m}.$$

Next, we construct an unbiased estimator of σ^2 . Recall that

$$\mathbb{E} \left[\sum_{i=1}^n (X_i - \bar{X})^2 \right] = (n-1)\sigma^2, \quad \mathbb{E} \left[\sum_{j=1}^m (Y_j - \bar{Y})^2 \right] = (m-1)\sigma^2.$$

Therefore, an unbiased estimator of σ^2 is

$$S^2 = \frac{\sum_{i=1}^n (X_i - \bar{X})^2 + \sum_{j=1}^m (Y_j - \bar{Y})^2}{n + m - 2}.$$

Finally, we compute the test statistic. We have $\hat{\mu}_A - \hat{\mu}_B = \bar{X} - \bar{Y}$. Since

$$\text{Var}(\bar{X} - \bar{Y}) = \frac{\sigma^2}{n} + \frac{\sigma^2}{m},$$

the standard deviation is

$$\sigma \sqrt{\frac{1}{n} + \frac{1}{m}},$$

and the standard error is

$$S \sqrt{\frac{1}{n} + \frac{1}{m}}.$$

Therefore

$$\frac{\bar{X} - \bar{Y}}{S \sqrt{\frac{1}{n} + \frac{1}{m}}} \sim t_{(n+m-2)}.$$

In general, the degrees of freedom equal the total sample size minus the number of **structural** (that is, **non-nuisance**) parameters estimated.

For the two-sample t -test, the same equivalence with the likelihood ratio test can be derived analogously to the one-sample case.

Lecture 14 — Tuesday, March 04, 2014

5. Regression and Categorical Data Analysis

Simple Linear Regression

Consider the model $Y_i \stackrel{\text{indep}}{\sim} N(\alpha + \beta X_i, \sigma^2)$ for $i = 1, \dots, n$, where the key hypothesis test of interest is $H_0 : \beta = 0$ versus $H_A : \beta \neq 0$.

We begin by constructing the maximum likelihood estimators for α and β . The likelihood and log-likelihood are

$$L(\alpha, \beta, \sigma^2) = \prod_{i=1}^n \frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{(Y_i - \alpha - \beta X_i)^2}{2\sigma^2}\right)$$

and

$$\ell(\alpha, \beta, \sigma^2) = -n \log(\sigma) - \frac{\sum_{i=1}^n (Y_i - \alpha - \beta X_i)^2}{2\sigma^2} + \text{constant},$$

respectively. Setting $\frac{\partial \ell}{\partial \alpha} = 0$ yields

$$\sum_{i=1}^n (Y_i - \alpha - \beta X_i) = 0 \implies \hat{\alpha}_{\text{MLE}} = \bar{Y} - \hat{\beta} \bar{X}.$$

Setting $\frac{\partial \ell}{\partial \beta} = 0$ yields

$$-\sum_{i=1}^n (Y_i - \alpha - \beta X_i) X_i = 0.$$

After substituting $\alpha = \bar{Y} - \hat{\beta} \bar{X}$, this becomes

$$\sum_{i=1}^n (Y_i - \bar{Y} + \hat{\beta} \bar{X} - \hat{\beta} X_i) X_i = 0,$$

and therefore

$$\hat{\beta}_{\text{MLE}} = \frac{\sum_{i=1}^n (Y_i - \bar{Y}) X_i}{\sum_{i=1}^n (X_i - \bar{X}) X_i}.$$

Remark 14.1 As a useful identity, note that

$$\sum_{i=1}^n (Y_i - \bar{Y}) \bar{X} = 0.$$

Indeed,

$$\bar{X} \sum_{i=1}^n (Y_i - \bar{Y}) = \bar{X} \left(\sum_{i=1}^n Y_i - n\bar{Y} \right) = 0.$$

Similarly,

$$\sum_{i=1}^n (X_i - \bar{X}) \bar{X} = 0.$$

Therefore we can rewrite

$$\hat{\beta}_{\text{MLE}} = \frac{\sum_{i=1}^n (Y_i - \bar{Y}) X_i - \sum_{i=1}^n (Y_i - \bar{Y}) \bar{X}}{\sum_{i=1}^n (X_i - \bar{X}) X_i - \sum_{i=1}^n (X_i - \bar{X}) \bar{X}} = \frac{\sum_{i=1}^n (X_i - \bar{X})(Y_i - \bar{Y})}{\sum_{i=1}^n (X_i - \bar{X})^2}.$$

This alternative form has a nice interpretation: the covariance is

$$\text{Cov}(X, Y) = \mathbb{E}[(X - \mathbb{E}[X])(Y - \mathbb{E}[Y])]$$

and the variance is

$$\text{Var}(X) = \mathbb{E}[(X - \mathbb{E}[X])^2].$$

Essentially, the numerator approximates $\text{Cov}(X, Y)$ while the denominator approximates $\text{Var}(X)$.

Next, setting $\frac{\partial \ell}{\partial \sigma} = 0$ yields

$$\frac{-n}{\sigma} + \frac{\sum_{i=1}^n (Y_i - \hat{\alpha}_{\text{MLE}} - \hat{\beta}_{\text{MLE}} X_i)^2}{\sigma^3} = 0,$$

which gives

$$\hat{\sigma}_{\text{MLE}}^2 = \frac{\sum_{i=1}^n (Y_i - \hat{\alpha}_{\text{MLE}} - \hat{\beta}_{\text{MLE}} X_i)^2}{n}.$$

An unbiased version of this estimator is

$$S^2 = \frac{\sum_{i=1}^n (Y_i - \hat{\alpha} - \hat{\beta} X_i)^2}{n - 2},$$

where the denominator $(n - 2)$ equals the sample size minus the number of parameters estimated from the mean part. The formal proof requires linear algebra, so we omit it here.

Next, we compute the standard error of $\hat{\beta}_{\text{MLE}}$, which requires first finding its variance. We have

$$\mathbb{E}[\hat{\beta}_{\text{MLE}}] = \mathbb{E}\left[\frac{\sum_{i=1}^n (X_i - \bar{X})Y_i}{\sum_{i=1}^n (X_i - \bar{X})^2}\right] = \mathbb{E}\left[\frac{\sum_{i=1}^n (X_i - \bar{X})(\alpha + \beta X_i - \alpha - \beta \bar{X})}{\sum_{i=1}^n (X_i - \bar{X})^2}\right] = \beta,$$

where we condition on $X_1, \dots, X_n$ and use that $\mathbb{E}[Y_i | X_i] = \alpha + \beta X_i$, which implies $\mathbb{E}[\bar{Y} | X_1, \dots, X_n] = \alpha + \beta \bar{X}$.

Hence $\hat{\beta}_{\text{MLE}}$ is unbiased for β . For the variance, write

$$\text{Var}(\hat{\beta}_{\text{MLE}}) = \text{Var}\left[\frac{\sum_{i=1}^n (X_i - \bar{X})Y_i}{\sum_{i=1}^n (X_i - \bar{X})^2}\right] = \text{Var}\left[\sum_{i=1}^n a_i Y_i\right],$$

where

$$a_i := \frac{X_i - \bar{X}}{\sum_{i=1}^n (X_i - \bar{X})^2}.$$

Finally,

$$\begin{aligned} \text{Var}(\hat{\beta}_{\text{MLE}}) &= \sum_{i=1}^n a_i^2 \text{Var}(Y_i) + \sum_{i=1}^n \sum_{\substack{j=1 \\ j \neq i}}^n a_i a_j \cdot 0 = \sum_{i=1}^n a_i^2 \sigma^2 = \sigma^2 \sum_{i=1}^n a_i^2 \\ &= \sigma^2 \sum_{i=1}^n \left[\frac{X_i - \bar{X}}{\sum_{i=1}^n (X_i - \bar{X})^2} \right]^2 = \sigma^2 \frac{\sum_{i=1}^n (X_i - \bar{X})^2}{\left[\sum_{i=1}^n (X_i - \bar{X})^2 \right]^2} = \frac{\sigma^2}{\sum_{i=1}^n (X_i - \bar{X})^2}. \end{aligned}$$

Thus the standard deviation of $\hat{\beta}_{\text{MLE}}$ is therefore

$$\frac{\sigma}{\sqrt{\sum_{i=1}^n (X_i - \bar{X})^2}},$$

and the corresponding standard error is

$$\frac{S}{\sqrt{\sum_{i=1}^n (X_i - \bar{X})^2}},$$

where

$$S^2 = \frac{\sum_{i=1}^n (Y_i - \hat{\alpha} - \hat{\beta}X_i)^2}{n-2}.$$

Finally, to test $\beta = 0$ versus $\beta \neq 0$, we use the t -test based on the statistic

$$\frac{\hat{\beta}_{\text{MLE}}}{S/\sqrt{\sum_{i=1}^n (X_i - \bar{X})^2}} \sim t_{(n-2)}.$$

If the researcher has control over the data collection process, the X_i values should be spread out as much as possible to maximize the denominator

$$\sum_{i=1}^n (X_i - \bar{X})^2,$$

which in turn minimizes $\text{Var}(\hat{\beta}_{\text{MLE}})$.

Lecture 15 — Thursday, March 06, 2014

Fitted Values and Prediction Intervals

Recall from the simple linear regression estimator formulas that

$$\hat{\beta}_{\text{MLE}} = \frac{\sum_{i=1}^n (X_i - \bar{X})(Y_i - \bar{Y})}{\sum_{i=1}^n (X_i - \bar{X})^2} = \frac{\sum_{i=1}^n (X_i - \bar{X})Y_i}{\sum_{i=1}^n (X_i - \bar{X})^2},$$

and

$$\hat{\alpha}_{\text{MLE}} = \bar{Y} - \hat{\beta}_{\text{MLE}}\bar{X}.$$

$\hat{Y}_i = \hat{\alpha} + \hat{\beta}X_i$ denotes the **fitted** (or **predicted**, or **estimated**) **value**, and $R_i = Y_i - \hat{Y}_i$ denotes the **residual**. The key observation is that $\hat{Y}_i$ is a random variable, so we can quantify its uncertainty by computing its variance. At $X = X_0$, we want to compute

$$\text{Var}(\hat{Y}_0) = \text{Var}(\hat{\alpha} + \hat{\beta}X_0) = \text{Var}(\hat{\alpha}) + X_0^2 \text{Var}(\hat{\beta}) + 2X_0 \text{Cov}(\hat{\alpha}, \hat{\beta}).$$

Earlier, in the variance calculation for the slope estimator, we computed

$$\text{Var}(\hat{\beta}) = \frac{\sigma^2}{\sum_{i=1}^n (X_i - \bar{X})^2}.$$

Now we compute the remaining terms. First,

$$\begin{aligned} \text{Var}(\hat{\alpha}) &= \text{Var}(\bar{Y} - \hat{\beta}\bar{X}) \\ &= \text{Var}(\bar{Y}) + \bar{X}^2 \text{Var}(\hat{\beta}) - 2\bar{X} \text{Cov}(\bar{Y}, \hat{\beta}) \\ &= \frac{\sigma^2}{n} + \bar{X}^2 \frac{\sigma^2}{\sum_{i=1}^n (X_i - \bar{X})^2} - 2\bar{X} \text{Cov}(\bar{Y}, \hat{\beta}). \end{aligned}$$

Also,

$$\text{Cov}(\bar{Y}, \hat{\beta}) = \text{Cov}\left(\sum_{i=1}^n \frac{1}{n} Y_i, \sum_{i=1}^n a_i Y_i\right) = \sum_{i=1}^n \frac{1}{n} a_i \text{Var}(Y_i) = \sum_{i=1}^n \frac{1}{n} a_i \sigma^2 = \frac{\sigma^2}{n} \sum_{i=1}^n a_i = 0.$$

Therefore,

$$\text{Var}(\hat{\alpha}) = \sigma^2 \left[\frac{1}{n} + \frac{\bar{X}^2}{\sum_{i=1}^n (X_i - \bar{X})^2} \right].$$

Similarly,

$$\text{Cov}(\hat{\alpha}, \hat{\beta}) = \text{Cov}(\bar{Y} - \hat{\beta}\bar{X}, \hat{\beta}) = \text{Cov}(\bar{Y}, \hat{\beta}) - \bar{X} \text{Cov}(\hat{\beta}, \hat{\beta}) = -\bar{X} \text{Var}(\hat{\beta}) = \frac{-\bar{X}\sigma^2}{\sum_{i=1}^n (X_i - \bar{X})^2}.$$

So

$$\begin{aligned} \text{Var}(\hat{Y}_0) &= \sigma^2 \left[\frac{1}{n} + \frac{\bar{X}^2}{\sum_{i=1}^n (X_i - \bar{X})^2} \right] + X_0^2 \frac{\sigma^2}{\sum_{i=1}^n (X_i - \bar{X})^2} + 2X_0 \frac{-\bar{X}\sigma^2}{\sum_{i=1}^n (X_i - \bar{X})^2} \\ &= \sigma^2 \left[\frac{1}{n} + \frac{\bar{X}^2 + X_0^2 - 2X_0\bar{X}}{\sum_{i=1}^n (X_i - \bar{X})^2} \right] \\ &= \sigma^2 \left[\frac{1}{n} + \frac{(X_0 - \bar{X})^2}{\sum_{i=1}^n (X_i - \bar{X})^2} \right]. \end{aligned}$$

This variance is minimized at $X_0 = \bar{X}$. As X_0 moves away from $\bar{X}$ (the center of the observed X values), $\text{Var}(\hat{Y}_0)$ increases, so prediction becomes less certain.

Observe that $\text{Cov}(\hat{\alpha}, \hat{\beta}) \geq 0$ when $\bar{X} \leq 0$ and $\text{Cov}(\hat{\alpha}, \hat{\beta}) \leq 0$ when $\bar{X} \geq 0$. Since $\bar{Y} = \hat{\alpha} + \hat{\beta}\bar{X}$, the point $(\bar{X}, \bar{Y})$ lies on the fitted line. When $\bar{X} \geq 0$, increasing the slope tends to decrease the intercept; when $\bar{X} \leq 0$, they tend to increase together. In this sense, $\bar{X}$ acts as a pivot.

Example 15.1 If $Y = \alpha X^\beta$, then X and Y have a **power-law** relationship. Taking logarithms gives

$$\log(Y) = \log(\alpha) + \beta \log(X).$$

So we can fit a linear regression of $\log(Y)$ on $\log(X)$, obtain estimates α' and β' , and then back-transform:

$$\hat{\beta} = \beta', \quad \hat{\alpha} = e^{\alpha'}.$$

Lecture 16 — Tuesday, March 11, 2014

Standardized Regression

In the simple linear regression setup, consider the standardized variables

$$U_i = \frac{X_i - \bar{X}}{\sqrt{\frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2}}, \quad V_i = \frac{Y_i - \bar{Y}}{\sqrt{\frac{1}{n-1} \sum_{i=1}^n (Y_i - \bar{Y})^2}}.$$

Then

$$\sum_{i=1}^n U_i = \frac{\sum_{i=1}^n (X_i - \bar{X})}{\sqrt{\frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2}} = 0,$$

which implies $\bar{U} = 0$. Similarly, $\bar{V} = 0$. Furthermore, the sample variance of the U_i satisfies

$$\frac{1}{n-1} \sum_{i=1}^n (U_i - \bar{U})^2 = \frac{1}{n-1} \sum_{i=1}^n U_i^2 = \frac{\sum_{i=1}^n (X_i - \bar{X})^2}{\frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2 \cdot (n-1)} = 1.$$

Likewise,

$$\frac{1}{n-1} \sum_{i=1}^n (V_i - \bar{V})^2 = 1.$$

Thus U_i measures how many sample standard deviation units X_i lies above or below $\bar{X}$.

When we fit the model $V_i \sim N(\tilde{\alpha} + \tilde{\beta}U_i, \sigma^2)$, we obtain

$$\tilde{\alpha} = \bar{V} - \tilde{\beta}\bar{U} = 0 \quad \text{and} \quad \tilde{\beta} = \frac{\sum_{i=1}^n (U_i - \bar{U})(V_i - \bar{V})}{\sum_{i=1}^n (U_i - \bar{U})^2} = \frac{\sum_{i=1}^n U_i V_i}{\sum_{i=1}^n U_i^2}.$$

Observe that

$$\tilde{\beta}^2 = \frac{\left(\sum_{i=1}^n U_i V_i\right)^2}{\left(\sum_{i=1}^n U_i^2\right)^2} \leq \frac{\left(\sum_{i=1}^n U_i^2\right)\left(\sum_{i=1}^n V_i^2\right)}{\left(\sum_{i=1}^n U_i^2\right)^2} = \frac{(n-1)(n-1)}{(n-1)^2} = 1.$$

Therefore $\tilde{\beta} \in [-1, 1]$, which implies that the slope never exceeds 45° . This observation underlies the name **regression to the mean**.

Pearson's Goodness-of-Fit Test

Consider

$$H_0 : p_1 = p_1^0, p_2 = p_2^0, \dots, p_k = p_k^0,$$

with the alternative H_A that at least one equality above fails. The likelihood ratio is

$$\Lambda = \frac{\mathcal{L}(\tilde{p}_1, \dots, \tilde{p}_k)}{\mathcal{L}(p_1^0, \dots, p_k^0)} = \frac{\frac{n!}{X_1! \dots X_k!} \tilde{p}_1^{X_1} \dots \tilde{p}_k^{X_k}}{\frac{n!}{X_1! \dots X_k!} (p_1^0)^{X_1} \dots (p_k^0)^{X_k}}.$$

Hence

$$2 \log \Lambda = 2 \sum_{k=1}^m X_k \log \left(\frac{\tilde{p}_k}{p_k^0} \right).$$

Since $\tilde{p}_k = \frac{X_k}{n}$, we can write

$$O_k := X_k = n\tilde{p}_k, \quad E_k := np_k^0,$$

to denote the observed and expected counts, respectively, so

$$2 \log \Lambda = 2 \sum_{k=1}^m O_k \log \left(\frac{O_k}{E_k} \right).$$

We reject H_0 if $2 \log \Lambda > C_\alpha$, where C_α is obtained from a $\chi^2_{(m-1)}$ reference distribution.

Example 16.1 Consider data on 63 sudden heart attacks:

M	Tu	W	Th	F	Sat	Sun
22	7	6	13	5	4	6

Here H_0 states that heart attacks are equally likely on all seven days, and H_A states otherwise. Under H_0 ,

$$E_k = 63 \cdot \frac{1}{7} = 9, \quad k = 1, \dots, 7.$$

Then

$$2 \log \Lambda = 2 \sum_{k=1}^7 O_k \log \left(\frac{O_k}{E_k} \right) = 2 \left[22 \log \left(\frac{22}{9} \right) + 7 \log \left(\frac{7}{9} \right) + \dots + 6 \log \left(\frac{6}{9} \right) \right] = 23.27.$$

The 5% cutoff from χ^2_6 is 12.59. If H_0 were true, values above 12.59 would occur only about 5% of the time. Since $23.27 > 12.59$, we reject H_0 .

Pearson's test statistic is

$$\sum_{k=1}^m \frac{(O_k - E_k)^2}{E_k} \sim \chi^2_{\text{degrees of freedom}}.$$

To derive the relationship to the likelihood ratio test, let

$$f(x) = x \log \left(\frac{x}{x_0} \right),$$

where x_0 is known. We use a two-term Taylor expansion. Since

$$f'(x) = \log \left(\frac{x}{x_0} \right) + 1, \quad f''(x) = \frac{1}{x},$$

we have

$$\begin{aligned} f(x) &\approx f(x_0) + f'(x_0)(x - x_0) + \frac{f''(x_0)}{2}(x - x_0)^2 \\ &= 0 + 1(x - x_0) + \frac{1}{2x_0}(x - x_0)^2 \end{aligned}$$

$$= x - x_0 + \frac{(x - x_0)^2}{2x_0}.$$

Therefore,

$$2 \log \Lambda \approx 2 \sum_{k=1}^m \left[(O_k - E_k) + \frac{1}{2E_k} (O_k - E_k)^2 \right].$$

Since

$$\sum_{k=1}^m O_k = n$$

and

$$\sum_{k=1}^m E_k = n,$$

we obtain

$$2 \log \Lambda \approx \sum_{k=1}^m \frac{(O_k - E_k)^2}{E_k}.$$

Example 16.2 Recall [Example 4.2](#), with genotype probabilities

$$\mathbb{P}(\text{AA}) = (1 - \theta)^2, \quad \mathbb{P}(\text{Aa}) = 2\theta(1 - \theta), \quad \mathbb{P}(\text{aa}) = \theta^2.$$

Given data with 37 individuals of type AA, 46 of type Aa, and 17 of type aa in a sample of 100, we test whether the Hardy-Weinberg model holds. The null hypothesis is that the population follows the Hardy-Weinberg parametrization. In [Example 4.2](#), we found that

$$\hat{\theta}_{\text{MLE}} = \frac{X_2 + 2X_3}{2n} = \frac{46 + 2 \cdot 17}{2 \cdot 100} = 0.4.$$

This yields expected counts

$$\mathbb{E}_1 = 36, \quad \mathbb{E}_2 = 48, \quad \mathbb{E}_3 = 16,$$

which are close to the observed values. To quantify the fit, we compute

$$2 \log \Lambda = 2 \left[37 \log \left(\frac{37}{36} \right) + 46 \log \left(\frac{46}{48} \right) + 17 \log \left(\frac{17}{16} \right) \right] = 0.1733.$$

The Pearson test statistic equals

$$\frac{(37 - 36)^2}{36} + \frac{(46 - 48)^2}{48} + \frac{(17 - 16)^2}{16} = 0.1736,$$

which is nearly identical. However, we must be careful with the degrees of freedom:

$$\dim(H_A) = 2, \quad \dim(H_0) = 1, \quad df = 2 - 1 = 1.$$

For 1 degree of freedom, the 5% cutoff is 3.84. Since

$$0.1733 < 3.84,$$

we accept the null hypothesis and conclude that the Hardy-Weinberg model provides a good fit.

We note three important observations. First, Pearson's test is a second-order Taylor approximation to the likelihood ratio test. Second, Pearson's test has an intuitive interpretation: it measures the magnitude of

$$\left| \frac{\hat{\theta}_{\text{MLE}} - \theta_0}{\text{stdev}(\hat{\theta}_{\text{MLE}})} \right|.$$

Third, Pearson's statistic is substantially easier to calculate by hand than the likelihood ratio test statistic, which explains its widespread use until the 1960s when computers became widely available.

Lecture 17 — Thursday, March 13, 2014

Applications of the Multinomial Model

We now look at several applications of the multinomial model: hospital quality assessment, authorship signatures in Jane Austen's writing (with analogous applications in music), and comparisons between two distributions.

Editorial Note This portion of the lecture has been lost.




Figure 8: Partition of a continuous density into intervals that induce multinomial cell probabilities.

If we divide a continuous distribution into n intervals, the resulting counts of observations from each interval form a multinomial model. This can help identify which distribution is plausible, provided the required parameters are known or can be estimated.

To make this precise, suppose $X_1, \dots, X_m$ are independent observations and we choose cut points

$$-\infty = a_0 < a_1 < \dots < a_n = \infty.$$

See Figure 8. Define interval counts

$$O_i = \sum_{r=1}^m \mathbf{1}\{a_{i-1} < X_r \leq a_i\}, \quad i = 1, \dots, n.$$

Under a candidate model F_θ , the cell probabilities are

$$p_i(\theta) = \mathbb{P}(a_{i-1} < X \leq a_i) = \int_{a_{i-1}}^{a_i} f_\theta(x) dx = F_\theta(a_i) - F_\theta(a_{i-1}),$$

so that

$$(O_1, \dots, O_n) \sim \text{Multinomial}(m; p_1(\theta), \dots, p_n(\theta)).$$

If θ is unknown, we can estimate it by $\hat{\theta}$, and then compare observed counts O_i with expected counts $E_i = mp_i(\hat{\theta})$. A standard summary is Pearson's statistic

$$X^2 = \sum_{i=1}^n \frac{(O_i - E_i)^2}{E_i},$$

or the likelihood-ratio version

$$G^2 = 2 \sum_{i=1}^n O_i \log \left(\frac{O_i}{E_i} \right).$$

Small values indicate that the observed bin counts are consistent with the candidate distribution; unusually large values indicate that the model allocates mass very differently from the data and is

therefore implausible. In large samples, these statistics are compared with a $\chi^2_{(n-1-k)}$ distribution, where k is the number of estimated parameters.

Another key application is testing independence. For discrete random variables, this is done with **contingency tables** (for example, smoking status versus cancer status). In a table with I rows and J columns, we have $\dim(H_0) = (I - 1) + (J - 1)$ and $\dim(H_A) = IJ - 1$. Hence

$$(IJ - 1) - (I - 1) - (J - 1) = \boxed{(I - 1)(J - 1)}.$$

This degrees-of-freedom formula is common, but it is not universally valid in every modelling setup.

Lecture 18 — Tuesday, March 18, 2014

6. Study Design and Sampling

Stratified Sampling and Paired Analysis

A researcher wishes to choose 10 blocks from 100 possible options. One approach is to select the most “representative” ones, yielding a **judgment sample**. Alternatively, use a random number generator to choose 10 blocks randomly, producing a **simple random sample**. A better approach divides the blocks into two groups called strata, then takes a simple random sample from each stratum (5 from small blocks, 5 from large blocks), yielding a **stratified random sample**. Judgment sampling tends to exhibit bias because human judgment is systematically flawed (psychologists continue to investigate why). The simple random sample reduces variance compared to judgment sampling. The stratified random sample reduces both variance and bias.

Consider a population $Z_1, Z_2, \dots, Z_{2n}$ with mean μ and variance σ^2 , divided into two strata: Stratum 1 contains $X_1, \dots, X_n$ with mean μ_1 and variance σ_1^2 , while Stratum 2 contains $Y_1, \dots, Y_n$ with mean μ_2 and variance σ_2^2 . For a simple random sample, the estimator $\bar{Z}$ satisfies

$$\mathbb{E}[\bar{Z}] = \mu, \quad \text{Var}(\bar{Z}) = \frac{\sigma^2}{2n}.$$

For a stratified sample, the estimator $\frac{\bar{X} + \bar{Y}}{2}$ satisfies

$$\mathbb{E} \left[\frac{\bar{X} + \bar{Y}}{2} \right] = \frac{\mu_1 + \mu_2}{2} = \mu,$$

while

$$\text{Var} \left(\frac{\bar{X} + \bar{Y}}{2} \right) = \frac{1}{4} [\text{Var}(\bar{X}) + \text{Var}(\bar{Y})] = \frac{\sigma_1^2 + \sigma_2^2}{4n},$$

with no covariance term because the two strata are sampled independently. We can have

$$\frac{\sigma_1^2 + \sigma_2^2}{4n} < \frac{\sigma^2}{2n}$$

when strata are internally homogeneous.

Example 18.1 Consider the toy population $\{1, 2, 3, 4\}$ divided into small = $\{1, 2\}$ and large = $\{3, 4\}$. We have

$$\mu = \frac{1 + 2 + 3 + 4}{4} = 2.5, \quad \mu_1 = \frac{1 + 2}{2} = 1.5, \quad \mu_2 = \frac{3 + 4}{2} = 3.5,$$

which satisfies

$$\frac{\mu_1 + \mu_2}{2} = \frac{1.5 + 3.5}{2} = 2.5 = \mu.$$

The variances are

$$\sigma^2 = \frac{(1 - 2.5)^2 + (2 - 2.5)^2 + (3 - 2.5)^2 + (4 - 2.5)^2}{4} = \frac{5}{4} = 1.25,$$

$$\sigma_1^2 = \frac{(1 - 1.5)^2 + (2 - 1.5)^2}{2} = 0.25,$$

$$\sigma_2^2 = \frac{(3 - 3.5)^2 + (4 - 3.5)^2}{2} = 0.25.$$

Therefore,

$$\frac{\sigma_1^2 + \sigma_2^2}{2} = 0.25 < 1.25 = \sigma^2,$$

demonstrating the variance reduction.

Paired Analysis

We now demonstrate mathematically how pairing reduces variance. Suppose

$$X_1, \dots, X_n \sim N(\mu_1, \sigma^2) \quad \text{and} \quad Y_1, \dots, Y_n \sim N(\mu_2, \sigma^2)$$

are independent samples. For an unpaired analysis, the estimator $\bar{X} - \bar{Y}$ has variance

$$\text{Var}(\bar{X} - \bar{Y}) = \text{Var}(\bar{X}) + \text{Var}(\bar{Y}) = \frac{\sigma^2}{n} + \frac{\sigma^2}{n} = \frac{2\sigma^2}{n}.$$

(There is no covariance term.) For a paired analysis, let $D_i = X_i - Y_i$ and estimate $\bar{D}$. We have

$$\mathbb{E}[\bar{D}] = \mathbb{E}[\bar{X} - \bar{Y}] = \mu_1 - \mu_2,$$

so $\bar{D}$ is also unbiased for $\mu_1 - \mu_2$. The variance is

$$\text{Var}(\bar{D}) = \left(\frac{1}{n}\right)^2 \sum_{i=1}^n \text{Var}(D_i),$$

where we note that $\text{Cov}(D_i, D_j) = 0$ for $i \neq j$. Since

$$\text{Var}(D_i) = \text{Var}(X_i - Y_i) = \text{Var}(X_i) + \text{Var}(Y_i) - 2\text{Cov}(X_i, Y_i),$$

and using the correlation

$$\text{Corr}(X_i, Y_i) = \frac{\text{Cov}(X_i, Y_i)}{\sqrt{\text{Var}(X_i)\text{Var}(Y_i)}} = \rho,$$

we obtain

$$\text{Var}(D_i) = \sigma^2 + \sigma^2 - 2\rho\sigma^2 = (2 - 2\rho)\sigma^2.$$

Therefore,

$$\text{Var}(\bar{D}) = \frac{2(1 - \rho)\sigma^2}{n}.$$

When $\rho > 0$ (which is the rationale for pairing), we have

$$\text{Var}(\bar{D}) < \text{Var}(\bar{X} - \bar{Y}).$$

Example 18.2 Suppose the correlation is 50%, that is, $\rho = 0.5$, which represents a mild correlation for “good pairs.” This yields a 50% reduction in variance, which translates to doubling the effective precision.

Lecture 19 — Thursday, March 20, 2014

Graphical Methods

Histograms and **scatter plots** are valuable visualization tools. For histograms, a common rule of thumb is that larger sample sizes permit more bins, but typically no more than $\sqrt{n}$ bins should be used. Box-and-whisker plots are particularly effective for comparing the distributions of several variables.

Good plotting practices are essential. For example, one should always use consistent scales across comparable plots. The principle of *garbage in, garbage out* applies: mathematics cannot remedy poor-quality data. There is no fully automatic method to detect problematic data; careful human inspection of plots and datasets remains necessary.

Empirical Studies

Chapter 3 of the official course notes discusses empirical study methodology. The **PPDAC cycle** describes the five stages of statistical inquiry: *Problem* (define the research question), *Plan* (design the study), *Data* (collect observations), *Analysis* (apply statistical methods), and *Conclusion* (interpret results and communicate findings).

In a study, the **population** is the group about which we wish to make inferences. An **attribute** is a characteristic of the population, such as average age, distribution of household income, or even a functional relationship between two other attributes.

The distinction between the target population and the study population is important. The **target population** is the full group to which we want our conclusions to apply, while the **study population** is the group that is actually accessible for sampling. In an ideal study these coincide, but in practice the study population is often narrower because of geography, timing, eligibility restrictions, or nonresponse. When the two differ, we must be careful about how far the results can be generalized beyond the group that was actually studied.

An **observational study** involves collecting data without controlling or manipulating variables, while an **experimental study** involves actively controlling variables.

Lecture 21 — Tuesday, March 25, 2014

7. Causation and Bayesian Statistics

Causation versus Association

We now turn to the interpretation of statistical relationships.

Causation means that, *all else being equal*, changing X produces changes in Y . These changes are typically probabilistic, representing shifts in the distribution of Y rather than deterministic outcomes. For example, a cancer drug may significantly improve survival rates without guaranteeing recovery for every patient. The key principle in establishing causation is **randomization**, which minimizes violations of the “all else equal” requirement.

Association, in contrast, can arise from three scenarios: X causes Y , or Y causes X , or an unseen **confounding factor** Z causes both X and Y .

An **experimental** (or **randomized**) **study** controls the randomization process. In a **double-blind study**, neither the physician nor the patient knows whether the treatment or placebo is being administered. This matters because subtle cues such as facial expressions can influence patient outcomes.

Blocking is a technique where participants are divided into blocks that control for known potential confounding factors (for example, age groups in a drug study).

In **observational studies**, we cannot randomize. For instance, to study whether smoking causes cancer, we cannot ethically assign people to smoke for 20 years. We can only analyze observational data. Chi-squared tests can assess association, but only association — *and not causation* — can be inferred. Probabilistic outcomes mean that some heavy smokers live long lives, while some healthy non-smokers develop cancer early.

Matching attempts to make comparison groups as similar as possible (for example, when studying gender differences, match participants by health status, weight, and other relevant factors).

Example 21.1 (1973 Berkeley Admission Controversy) In 1973, admission data from UC Berkeley showed that 46% of male applicants were admitted compared to only 30% of female applicants. However, when a Berkeley statistician analyzed the data department by department (a form of matching), the analysis revealed that in most departments, women were slightly favored in admissions. The apparent discrepancy arose because women disproportionately applied to highly competitive departments, while men applied more frequently to less competitive departments. The department selectivity was the confounding factor. This phenomenon is known as **Simpson's paradox**.

Example 21.1 illustrates a fundamental limitation of observational studies: if department-level data had been unavailable, the misleading aggregate pattern would have been impossible to detect. The key lesson — perhaps more important than technical details like the likelihood ratio test — is to always remember these principles when interpreting statistical associations.

Bayesian Statistics

The Bayesian framework treats the parameter θ as a random variable. We denote $\pi(\theta)$ as the **prior distribution** and $\pi(\theta \mid x_1, \dots, x_n)$ as the **posterior distribution**, representing our beliefs before and after observing the data, respectively. The central task in Bayesian statistics is to update from $\pi(\theta)$ to $\pi(\theta \mid x_1, \dots, x_n)$ using Bayes' theorem:

$$\pi(\theta \mid x_1, \dots, x_n) = \frac{f(x_1, \dots, x_n \mid \theta) \cdot \pi(\theta)}{\int f(x_1, \dots, x_n \mid \theta) \cdot \pi(\theta) \, d\theta}.$$

The numerator involves $f(x_1, \dots, x_n \mid \theta)$, which is the likelihood of θ . The denominator is often computationally challenging, especially when θ is a vector with thousands of components, making even numerical evaluation difficult.

Once the posterior is obtained, however, we can compute point estimates (such as $\mathbb{E}[\theta \mid x_1, \dots, x_n]$ or the posterior mode) or **credible regions**. Unlike frequentist confidence intervals, Bayesian credible regions directly represent our uncertainty about θ given the data, since we have the entire posterior distribution.

Example 21.2 Suppose $X \sim \text{Binomial}(n, p)$ and we are interested in the unknown parameter p . We first need to specify a prior distribution. Without prior knowledge about p , a

reasonable choice is the uniform prior $p \sim \text{Uniform}(0, 1)$. Then

$$\pi(p | x) = \frac{f(x | p) \cdot \pi(p)}{\int f(x | p) \cdot \pi(p) \, dp} = \frac{\binom{n}{x} p^x (1-p)^{n-x} \cdot 1}{\int_0^1 \binom{n}{x} p^x (1-p)^{n-x} \cdot 1 \, dp}.$$

To evaluate this expression, we require properties of the beta distribution.

We say that Z has a **beta distribution** and write $Z \sim \text{Beta}(\alpha, \beta)$ if

$$f(z) = \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha)\Gamma(\beta)} z^{\alpha-1} (1-z)^{\beta-1}$$

for $z \in (0, 1)$. The expected value is

$$\begin{aligned} \mathbb{E}[Z] &= \int_0^1 z \cdot f(z) \, dz = \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha)\Gamma(\beta)} \int_0^1 z \cdot z^{\alpha-1} (1-z)^{\beta-1} \, dz \\ &= \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha)\Gamma(\beta)} \underbrace{\int_0^1 z^{(\alpha+1)-1} (1-z)^{\beta-1} \, dz}_{\frac{\Gamma(\alpha+1)\Gamma(\beta)}{\Gamma(\alpha+1+\beta)} \text{ since the Beta}(\alpha+1, \beta) \text{ density integrates to 1}} \\ &= \frac{\alpha}{\alpha + \beta}. \end{aligned}$$

Example 21.3 Returning to the denominator in [Example 21.2](#), which resembles the beta distribution, it is equal to

$$\begin{aligned} &\int_0^1 \binom{n}{x} p^x (1-p)^{n-x} \cdot 1 \, dp \\ &= \frac{\Gamma(x+1) \cdot \Gamma(n-x+1)}{\Gamma(n+2)} \binom{n}{x} \cdot \underbrace{\int_0^1 \frac{\Gamma(n+2)}{\Gamma(x+1) \cdot \Gamma(n-x+1)} p^{(x+1)-1} (1-p)^{(n-x+1)-1} \, dp}_{=1} \\ &= \binom{n}{x} \frac{\Gamma(x+1)\Gamma(n-x+1)}{\Gamma(n+2)} \\ &= \frac{n!}{x!(n-x)!} \cdot \frac{x! \cdot (n-x)!}{(n+1)!} \end{aligned}$$

$$\begin{aligned}
&= \frac{n!}{(n+1)!} \\
&= \frac{1}{n+1}.
\end{aligned}$$

Thus

$$\pi(p | x) = (n+1) \binom{n}{x} p^x (1-p)^{n-x} \cdot 1 = \frac{\Gamma(n+2)}{\Gamma(x+1)\Gamma(n-x+1)} p^{(x+1)-1} (1-p)^{(n-x+1)-1}.$$

We conclude that the posterior distribution is $p | x \sim \text{Beta}(x+1, n-x+1)$.

Lecture 22 — Thursday, March 27, 2014

Conjugate Priors and Jeffrey's Rule

Example 22.1 Continuing from [Example 21.2](#), we saw that when $X \sim \text{Binomial}(n, p)$ and the prior on p is $\text{Uniform}(0, 1)$, the posterior is $\text{Beta}(x+1, n-x+1)$.

The posterior expectation is therefore

$$\mathbb{E}[p | x] = \frac{x+1}{n+2}.$$

By comparison, $\hat{p}_{\text{MLE}} = \frac{x}{n}$. These are close for large n , but can differ noticeably for small n . For one observation (i.e., $n = 1$) we have the following tabulation:

x	0	1
MLE	0	1
Bayes	1/3	2/3

Observe that the $\text{Unif}(0, 1)$ distribution is exactly the $\text{Beta}(1, 1)$ distribution. The prior mean is $\frac{1}{2}$, and the posterior mean $\mathbb{E}[p | x]$ combines information from both the prior mean and the likelihood.

The beta distribution thus is a **conjugate prior** for the binomial family, meaning the

posterior remains a beta distribution. In particular, if the prior is $\text{Beta}(\alpha, \beta)$ so that

$$\pi(p) \propto p^{\alpha-1}(1-p)^{\beta-1},$$

and the model for the data is $\text{Binomial}(n, p)$ so that

$$f(x | p) = \binom{n}{x} p^x (1-p)^{n-x},$$

then

$$\pi(p | x) \propto f(x | p)\pi(p) \propto p^{x+\alpha-1}(1-p)^{n-x+\beta-1},$$

so the posterior is $\text{Beta}(x + \alpha, n - x + \beta)$. The posterior mean is

$$\mathbb{E}[p | x] = \frac{x + \alpha}{n + \alpha + \beta}.$$

Common conjugate pairs include beta-binomial, gamma-Poisson, and normal-normal; in each case, conjugacy is straightforward to check.

How do we choose a prior? A common idea is to choose a “most ignorant” prior that encodes minimal information. For binomial p , the $\text{Beta}(1, 1)$ distribution (or equivalently, the $\text{Unif}(0, 1)$ distribution) is a flat prior. A key issue is that flat priors are generally not invariant under reparametrization.

Example 22.2 Let $\pi_p(p) = 1$ for $p \in (0, 1)$ and set $Q = -\log(p) \in [0, \infty)$. Then

$$F_Q(q) = \mathbb{P}(Q \leq q) = \mathbb{P}(-\log(p) \leq q) = \mathbb{P}(p \geq e^{-q}) = 1 - e^{-q},$$

so

$$\pi_Q(q) = \frac{d}{dq} F_Q(q) = e^{-q}.$$

So the prior on Q is $\text{Exponential}(1)$, which is decidedly not flat. Thus a flat prior is not invariant to a change of scale. More generally, for a monotone transformation $Q = g(p)$,

$$F_Q(q) = F_p(g^{-1}(q)), \quad \pi_Q(q) = \pi_p(g^{-1}(q)) \left| \frac{d}{dq} g^{-1}(q) \right|,$$

which is the usual **change-of-variables formula** for priors.

Jeffreys' rule says that to obtain invariance under reparametrization, choose

$$\pi(\theta) \propto \sqrt{-\mathbb{E} \left[\frac{\partial^2}{\partial \theta^2} l(\theta) \right]}.$$

This quantity is based on the Fisher information. If we reparametrize with $\phi = g(\theta)$ and apply Jeffreys' rule, then

$$\pi(\phi) = \sqrt{-\mathbb{E} \left[\frac{\partial^2}{\partial \phi^2} l(\phi) \right]} = \sqrt{-\mathbb{E} \left[\frac{\partial^2}{\partial \theta^2} l(\theta) \cdot \left(\frac{d\theta}{d\phi} \right)^2 \right]} = \sqrt{-\mathbb{E} \left[\frac{\partial^2}{\partial \theta^2} l(\theta) \right]} \cdot \left| \frac{d\theta}{d\phi} \right| = \pi(\theta) \cdot \left| \frac{d}{d\phi} g^{-1}(\phi) \right|,$$

which matches the change-of-variables formula.

Example 22.3 For $X \sim \text{Binomial}(n, p)$,

$$L(p) = \binom{n}{x} p^x (1-p)^{n-x}, \quad l(p) = \text{const} + x \log(p) + (n-x) \log(1-p),$$

so

$$l'(p) = \frac{x}{p} - \frac{n-x}{1-p}, \quad l''(p) = -\frac{x}{p^2} - \frac{n-x}{(1-p)^2},$$

and therefore

$$-\mathbb{E} [l''(p)] = \frac{\mathbb{E}[X]}{p^2} + \frac{n - \mathbb{E}[X]}{(1-p)^2} = \frac{n}{p(1-p)}.$$

Jeffreys' rule yields

$$\pi(p) \propto \frac{1}{\sqrt{p(1-p)}},$$

so Jeffreys' prior is $\text{Beta}(\frac{1}{2}, \frac{1}{2})$. Under this prior, the posterior mean is

$$\mathbb{E}[p | x] = \frac{x + 1/2}{n + 1},$$

compared with

$$\mathbb{E}[p | x] = \frac{x + 1}{n + 2}$$

under the uniform prior. One interpretation of a $\text{Beta}(\alpha, \beta)$ prior is that it acts like $(\alpha + \beta)$ pseudo-observations, of which α are successes.

Lecture 23 — Tuesday, April 01, 2014

Empirical Bayes

Let $Y_i \stackrel{\text{indep}}{\sim} f(y_i | \theta_i)$ for $i = 1, \dots, n$, and suppose $\theta_i \sim \pi(\theta | \lambda)$ with common **hyperparameter** λ . In simple Bayes, λ is fixed in advance. In **hierarchical Bayes**, one adds a prior on λ . Here we focus instead on **empirical Bayes**.

The marginal (unconditional) distribution of Y_i is

$$m(y_i) = \int f(y_i | \theta_i) \pi(\theta_i | \lambda) d\theta_i.$$

This is an instance of the law of total probability. After integrating out θ_i , $m(y_i)$ depends only on λ . The empirical Bayes procedure first estimates λ from $\prod_{i=1}^n m(y_i)$ by, for example, maximum likelihood or the method of moments, and then uses the posterior $\pi(\theta_i | y_i, \hat{\lambda})$, which depends on the data $Y_1, \dots, Y_n$ through the estimate $\hat{\lambda}$.

Example 23.1 If $Y_i \sim \text{Binomial}(n, p_i)$ and $p_i \sim \text{Beta}(\alpha, \beta)$, then the posterior expectation is

$$\mathbb{E}[p_i | Y_i] = \frac{Y_i + \alpha}{n + \alpha + \beta}.$$

The vanilla Bayes formulation fixes α and β (for example at 1 and 1 or 1/2 and 1/2), while empirical Bayes estimates them as $\hat{\alpha}$ and $\hat{\beta}$, respectively, and then uses

$$\frac{Y_i + \hat{\alpha}}{n + \hat{\alpha} + \hat{\beta}}.$$

Example 23.2 Let $Y_i \stackrel{\text{indep}}{\sim} N(\theta_i, \sigma^2)$ with known σ^2 , and prior $\theta_i \sim N(\mu, \tau^2)$. The posterior is

$$\pi(\theta_i | Y_i) = \frac{f(y_i | \theta_i) \pi(\theta_i | \mu)}{\int f(y_i | \theta_i) \pi(\theta_i | \mu) d\theta_i}$$

$$= \frac{\frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{(y_i - \theta_i)^2}{2\sigma^2}\right) \cdot \frac{1}{\sqrt{2\pi}\tau} \exp\left(-\frac{(\theta_i - \mu)^2}{2\tau^2}\right)}{\int \frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{(y_i - \theta_i)^2}{2\sigma^2}\right) \cdot \frac{1}{\sqrt{2\pi}\tau} \exp\left(-\frac{(\theta_i - \mu)^2}{2\tau^2}\right) d\theta_i}.$$

The denominator works out to the density of the $N(\mu, \sigma^2 + \tau^2)$ distribution, which is the marginal distribution of $y_i | \mu$. Using this, we obtain

$$\pi(\theta_i | Y_i) = \frac{\frac{1}{\sqrt{2\pi}\sigma\tau}}{\frac{1}{\sqrt{2\pi}\sqrt{\sigma^2 + \tau^2}}} \exp\left(-\left[\frac{(y_i - \theta_i)^2}{2\sigma^2} + \frac{(\theta_i - \mu)^2}{2\tau^2} - \frac{(y_i - \mu)^2}{2(\sigma^2 + \tau^2)}\right]\right).$$

Now simplify the exponential term. Let $B = \frac{\sigma^2}{\sigma^2 + \tau^2}$. Then the negative of the argument in the exponential term is

$$\begin{aligned} \arg &= \left[\frac{(y_i - \theta_i)^2}{2\sigma^2} + \frac{(\theta_i - \mu)^2}{2\tau^2} - \frac{(y_i - \mu)^2}{2(\sigma^2 + \tau^2)}\right] \cdot \frac{\sigma^2\tau^2}{\sigma^2 + \tau^2} = \dots \\ &= \left[\frac{(1-B)(y_i - \theta_i)^2 + B(\theta_i - \mu)^2 - B(1-B)(y_i - \mu)^2}{2B(1-B)}\right] \frac{1}{\sigma^2 + \tau^2} \\ &= \left[\frac{(1-B)(y_i^2 - 2\theta_i y_i + \theta_i^2) + B(\theta_i^2 - 2\theta_i \mu + \mu^2)}{2B(1-B)}\right] \frac{1}{\sigma^2 + \tau^2} \\ &\quad - \left[\frac{B(1-B)(y_i^2 - 2y_i \mu + \mu^2)}{2B(1-B)}\right] \frac{1}{\sigma^2 + \tau^2} \\ &= \left[\frac{\theta_i^2 - 2[(1-B)y_i + B\mu]\theta_i + (1-B)y_i^2 + B\mu^2}{2B(1-B)}\right] \frac{1}{\sigma^2 + \tau^2} \\ &\quad + \left[\frac{-(1-B)By_i^2 + 2B(1-B)y_i\mu - B(1-B)\mu^2}{2B(1-B)}\right] \frac{1}{\sigma^2 + \tau^2} \\ &= \left[\frac{(\theta_i - [(1-B)y_i + B\mu])^2}{2B(1-B)}\right] \frac{1}{\sigma^2 + \tau^2} \\ &= \frac{(\theta_i - [(1-B)y_i + B\mu])^2}{2} \cdot \frac{\sigma^2 + \tau^2}{\sigma^2\tau^2} \end{aligned}$$

Then

$$\pi(\theta_i | Y_i) = \frac{1}{\sqrt{2\pi} \frac{\sigma\tau}{\sqrt{\sigma^2 + \tau^2}}} \exp\left(-\frac{(\theta_i - [(1-B)y_i + B\mu])^2}{2} \cdot \frac{\sigma^2 + \tau^2}{\sigma^2\tau^2}\right)$$

so the posterior is the

$$N\left((1-B)Y_i + B\mu, \frac{\sigma^2\tau^2}{\sigma^2 + \tau^2}\right)$$

distribution.

In simple Bayes, μ is specified in advance, giving the posterior mean

$$(1-B)Y_i + B\mu,$$

where $B = \frac{\sigma^2}{\sigma^2 + \tau^2}$. On the other hand, in empirical Bayes, we estimate μ from $Y_1, \dots, Y_n \sim N(\mu, \sigma^2 + \tau^2)$, so $\hat{\mu} = \bar{Y}$ if we use maximum likelihood or the method of moments. Then the posterior mean of θ_i is

$$(1-B)Y_i + B\bar{Y} = \bar{Y} + (1-B)(Y_i - \bar{Y}).$$

The key feature is that the posterior of θ_i depends not only on Y_i , but also on all $Y_1, \dots, Y_n$ through $\bar{Y}$.

Example 23.3 In [Example 23.2](#), if τ^2 is unknown, we could instead estimate $\sigma^2 + \tau^2$ from the marginal model $Y_i \sim N(\mu, \sigma^2 + \tau^2)$. From the one-sample t -test framework,

$$\frac{\sum_{i=1}^n (Y_i - \bar{Y})^2}{\sigma^2 + \tau^2} \sim \chi_{(n-1)}^2, \quad \frac{\sigma^2 + \tau^2}{\sum_{i=1}^n (Y_i - \bar{Y})^2} \sim \text{Inv-}\chi_{(n-1)}^2.$$

The latter distribution is an **inverse-chi-squared** distribution with $n-1$ degrees of freedom. If $V \sim \text{Inv-}\chi_{(n)}^2$, then $\mathbb{E}[V] = \frac{1}{(n-2)}$, so

$$\mathbb{E}\left[\frac{\sigma^2 + \tau^2}{\sum_{i=1}^n (Y_i - \bar{Y})^2}\right] = \frac{1}{n-3}.$$

Hence

$$\frac{n-3}{\sum_{i=1}^n (Y_i - \bar{Y})^2}$$

is unbiased for $\frac{1}{(\sigma^2 + \tau^2)}$, leading to the empirical Bayes estimator

$$\hat{\theta}_i = \bar{Y} + \left(1 - \frac{\sigma^2(n-3)}{\sum_{i=1}^n (Y_i - \bar{Y})^2} \right) (Y_i - \bar{Y}).$$

This is called the **James-Stein estimator** for θ_i .

Lecture 24 — Thursday, April 03, 2014

Posterior Odds

Frequentist hypothesis tests are calibrated by the significance level and power, but they do not directly assign probabilities to competing hypotheses after observing data. In a Bayesian framework, we begin with prior probabilities for H_0 and H_A , and then update them using the observed result of a test. The key quantity is the **posterior odds**

$$\frac{\mathbb{P}(H_A \text{ true} \mid E)}{\mathbb{P}(H_0 \text{ true} \mid E)},$$

where E denotes the event that the data favor H_A (equivalently, that the testing procedure rejects H_0).

Using Bayes' theorem,

$$\mathbb{P}(H_A \text{ true} \mid E) = \frac{\mathbb{P}(E \mid H_A) \mathbb{P}(H_A)}{\mathbb{P}(E)}.$$

An analogous expression holds for H_0 . Taking the ratio gives

$$\frac{\mathbb{P}(H_A \text{ true} \mid E)}{\mathbb{P}(H_0 \text{ true} \mid E)} = \frac{\mathbb{P}(E \mid H_A)}{\mathbb{P}(E \mid H_0)} \cdot \frac{\mathbb{P}(H_A)}{\mathbb{P}(H_0)} = \frac{\text{power}}{\text{significance level}} \cdot \frac{\mathbb{P}(H_A)}{\mathbb{P}(H_0)}.$$

The term

$$\frac{\mathbb{P}(H_A)}{\mathbb{P}(H_0)}$$

is called the **prior odds**, and the full ratio above is the **posterior odds**. Thus the testing procedure contributes an evidence multiplier $\frac{\text{power}}{\text{significance level}}$ that updates prior odds into posterior odds.

For instance, if the prior odds are 1 : 1, the power is 0.80, and the significance level is 0.05, then the posterior odds become 16 : 1 in favor of H_A .

Appendix: Lecture and Tutorial Index

1. Lecture 1 — Tuesday, January 07, 2014
2. Lecture 2 — Thursday, January 09, 2014
3. Lecture 3 — Tuesday, January 14, 2014
4. Lecture 4 — Thursday, January 16, 2014
5. Lecture 5 — Tuesday, January 21, 2014
6. Lecture 6 — Thursday, January 23, 2014
7. Lecture 7 — Tuesday, January 28, 2014
8. Lecture 8 — Thursday, January 30, 2014
9. Lecture 9 — Tuesday, February 04, 2014
10. Lecture 10 — Thursday, February 06, 2014
11. Lecture 11 — Tuesday, February 11, 2014
12. Lecture 12 — Thursday, February 13, 2014
13. Lecture 13 — Tuesday, February 25, 2014
14. Lecture 14 — Tuesday, March 04, 2014
15. Lecture 15 — Thursday, March 06, 2014
16. Lecture 16 — Tuesday, March 11, 2014
17. Lecture 17 — Thursday, March 13, 2014
18. Lecture 18 — Tuesday, March 18, 2014
19. Lecture 19 — Thursday, March 20, 2014
20. Lecture 21 — Tuesday, March 25, 2014
21. Lecture 22 — Thursday, March 27, 2014
22. Lecture 23 — Tuesday, April 01, 2014
23. Lecture 24 — Thursday, April 03, 2014