

Topos Theory V: Sheaves

Robbert Liu

October 19, 2022

- Grothendieck Sites

Contents

- Grothendieck Sites
- Examples of Sites

Contents

- Grothendieck Sites
- Examples of Sites
- Sheaves on a Site

Generalized Topologies

- Grothendieck observes the following duality between Galois theory and covering space theory:

Galois Theory	Covering Space Theory
Field K of characteristic 0	(Locally) arcwise connected space X
Normal extension $m : K \hookrightarrow N$	Covering space $\rho : Y \twoheadrightarrow X$
Galois group of N/K	Covering group of ρ
Field automorphisms σ of N/K	Deck transformations $\sigma : Y \rightarrow Y$
Factorizations $K \hookrightarrow N_\sigma \hookrightarrow N$	Factorizations $Y \twoheadrightarrow Y_\sigma \twoheadrightarrow X$
$Y \otimes_K N \cong \bigotimes_i N$, N splitting	$Y \times_X U \rightarrow U \cong \prod_i U$ for some $U \ni x$
$K \rightarrow N$	$U \hookrightarrow X$

Generalized Topologies

- Grothendieck observes the following duality between Galois theory and covering space theory:

Galois Theory	Covering Space Theory
Field K of characteristic 0	(Locally) arcwise connected space X
Normal extension $m : K \twoheadrightarrow N$	Covering space $\rho : Y \twoheadrightarrow X$
Galois group of N/K	Covering group of ρ
Field automorphisms σ of N/K	Deck transformations $\sigma : Y \rightarrow Y$
Factorizations $K \twoheadrightarrow N_\sigma \twoheadrightarrow N$	Factorizations $Y \twoheadrightarrow Y_\sigma \twoheadrightarrow X$
$Y \otimes_K N \cong \bigotimes_i N$, N splitting	$Y \times_X U \rightarrow U \cong \prod_i U$ for some $U \ni x$
$K \rightarrow N$	$U \twoheadrightarrow X$

- Grothendieck.* We need a more general notion of topology where the primitive notions are not open sets $U \twoheadrightarrow X$, but more general maps.

Grothendieck Topologies

- *Idea.* Define a “topology” on a category by assigning each object its set of “open covers” (in actuality: sieves).

Grothendieck Topologies

- *Idea.* Define a “topology” on a category by assigning each object its set of “open covers” (in actuality: sieves).
- *Definition.* A **Grothendieck topology** on a category \mathbf{C} is a functor J assigning each object C a collection $J(C)$ of sieves on C such that:

Grothendieck Topologies

- *Idea.* Define a “topology” on a category by assigning each object its set of “open covers” (in actuality: sieves).
- *Definition.* A **Grothendieck topology** on a category \mathbf{C} is a functor J assigning each object C a collection $J(C)$ of sieves on C such that:
 - 1 $J(C)$ contains the maximal sieve C .

Grothendieck Topologies

- *Idea.* Define a “topology” on a category by assigning each object its set of “open covers” (in actuality: sieves).
- *Definition.* A **Grothendieck topology** on a category \mathbf{C} is a functor J assigning each object C a collection $J(C)$ of sieves on C such that:
 - 1 $J(C)$ contains the maximal sieve C .
 - 2 *Stability Axiom.* If $S \in J(C)$, then $h^*(S) \in J(D)$ for any morphism $h : D \rightarrow C$.

Grothendieck Topologies

- *Idea.* Define a “topology” on a category by assigning each object its set of “open covers” (in actuality: sieves).
- *Definition.* A **Grothendieck topology** on a category \mathbf{C} is a functor J assigning each object C a collection $J(C)$ of sieves on C such that:
 - 1 $J(C)$ contains the maximal sieve C .
 - 2 *Stability Axiom.* If $S \in J(C)$, then $h^*(S) \in J(D)$ for any morphism $h : D \rightarrow C$.
 - 3 *Transitivity Axiom.* If $S \in J(C)$ and R is any sieve on C such that $h^*(R) \in J(D)$ for all $h : D \rightarrow C \in S$, then $R \in J(C)$.

Grothendieck Topologies

- *Idea.* Define a “topology” on a category by assigning each object its set of “open covers” (in actuality: sieves).
- *Definition.* A **Grothendieck topology** on a category \mathbf{C} is a functor J assigning each object C a collection $J(C)$ of sieves on C such that:
 - 1 $J(C)$ contains the maximal sieve C .
 - 2 *Stability Axiom.* If $S \in J(C)$, then $h^*(S) \in J(D)$ for any morphism $h : D \rightarrow C$.
 - 3 *Transitivity Axiom.* If $S \in J(C)$ and R is any sieve on C such that $h^*(R) \in J(D)$ for all $h : D \rightarrow C \in S$, then $R \in J(C)$.
- A tuple (\mathbf{C}, J) , \mathbf{C} small, is called a **(Grothendieck) site**.

Sieves as Covers

- *Alternative terminology.* we say S covers C if $S \in J(C)$, and S covers $f : D \rightarrow C$ if $S \in J(C)$ and f^*S covers D .

Sieves as Covers

- *Alternative terminology.* we say S covers C if $S \in J(C)$, and S covers $f : D \rightarrow C$ if $S \in J(C)$ and f^*S covers D .
- We reformulate the axioms as follows:
 - 1 If S is a sieve on C and $f : D \rightarrow C \in S$, then S covers $f : D \rightarrow C$.

Sieves as Covers

- *Alternative terminology.* we say S covers C if $S \in J(C)$, and S covers $f : D \rightarrow C$ if $S \in J(C)$ and f^*S covers D .
- We reformulate the axioms as follows:
 - 1 If S is a sieve on C and $f : D \rightarrow C \in S$, then S covers $f : D \rightarrow C$.
 - 2 *Stability.* If S covers $f : D \rightarrow C$, then it also covers $fg : E \rightarrow C$ for any morphism $g : E \rightarrow D$

Sieves as Covers

- *Alternative terminology.* we say S covers C if $S \in J(C)$, and S covers $f : D \rightarrow C$ if $S \in J(C)$ and f^*S covers D .
- We reformulate the axioms as follows:
 - 1 If S is a sieve on C and $f : D \rightarrow C \in S$, then S covers $f : D \rightarrow C$.
 - 2 *Stability.* If S covers $f : D \rightarrow C$, then it also covers $fg : E \rightarrow C$ for any morphism $g : E \rightarrow D$
 - 3 *Transitivity.* If S covers $f : D \rightarrow C$ and R is a sieve on C which covers all morphisms in S , then R covers f .

Preliminary Consequences

- *Consequence of Transitivity.* If S covers C , and if we have a cover R_f of each $f : D_f \rightarrow C$ in S , then the set of composites fg with $f \in S$ and $g \in R_f$ covers C .

Preliminary Consequences

- *Consequence of Transitivity.* If S covers C , and if we have a cover R_f of each $f : D_f \rightarrow C$ in S , then the set of composites fg with $f \in S$ and $g \in R_f$ covers C .
- If R and S cover $g : D \rightarrow C$, then $R \cap S$ covers g .

Preliminary Consequences

- *Consequence of Transitivity.* If S covers C , and if we have a cover R_f of each $f : D_f \rightarrow C$ in S , then the set of composites fg with $f \in S$ and $g \in R_f$ covers C .
- If R and S cover $g : D \rightarrow C$, then $R \cap S$ covers g .
- *Proof.* If $f : D \rightarrow C \in R$, then $f^*(R \cap S) = f^*(S)$, since any morphism in $f^*(S)$, when composed with $f \in R$, is also an element of R . Hence, $f^*(R \cap S)$ covers D .

Basis for a Grothendieck Topology

- Equivalently, we can define a topology by specifying a **basis**.

Basis for a Grothendieck Topology

- Equivalently, we can define a topology by specifying a **basis**.
- *Definition.* A **basis** for a Grothendieck topology is a function K which assigns to each object C of a category a family of morphisms with codomain C , such that

Basis for a Grothendieck Topology

- Equivalently, we can define a topology by specifying a **basis**.
- *Definition.* A **basis** for a Grothendieck topology is a function K which assigns to each object C of a category a family of morphisms with codomain C , such that
 - 1 If $f : C' \rightarrow C$ is an isomorphism, then $\{f\} \in K(C)$.

Basis for a Grothendieck Topology

- Equivalently, we can define a topology by specifying a **basis**.
- *Definition.* A **basis** for a Grothendieck topology is a function K which assigns to each object C of a category a family of morphisms with codomain C , such that
 - 1 If $f : C' \rightarrow C$ is an isomorphism, then $\{f\} \in K(C)$.
 - 2 *Stability.* If $\{f_i : C_i \rightarrow C \mid i \in I\}$ is in $K(C)$, then for any morphism $g : D \rightarrow C$, the family of pullbacks $\pi_2 : C_i \times_C D \rightarrow D \mid i \in I$ is in $K(D)$.

Basis for a Grothendieck Topology

- Equivalently, we can define a topology by specifying a **basis**.
- *Definition.* A **basis** for a Grothendieck topology is a function K which assigns to each object C of a category a family of morphisms with codomain C , such that
 - 1 If $f : C' \rightarrow C$ is an isomorphism, then $\{f\} \in K(C)$.
 - 2 *Stability.* If $\{f_i : C_i \rightarrow C \mid i \in I\}$ is in $K(C)$, then for any morphism $g : D \rightarrow C$, the family of pullbacks $\pi_2 : C_i \times_C D \rightarrow D \mid i \in I$ is in $K(D)$.
 - 3 *Transitivity.* If $\{f_i : C_i \rightarrow C \mid i \in I\}$ is in $K(C)$ and if, for each i , $K(C_i)$ contains a family $\{g_{ij} : D_{ij} \rightarrow C_i \mid j \in I_i\}$, then the family of composites $\{f_i g_{ij}\}$ is in $K(C)$.

Basis for a Grothendieck Topology

- Equivalently, we can define a topology by specifying a **basis**.
- *Definition.* A **basis** for a Grothendieck topology is a function K which assigns to each object C of a category a family of morphisms with codomain C , such that
 - 1 If $f : C' \rightarrow C$ is an isomorphism, then $\{f\} \in K(C)$.
 - 2 *Stability.* If $\{f_i : C_i \rightarrow C \mid i \in I\}$ is in $K(C)$, then for any morphism $g : D \rightarrow C$, the family of pullbacks $\pi_2 : C_i \times_C D \rightarrow D \mid i \in I$ is in $K(D)$.
 - 3 *Transitivity.* If $\{f_i : C_i \rightarrow C \mid i \in I\}$ is in $K(C)$ and if, for each i , $K(C_i)$ contains a family $\{g_{ij} : D_{ij} \rightarrow C_i \mid j \in I_i\}$, then the family of composites $\{f_i g_{ij}\}$ is in $K(C)$.
- From K , we get a unique topology J , such that $S \in J(C)$ iff S contains some $R \in K(C)$.

Examples of Topologies

- Given any topological space X , the open set category $\mathcal{O}(X)$ has the topology assigning each object U its set of covering sieves.

Examples of Topologies

- Given any topological space X , the open set category $\mathcal{O}(X)$ has the topology assigning each object U its set of covering sieves.
- *Trivial Topology.* Given any category \mathbf{C} , the coarsest topology on \mathbf{C} is the **trivial topology** $\tau_{\mathbf{C}}$ containing only the maximal sieve $\tau_{\mathbf{C}}$.

Examples of Topologies

- Given any topological space X , the open set category $\mathcal{O}(X)$ has the topology assigning each object U its set of covering sieves.
- *Trivial Topology.* Given any category \mathbf{C} , the coarsest topology on \mathbf{C} is the **trivial topology** $\tau_{\mathbf{C}}$ containing only the maximal sieve $\tau_{\mathbf{C}}$.
- *Open Cover Topology.* Given a small subcategory $\mathbf{T} \leq \mathbf{Top}$, which is closed under finite limits and restriction to open subspaces, the **open cover topology** is generated by the basis K defined by $\{\iota_i : Y_i \hookrightarrow X \mid i \in I\} \in K(X)$, iff each Y_i is an open subspace of X with the inclusion ι_i , and $\bigcup_i Y_i = X$.

Examples of Topologies

- Given any topological space X , the open set category $\mathcal{O}(X)$ has the topology assigning each object U its set of covering sieves.
- *Trivial Topology.* Given any category \mathbf{C} , the coarsest topology on \mathbf{C} is the **trivial topology** $\tau_{\mathbf{C}}$ containing only the maximal sieve $\tau_{\mathbf{C}}$.
- *Open Cover Topology.* Given a small subcategory $\mathbf{T} \leq \mathbf{Top}$, which is closed under finite limits and restriction to open subspaces, the **open cover topology** is generated by the basis K defined by $\{\iota_i : Y_i \hookrightarrow X \mid i \in I\} \in K(X)$, iff each Y_i is an open subspace of X with the inclusion ι_i , and $\bigcup_i Y_i = X$.

Examples of Topologies

- Given any topological space X , the open set category $\mathcal{O}(X)$ has the topology assigning each object U its set of covering sieves.
- *Trivial Topology.* Given any category \mathbf{C} , the coarsest topology on \mathbf{C} is the **trivial topology** $\tau_{\mathbf{C}}$ containing only the maximal sieve $\tau_{\mathbf{C}}$.
- *Open Cover Topology.* Given a small subcategory $\mathbf{T} \leq \mathbf{Top}$, which is closed under finite limits and restriction to open subspaces, the **open cover topology** is generated by the basis K defined by $\{\iota_i : Y_i \hookrightarrow X \mid i \in I\} \in K(X)$, iff each Y_i is an open subspace of X with the inclusion ι_i , and $\bigcup_i Y_i = X$.
- A finer category on \mathbf{T} is generated by the basis K assigning to X the set of all families $\{f_i : Y_i \rightarrow X \mid i \in I\}$ such that $f : \coprod_i Y_i \rightarrow X$ is an open surjection.

More Examples

- *Sup Topology.* A complete Heyting algebra is a HA admitting sups and infs over any family. The **sup topology** on a cHa is generated by the basis K , defined by $\{a_i : i \in I\} \in K(c)$ if $\bigwedge_{i \in I} a_i = c$; this generalizes the open set topology to any complete Heyting algebra. Stability follows from

$$\bigwedge_{i \in I} (b \wedge a_i) = b \wedge \bigwedge_{i \in I} a_i$$

More Examples

- *Sup Topology.* A complete Heyting algebra is a HA admitting sups and infs over any family. The **sup topology** on a cHa is generated by the basis K , defined by $\{a_i : i \in I\} \in K(c)$ if $\bigwedge_{i \in I} a_i = c$; this generalizes the open set topology to any complete Heyting algebra. Stability follows from

$$\bigwedge_{i \in I} (b \wedge a_i) = b \wedge \bigwedge_{i \in I} a_i$$

- *Dense Topology.* Define the **dense topology** J on a category \mathbf{C} by $S \in J(C)$ if for any $f : D \rightarrow C$, there is a $g : E \rightarrow D$ such that $fg \in S$. When the category is a poset P , $J(p)$ is simply the set of dense subsets below $p \in P$.

More Examples

- *Sup Topology.* A complete Heyting algebra is a HA admitting sups and infs over any family. The **sup topology** on a cHa is generated by the basis K , defined by $\{a_i : i \in I\} \in K(c)$ if $\bigwedge_{i \in I} a_i = c$; this generalizes the open set topology to any complete Heyting algebra. Stability follows from

$$\bigwedge_{i \in I} (b \wedge a_i) = b \wedge \bigwedge_{i \in I} a_i$$

- *Dense Topology.* Define the **dense topology** J on a category \mathbf{C} by $S \in J(C)$ if for any $f : D \rightarrow C$, there is a $g : E \rightarrow D$ such that $fg \in S$. When the category is a poset P , $J(p)$ is simply the set of dense subsets below $p \in P$.
- *Atomic Topology.* The **atomic category** is defined by $S \in J(C)$ iff S is a nonempty sieve.

The Zariski Site

- Given an ideal $I \subseteq \mathbb{C}[x_1, \dots, x_n]$, we define $V(I) = \{x \in \mathbb{C}^n : I \text{ vanishes on } x\}$.
 $V(I)$ is called a **complex affine variety**.

The Zariski Site

- Given an ideal $I \subseteq \mathbb{C}[x_1, \dots, x_n]$, we define $V(I) = \{x \in \mathbb{C}^n : I \text{ vanishes on } x\}$. $V(I)$ is called a **complex affine variety**.
- We can then form a topology on \mathbb{C}^n called the **Zariski topology** by specifying the closed sets to be all $V(I)$. The topology is generated by the subbasis of sets of the form $V(P)^c$, for prime ideals P , which is the nonvanishing set of P .

The Zariski Site

- Given an ideal $I \subseteq \mathbb{C}[x_1, \dots, x_n]$, we define $V(I) = \{x \in \mathbb{C}^n : I \text{ vanishes on } x\}$. $V(I)$ is called a **complex affine variety**.
- We can then form a topology on \mathbb{C}^n called the **Zariski topology** by specifying the closed sets to be all $V(I)$. The topology is generated by the subbasis of sets of the form $V(P)^c$, for prime ideals P , which is the nonvanishing set of P .
- The Zariski topology has a structure sheaf \mathcal{O} , defined by letting $\mathcal{O}(U)$ to be the set of all rational functions h over \mathbb{C} which are *defined* at each point of U .

The Zariski Site

- Given an ideal $I \subseteq \mathbb{C}[x_1, \dots, x_n]$, we define $V(I) = \{x \in \mathbb{C}^n : I \text{ vanishes on } x\}$. $V(I)$ is called a **complex affine variety**.
- We can then form a topology on \mathbb{C}^n called the **Zariski topology** by specifying the closed sets to be all $V(I)$. The topology is generated by the subbasis of sets of the form $V(P)^c$, for prime ideals P , which is the nonvanishing set of P .
- The Zariski topology has a structure sheaf \mathcal{O} , defined by letting $\mathcal{O}(U)$ to be the set of all rational functions h over \mathbb{C} which are *defined* at each point of U .
- Moreover, \mathcal{O} is a sheaf of rings, with each stalk being a local ring, i.e. contains a unique maximal ideal.

The Zariski Site

- Given an ideal $I \subseteq \mathbb{C}[x_1, \dots, x_n]$, we define $V(I) = \{x \in \mathbb{C}^n : I \text{ vanishes on } x\}$. $V(I)$ is called a **complex affine variety**.
- We can then form a topology on \mathbb{C}^n called the **Zariski topology** by specifying the closed sets to be all $V(I)$. The topology is generated by the subbasis of sets of the form $V(P)^c$, for prime ideals P , which is the nonvanishing set of P .
- The Zariski topology has a structure sheaf \mathcal{O} , defined by letting $\mathcal{O}(U)$ to be the set of all rational functions h over \mathbb{C} which are *defined* at each point of U .
- Moreover, \mathcal{O} is a sheaf of rings, with each stalk being a local ring, i.e. contains a unique maximal ideal.
- The general construction can be done for a commutative ring k .

The Zariski Site (Cont.)

- Given a commutative unital ring A , define the **localization** $A_{[a^{-1}]}$ containing elements b/a^n , where $b \in A$ and $n \in \mathbb{N}$.

The Zariski Site (Cont.)

- Given a commutative unital ring A , define the **localization** $A_{[a^{-1}]}$ containing elements b/a^n , where $b \in A$ and $n \in \mathbb{N}$.
- Instead of working with varieties $V(I)$, we define a **finitely presented k -algebra** A to be of the form $k[x_1, \dots, x_m]/(f_1, \dots, f_m)$, where $f_i \in k[x_1, \dots, x_m]$.

The Zariski Site (Cont.)

- Given a commutative unital ring A , define the **localization** $A_{[a^{-1}]}$ containing elements b/a^n , where $b \in A$ and $n \in \mathbb{N}$.
- Instead of working with varieties $V(I)$, we define a **finitely presented k -algebra** A to be of the form $k[x_1, \dots, x_m]/(f_1, \dots, f_m)$, where $f_i \in k[x_1, \dots, x_m]$.
- This construction provides more information than varieties: the k -algebra defined by $x - y$ is different from that defined by $(x - y)^2$.

The Zariski Site (Cont.)

- Given a commutative unital ring A , define the **localization** $A_{[a^{-1}]}$ containing elements b/a^n , where $b \in A$ and $n \in \mathbb{N}$.
- Instead of working with varieties $V(I)$, we define a **finitely presented k -algebra** A to be of the form $k[x_1, \dots, x_m]/(f_1, \dots, f_m)$, where $f_i \in k[x_1, \dots, x_m]$.
- This construction provides more information than varieties: the k -algebra defined by $x - y$ is different from that defined by $(x - y)^2$.
- Let $(k - \text{Alg})_{fp}$ denote the category of all these algebras. The objects are no longer point-sets but algebras, so we cannot define functions at points.

The Zariski Site (Cont.)

- Given a commutative unital ring A , define the **localization** $A_{[a^{-1}]}$ containing elements b/a^n , where $b \in A$ and $n \in \mathbb{N}$.
- Instead of working with varieties $V(I)$, we define a **finitely presented k -algebra** A to be of the form $k[x_1, \dots, x_m]/(f_1, \dots, f_m)$, where $f_i \in k[x_1, \dots, x_m]$.
- This construction provides more information than varieties: the k -algebra defined by $x - y$ is different from that defined by $(x - y)^2$.
- Let $(k - \text{Alg})_{fp}$ denote the category of all these algebras. The objects are no longer point-sets but algebras, so we cannot define functions at points.
- We can instead define a Grothendieck topology on $(k - \text{Alg})_{fp}$. Covers of a k -algebra A will be determined by elements a_1, \dots, a_n such that $(a_1, \dots, a_n) = A$. Each cover will be the dual of a family of the form $\{A \rightarrow A[a_i^{-1}]\}_i$.

The Zariski Site (Cont.)

- Given a commutative unital ring A , define the **localization** $A_{[a^{-1}]}$ containing elements b/a^n , where $b \in A$ and $n \in \mathbb{N}$.
- Instead of working with varieties $V(I)$, we define a **finitely presented k -algebra** A to be of the form $k[x_1, \dots, x_m]/(f_1, \dots, f_m)$, where $f_i \in k[x_1, \dots, x_m]$.
- This construction provides more information than varieties: the k -algebra defined by $x - y$ is different from that defined by $(x - y)^2$.
- Let $(k - \text{Alg})_{fp}$ denote the category of all these algebras. The objects are no longer point-sets but algebras, so we cannot define functions at points.
- We can instead define a Grothendieck topology on $(k - \text{Alg})_{fp}$. Covers of a k -algebra A will be determined by elements a_1, \dots, a_n such that $(a_1, \dots, a_n) = A$. Each cover will be the dual of a family of the form $\{A \rightarrow A[a_i^{-1}]\}_i$.
- The last step is to define a suitable structure sheaf on $(k - \text{Alg})_{fp}$ which acts like “the ring of functions on each neighbourhood”.

Sheaves on a Site

- Let $P : \mathbf{C}^{\text{op}} \rightarrow \mathbf{Set}$ be a presheaf.

Sheaves on a Site

- Let $P : \mathcal{C}^{\text{op}} \rightarrow \mathbf{Set}$ be a presheaf.
- Given a covering sieve S of an object C , **matching family** for S of elements of P is a family $\{x_f \mid f : D \rightarrow C \in S\}$, such that $x_f \cdot g = x_{fg}$ for any morphism $g : E \rightarrow D$.

Sheaves on a Site

- Let $P : \mathbf{C}^{\text{op}} \rightarrow \mathbf{Set}$ be a presheaf.
- Given a covering sieve S of an object C , **matching family** for S of elements of P is a family $\{x_f \mid f : D \rightarrow C \in S\}$, such that $x_f \cdot g = x_{fg}$ for any morphism $g : E \rightarrow D$.
- An **amalgamation** of such a matching family is an element $x \in P(C)$ such that $x \cdot f = x_f$ for all $f \in S$.

Sheaves on a Site

- Let $P : \mathbf{C}^{\text{op}} \rightarrow \mathbf{Set}$ be a presheaf.
- Given a covering sieve S of an object C , **matching family** for S of elements of P is a family $\{x_f \mid f : D \rightarrow C \in S\}$, such that $x_f \cdot g = x_{fg}$ for any morphism $g : E \rightarrow D$.
- An **amalgamation** of such a matching family is an element $x \in P(C)$ such that $x \cdot f = x_f$ for all $f \in S$.
- Then, P is a sheaf precisely when all matching families for any cover of any object of \mathbf{C} has a unique amalgamation.

Sheaves on a Site

- Let $P : \mathbf{C}^{\text{op}} \rightarrow \mathbf{Set}$ be a presheaf.
- Given a covering sieve S of an object C , **matching family** for S of elements of P is a family $\{x_f \mid f : D \rightarrow C \in S\}$, such that $x_f \cdot g = x_{fg}$ for any morphism $g : E \rightarrow D$.
- An **amalgamation** of such a matching family is an element $x \in P(C)$ such that $x \cdot f = x_f$ for all $f \in S$.
- Then, P is a sheaf precisely when all matching families for any cover of any object of \mathbf{C} has a unique amalgamation.
- Regarding a sieve as a subfunctor of $\mathcal{Y}(C)$, a matching family $f \mapsto x_f$ is just a natural transformation $S \rightarrow P$; the above conditions reduces to if

$$\text{Hom}(S, P) \cong \text{Hom}(\mathcal{Y}(C), P)$$

Sheaves on a Site

- Let $P : \mathbf{C}^{\text{op}} \rightarrow \mathbf{Set}$ be a presheaf.
- Given a covering sieve S of an object C , **matching family** for S of elements of P is a family $\{x_f \mid f : D \rightarrow C \in S\}$, such that $x_f \cdot g = x_{fg}$ for any morphism $g : E \rightarrow D$.
- An **amalgamation** of such a matching family is an element $x \in P(C)$ such that $x \cdot f = x_f$ for all $f \in S$.
- Then, P is a sheaf precisely when all matching families for any cover of any object of \mathbf{C} has a unique amalgamation.
- Regarding a sieve as a subfunctor of $\mathcal{J}(C)$, a matching family $f \mapsto x_f$ is just a natural transformation $S \rightarrow P$; the above conditions reduces to if

$$\text{Hom}(S, P) \cong \text{Hom}(\mathcal{J}(C), P)$$

- This is true if we have a family of equalizers:

$$P(C) \xrightarrow{x \mapsto \{x_f\}_f} \prod_{f \in S} P(\text{dom } f) \xrightleftharpoons[x_f \mapsto x_f \cdot g]{x_f \mapsto x_{fg}} \prod_{\substack{f \in S \\ \text{dom } f = \text{cod } g}} P(\text{dom } g)$$

- Again, it suffices to describe a sheaf on a basis K . If $R = \{f_i : C_i \rightarrow C \mid i \in I\}$ is a K -cover of C , a matching family is one such that $x_i \cdot \pi_{ij}^1 = x_j \cdot \pi_{ij}^2$ always holds. An amalgamation for $\{x_i\}$ is a compatible element x .

- Again, it suffices to describe a sheaf on a basis K . If $R = \{f_i : C_i \rightarrow C \mid i \in I\}$ is a K -cover of C , a matching family is one such that $x_i \cdot \pi_{ij}^1 = x_j \cdot \pi_{ij}^2$ always holds. An amalgamation for $\{x_i\}$ is a compatible element x .
- *Proposition.* Let P be a presheaf on \mathbf{C} with basis K . Then, P is a sheaf iff for any cover $\{f_i : C_i \rightarrow C \mid i \in I\}$ in the basis K , any matching family $\{x_i\}_i$ has a unique amalgamation.

- Again, it suffices to describe a sheaf on a basis K . If $R = \{f_i : C_i \rightarrow C \mid i \in I\}$ is a K -cover of C , a matching family is one such that $x_i \cdot \pi_{ij}^1 = x_j \cdot \pi_{ij}^2$ always holds. An amalgamation for $\{x_i\}$ is a compatible element x .
- *Proposition.* Let P be a presheaf on \mathbf{C} with basis K . Then, P is a sheaf iff for any cover $\{f_i : C_i \rightarrow C \mid i \in I\}$ in the basis K , any matching family $\{x_i\}_i$ has a unique amalgamation.
- *Definition.* If J is a topology on \mathbf{C} such that all representable presheaves on \mathbf{C} are sheaves, then J is called **subcanonical**. The **canonical** topology is the largest subcanonical topology on \mathbf{C} . e.g. the open surjection topology is canonical.

- Again, it suffices to describe a sheaf on a basis K . If $R = \{f_i : C_i \rightarrow C \mid i \in I\}$ is a K -cover of C , a matching family is one such that $x_i \cdot \pi_{ij}^1 = x_j \cdot \pi_{ij}^2$ always holds. An amalgamation for $\{x_i\}$ is a compatible element x .
- *Proposition.* Let P be a presheaf on \mathbf{C} with basis K . Then, P is a sheaf iff for any cover $\{f_i : C_i \rightarrow C \mid i \in I\}$ in the basis K , any matching family $\{x_i\}_i$ has a unique amalgamation.
- *Definition.* If J is a topology on \mathbf{C} such that all representable presheaves on \mathbf{C} are sheaves, then J is called **subcanonical**. The **canonical** topology is the largest subcanonical topology on \mathbf{C} . e.g. the open surjection topology is canonical.
- *Proposition.* Let $I \rightarrow \mathbf{C}$ be a diagram of presheaves P_i on (\mathbf{C}, J) . Then if P_i are all sheaves, so is the limit $\lim_{\leftarrow} P_i$.

- Again, it suffices to describe a sheaf on a basis K . If $R = \{f_i : C_i \rightarrow C \mid i \in I\}$ is a K -cover of C , a matching family is one such that $x_i \cdot \pi_{ij}^1 = x_j \cdot \pi_{ij}^2$ always holds. An amalgamation for $\{x_i\}$ is a compatible element x .
- *Proposition.* Let P be a presheaf on \mathbf{C} with basis K . Then, P is a sheaf iff for any cover $\{f_i : C_i \rightarrow C \mid i \in I\}$ in the basis K , any matching family $\{x_i\}_i$ has a unique amalgamation.
- *Definition.* If J is a topology on \mathbf{C} such that all representable presheaves on \mathbf{C} are sheaves, then J is called **subcanonical**. The **canonical** topology is the largest subcanonical topology on \mathbf{C} . e.g. the open surjection topology is canonical.
- *Proposition.* Let $I \rightarrow \mathbf{C}$ be a diagram of presheaves P_i on (\mathbf{C}, J) . Then if P_i are all sheaves, so is the limit $\lim_{\leftarrow} P_i$.
- *Definition.* A **Grothendieck topos** is a category which is equivalent to the category $\text{Sh}(\mathbf{C}, J)$ of sheaves on some site (\mathbf{C}, J) .