

Sections over Vector Bundles and the Serre-Swan Theorem

Robbert Liu

April 14, 2022

Introduction.

Introduction.

- Vector bundles, family of finite dimensional vector spaces continuously parametrized by a base space. Locally looks like a “bundle of vectors spaces”.

Introduction.

- Vector bundles, family of finite dimensional vector spaces continuously parametrized by a base space. Locally looks like a “bundle of vectors spaces”.
- Sections, continuous choice of a vector at each vector space. Can be added and scaled by continuous functions.

Introduction.

- Vector bundles, family of finite dimensional vector spaces continuously parametrized by a base space. Locally looks like a “bundle of vectors spaces”.
- Sections, continuous choice of a vector at each vector space. Can be added and scaled by continuous functions.
- Recurring theme: algebraic vs geometric perspective.

Introduction.

- Vector bundles, family of finite dimensional vector spaces continuously parametrized by a base space. Locally looks like a “bundle of vectors spaces”.
- Sections, continuous choice of a vector at each vector space. Can be added and scaled by continuous functions.
- Recurring theme: algebraic vs geometric perspective.
- The Serre-Swan provides a way to switch between these two perspectives.

Vector bundles.

Vector bundles.

- Let $K \in \{\mathbb{R}, \mathbb{C}\}$. A topological K -vector bundle is a continuous surjective map $p : E \rightarrow B$ between topological spaces such that

Vector bundles.

- Let $K \in \{\mathbb{R}, \mathbb{C}\}$. A topological K -vector bundle is a continuous surjective map $p : E \rightarrow B$ between topological spaces such that
 - ▶ Each fiber $p^{-1}(x)$ is a finite dimensional vector space over K .

Vector bundles.

- Let $K \in \{\mathbb{R}, \mathbb{C}\}$. A topological K -vector bundle is a continuous surjective map $p : E \rightarrow B$ between topological spaces such that
 - ▶ Each fiber $p^{-1}(x)$ is a finite dimensional vector space over K .
 - ▶ Each $x \in B$ has a neighbourhood U and a homeomorphism $h : p^{-1}(U) \rightarrow U \times K^n$, such that h is linear when restricted to each fiber.

Vector bundles.

- Let $K \in \{\mathbb{R}, \mathbb{C}\}$. A topological K -vector bundle is a continuous surjective map $p : E \rightarrow B$ between topological spaces such that
 - ▶ Each fiber $p^{-1}(x)$ is a finite dimensional vector space over K .
 - ▶ Each $x \in B$ has a neighbourhood U and a homeomorphism $h : p^{-1}(U) \rightarrow U \times K^n$, such that h is linear when restricted to each fiber.
 - ▶ h is called a “local trivialization” around x .

Vector bundles.

- Let $K \in \{\mathbb{R}, \mathbb{C}\}$. A topological K -vector bundle is a continuous surjective map $p : E \rightarrow B$ between topological spaces such that
 - ▶ Each fiber $p^{-1}(x)$ is a finite dimensional vector space over K .
 - ▶ Each $x \in B$ has a neighbourhood U and a homeomorphism $h : p^{-1}(U) \rightarrow U \times K^n$, such that h is linear when restricted to each fiber.
 - ▶ h is called a “local trivialization” around x .
- *Remark.* the dimension $n(x) = \dim p^{-1}(y)$ of fibers around x is locally constant, so dimension is constant on connected components.

Vector bundles.

- Let $K \in \{\mathbb{R}, \mathbb{C}\}$. A topological K -vector bundle is a continuous surjective map $p : E \rightarrow B$ between topological spaces such that
 - ▶ Each fiber $p^{-1}(x)$ is a finite dimensional vector space over K .
 - ▶ Each $x \in B$ has a neighbourhood U and a homeomorphism $h : p^{-1}(U) \rightarrow U \times K^n$, such that h is linear when restricted to each fiber.
 - ▶ h is called a “local trivialization” around x .
- *Remark.* the dimension $n(x) = \dim p^{-1}(y)$ of fibers around x is locally constant, so dimension is constant on connected components.
- A map between vector bundles $p_1 : E_1 \rightarrow B, p_2 : E_2 \rightarrow B$ is a continuous map $f : E_1 \rightarrow E_2$ sending $p_1^{-1}(x)$ to $p_2^{-1}(x)$, which restricts to a linear map on each fiber.

Examples.

- The trivial bundle: $p : B \times U^k \rightarrow B$.

Examples.

- The trivial bundle: $p : B \times U^k \rightarrow B$.
- The tangent bundle $p : TS^n \rightarrow S^n$ of S^n in \mathbb{R}^{n+1} , containing elements $(x, v) \subseteq S^n \times \mathbb{R}^{n+1}$ such that v is tangent to x .

Examples.

- The trivial bundle: $p : B \times U^k \rightarrow B$.
- The tangent bundle $p : TS^n \rightarrow S^n$ of S^n in \mathbb{R}^{n+1} , containing elements $(x, v) \subseteq S^n \times \mathbb{R}^{n+1}$ such that v is tangent to x .
- The normal bundle $p : NS^n \rightarrow S^n$ of S^n in \mathbb{R}^{n+1} , containing elements (x, tx) .

Examples.

- The trivial bundle: $p : B \times U^k \rightarrow B$.
- The tangent bundle $p : TS^n \rightarrow S^n$ of S^n in \mathbb{R}^{n+1} , containing elements $(x, v) \subseteq S^n \times \mathbb{R}^{n+1}$ such that v is tangent to x .
- The normal bundle $p : NS^n \rightarrow S^n$ of S^n in \mathbb{R}^{n+1} , containing elements (x, tx) .
- The Möbius strip $p : \mu \rightarrow S^1$ and the annulus $p : \alpha \rightarrow S^1$.

Examples.

- The trivial bundle: $p : B \times U^k \rightarrow B$.
- The tangent bundle $p : TS^n \rightarrow S^n$ of S^n in \mathbb{R}^{n+1} , containing elements $(x, v) \subseteq S^n \times \mathbb{R}^{n+1}$ such that v is tangent to x .
- The normal bundle $p : NS^n \rightarrow S^n$ of S^n in \mathbb{R}^{n+1} , containing elements (x, tx) .
- The Möbius strip $p : \mu \rightarrow S^1$ and the annulus $p : \alpha \rightarrow S^1$.
- The canonical line bundle $p : E \rightarrow \mathbb{R}P^n$. $\mathbb{R}P^n$ can be viewed as the space of lines in \mathbb{R}^{n+1} intersecting 0; $E \subseteq \mathbb{R}P^n \times \mathbb{R}^{n+1}$ contains elements (ℓ, v) , where $v \in \ell$.

Sections.

- Given $p : E \rightarrow B$ and $U \subseteq B$, a section over U is a continuous choice $s : U \rightarrow E$ of a vector at each fiber of U . Sections over B are simply called sections.

Sections.

- Given $p : E \rightarrow B$ and $U \subseteq B$, a section over U is a continuous choice $s : U \rightarrow E$ of a vector at each fiber of U . Sections over B are simply called sections.

Sections.

- Given $p : E \rightarrow B$ and $U \subseteq B$, a section over U is a continuous choice $s : U \rightarrow E$ of a vector at each fiber of U . Sections over B are simply called sections.
- (*Lemma 1.6*) Using a local trivialization around $x \in B$, we can pull back a basis of $B \times \mathbb{R}^n$ to sections s_1, \dots, s_n over $U \ni x$ which are a basis at every fiber of $y \in U$.

Sections.

- Given $p : E \rightarrow B$ and $U \subseteq B$, a section over U is a continuous choice $s : U \rightarrow E$ of a vector at each fiber of U . Sections over B are simply called sections.
- (*Lemma 1.6*) Using a local trivialization around $x \in B$, we can pull back a basis of $B \times \mathbb{R}^n$ to sections s_1, \dots, s_n over $U \ni x$ which are a basis at every fiber of $y \in U$.
- (*Lemma 1.7*) Sections which are linearly independent at x remain linearly independent around x .

Sections.

- Given $p : E \rightarrow B$ and $U \subseteq B$, a section over U is a continuous choice $s : U \rightarrow E$ of a vector at each fiber of U . Sections over B are simply called sections.
- (*Lemma 1.6*) Using a local trivialization around $x \in B$, we can pull back a basis of $B \times \mathbb{R}^n$ to sections s_1, \dots, s_n over $U \ni x$ which are a basis at every fiber of $y \in U$.
- (*Lemma 1.7*) Sections which are linearly independent at x remain linearly independent around x .
- The set $\Gamma(E)$ of sections of $p : E \rightarrow B$ has a natural module structure over $C(B)$.

Subbundles.

- If $p : E \rightarrow B, p : E' \rightarrow B$ are vector bundles such that $E' \subseteq E$, then E' is a subbundle of E .

Subbundles.

- If $p : E \rightarrow B, p : E' \rightarrow B$ are vector bundles such that $E' \subseteq E$, then E' is a subbundle of E .
- *Question.* If $f : E_1 \rightarrow E_2$ is a bundle map, are $\text{im } f, \ker f$ always subbundles?

Subbundles.

- If $p : E \rightarrow B, p : E' \rightarrow B$ are vector bundles such that $E' \subseteq E$, then E' is a subbundle of E .
- *Question.* If $f : E_1 \rightarrow E_2$ is a bundle map, are $\text{im } f, \ker f$ always subbundles?
- No!

Subbundles.

- If $p : E \rightarrow B, p : E' \rightarrow B$ are vector bundles such that $E' \subseteq E$, then E' is a subbundle of E .
- *Question.* If $f : E_1 \rightarrow E_2$ is a bundle map, are $\text{im } f, \ker f$ always subbundles?
- No!
- Take $f : [0, 1] \times \mathbb{R} \rightarrow [0, 1] \times \mathbb{R}$ defined by $(t, x) \mapsto (t, tx)$, where $p : [0, 1] \times \mathbb{R} \rightarrow [0, 1]$ is a trivial bundle.

Subbundles.

- If $p : E \rightarrow B, p : E' \rightarrow B$ are vector bundles such that $E' \subseteq E$, then E' is a subbundle of E .
- *Question.* If $f : E_1 \rightarrow E_2$ is a bundle map, are $\operatorname{im} f, \ker f$ always subbundles?
- No!
- Take $f : [0, 1] \times \mathbb{R} \rightarrow [0, 1] \times \mathbb{R}$ defined by $(t, x) \mapsto (t, tx)$, where $p : [0, 1] \times \mathbb{R} \rightarrow [0, 1]$ is a trivial bundle.
- (*Proposition 1.10*) The fibers of $\operatorname{im} f$ have locally constant dimension $\iff \operatorname{im} f$ is a subbundle $\iff \ker f$ is a subbundle \iff the fibers of $\ker f$ have locally constant dimension.

Direct Sum.

- If $p_1 : E_1 \rightarrow B, p_2 : E_2 \rightarrow B$ are vector bundles, we can construct their direct sum $p : E_1 \oplus E_2 \rightarrow B$.

Direct Sum.

- If $p_1 : E_1 \rightarrow B, p_2 : E_2 \rightarrow B$ are vector bundles, we can construct their direct sum $p : E_1 \oplus E_2 \rightarrow B$.
- Fibers are direct sums $p_1^{-1}(x) \oplus p_2^{-1}(x)$, and p projects to x .

Direct Sum.

- If $p_1 : E_1 \rightarrow B, p_2 : E_2 \rightarrow B$ are vector bundles, we can construct their direct sum $p : E_1 \oplus E_2 \rightarrow B$.
- Fibers are direct sums $p_1^{-1}(x) \oplus p_2^{-1}(x)$, and p projects to x .
- Local trivializations are products of local trivializations for E_1, E_2 .

Direct Sum.

- If $p_1 : E_1 \rightarrow B, p_2 : E_2 \rightarrow B$ are vector bundles, we can construct their direct sum $p : E_1 \oplus E_2 \rightarrow B$.
- Fibers are direct sums $p_1^{-1}(x) \oplus p_2^{-1}(x)$, and p projects to x .
- Local trivializations are products of local trivializations for E_1, E_2 .
- Anomalous example: direct sum of a trivial bundle and nontrivial bundle which is trivial.

Direct Sum.

- If $p_1 : E_1 \rightarrow B, p_2 : E_2 \rightarrow B$ are vector bundles, we can construct their direct sum $p : E_1 \oplus E_2 \rightarrow B$.
- Fibers are direct sums $p_1^{-1}(x) \oplus p_2^{-1}(x)$, and p projects to x .
- Local trivializations are products of local trivializations for E_1, E_2 .
- Anomalous example: direct sum of a trivial bundle and nontrivial bundle which is trivial.
- *Question.* is a subbundle of a vector bundle always a summand?

Inner product.

- An inner product on $p : E \rightarrow B$ is a continuously varying map $\langle, \rangle : E \oplus E \rightarrow K$ which restricts to inner product on each fiber.

Inner product.

- An inner product on $p : E \rightarrow B$ is a continuously varying map $\langle, \rangle : E \oplus E \rightarrow K$ which restricts to inner product on each fiber.
- We can use an inner product on E to construct a projection $f : E \rightarrow E'$, if $E' \subseteq E$ is a subbundle.

Inner product.

- An inner product on $p : E \rightarrow B$ is a continuously varying map $\langle, \rangle : E \oplus E \rightarrow K$ which restricts to inner product on each fiber.
- We can use an inner product on E to construct a projection $f : E \rightarrow E'$, if $E' \subseteq E$ is a subbundle.
- Existence of an inner product is guaranteed if B is paracompact.

Inner product.

- An inner product on $p : E \rightarrow B$ is a continuously varying map $\langle, \rangle : E \oplus E \rightarrow K$ which restricts to inner product on each fiber.
- We can use an inner product on E to construct a projection $f : E \rightarrow E'$, if $E' \subseteq E$ is a subbundle.
- Existence of an inner product is guaranteed if B is paracompact.
- Theme: vector bundles decompose and coalesce nicely if B is nice.

The functor Γ .

- Recall that $\Gamma(E)$ is the set of sections of $p : E \rightarrow B$, which is a $C(B)$ -module.

The functor Γ .

- Recall that $\Gamma(E)$ is the set of sections of $p : E \rightarrow B$, which is a $C(B)$ -module.
- $\Gamma : \mathbf{Vect}_B \rightarrow \mathbf{Mod}_{C(B)}$ is a functor, with $F(f) : s(x) \mapsto fs(x)$.

The functor Γ .

- Recall that $\Gamma(E)$ is the set of sections of $p : E \rightarrow B$, which is a $C(B)$ -module.
- $\Gamma : \mathbf{Vect}_B \rightarrow \mathbf{Mod}_{C(B)}$ is a functor, with $F(f) : s(x) \mapsto fs(x)$.
- Γ respects the finite additive structures of \mathbf{Vect}_B and $\mathbf{Mod}_{C(B)}$:
 $\Gamma(E_1 \oplus E_2) = \Gamma(E_1) \oplus \Gamma(E_2)$.

The functor Γ .

- Recall that $\Gamma(E)$ is the set of sections of $p : E \rightarrow B$, which is a $C(B)$ -module.
- $\Gamma : \mathbf{Vect}_B \rightarrow \mathbf{Mod}_{C(B)}$ is a functor, with $F(f) : s(x) \mapsto fs(x)$.
- Γ respects the finite additive structures of \mathbf{Vect}_B and $\mathbf{Mod}_{C(B)}$:
 $\Gamma(E_1 \oplus E_2) = \Gamma(E_1) \oplus \Gamma(E_2)$.
- Quick computation: $\Gamma(B \times K^n) = \Gamma(\bigoplus_{i=1}^n B \times K) = \bigoplus_{i=1}^n \Gamma(B \times K) = C(B)^n$.

The Serre-Swan theorem.

- A projective $C(B)$ -module M is such that $M \oplus N = C(B)^n$.

The Serre-Swan theorem.

- A projective $C(B)$ -module M is such that $M \oplus N = C(B)^n$.
- (*Theorem 1*) If B is normal, then $\Gamma : \text{mor}(E_1, E_2) \rightarrow \text{mor}(\Gamma(E_1), \Gamma(E_2))$ is an isomorphism on sets of morphisms.

The Serre-Swan theorem.

- A projective $C(B)$ -module M is such that $M \oplus N = C(B)^n$.
- (*Theorem 1*) If B is normal, then $\Gamma : \text{mor}(E_1, E_2) \rightarrow \text{mor}(\Gamma(E_1), \Gamma(E_2))$ is an isomorphism on sets of morphisms.
- (*Theorem 2*) If B is compact Hausdorff, then Γ is an isomorphism onto the subcategory of finitely generated projective modules.

The Serre-Swan theorem.

- A projective $C(B)$ -module M is such that $M \oplus N = C(B)^n$.
- (*Theorem 1*) If B is normal, then $\Gamma : \text{mor}(E_1, E_2) \rightarrow \text{mor}(\Gamma(E_1), \Gamma(E_2))$ is an isomorphism on sets of morphisms.
- (*Theorem 2*) If B is compact Hausdorff, then Γ is an isomorphism onto the subcategory of finitely generated projective modules.
- Conclusion: both perspectives study essentially the same thing.

The Serre-Swan theorem I ($\text{mor}(E_1, E_2) \cong \text{mor}(\Gamma(E_1), \Gamma(E_2))$).

- Injectivity: if $\Gamma(f), \Gamma(g)$ coincide on sections $s : B \rightarrow E$, we want to show that f, g coincide on vectors $y \in E$.

The Serre-Swan theorem I ($\text{mor}(E_1, E_2) \cong \text{mor}(\Gamma(E_1), \Gamma(E_2))$).

- Injectivity: if $\Gamma(f), \Gamma(g)$ coincide on sections $s : B \rightarrow E$, we want to show that f, g coincide on vectors $y \in E$.

1 *Idea.* Construct a section s such that $s(p(y)) = y$.

The Serre-Swan theorem I ($\text{mor}(E_1, E_2) \cong \text{mor}(\Gamma(E_1), \Gamma(E_2))$).

- Injectivity: if $\Gamma(f), \Gamma(g)$ coincide on sections $s : B \rightarrow E$, we want to show that f, g coincide on vectors $y \in E$.
 - 1 *Idea.* Construct a section s such that $s(p(y)) = y$.
 - 2 Construct s locally.

The Serre-Swan theorem I ($\text{mor}(E_1, E_2) \cong \text{mor}(\Gamma(E_1), \Gamma(E_2))$).

- Injectivity: if $\Gamma(f), \Gamma(g)$ coincide on sections $s : B \rightarrow E$, we want to show that f, g coincide on vectors $y \in E$.
 - 1 *Idea.* Construct a section s such that $s(p(y)) = y$.
 - 2 Construct s locally.
 - 3 use a Urysohn function to extend s to all of E .

The Serre-Swan theorem I ($\text{mor}(E_1, E_2) \cong \text{mor}(\Gamma(E_1), \Gamma(E_2))$).

- Injectivity: if $\Gamma(f), \Gamma(g)$ coincide on sections $s : B \rightarrow E$, we want to show that f, g coincide on vectors $y \in E$.
 - 1 *Idea.* Construct a section s such that $s(p(y)) = y$.
 - 2 Construct s locally.
 - 3 use a Urysohn function to extend s to all of E .
- Surjectivity: for $F : \Gamma(E_1) \rightarrow \Gamma(E_2)$, find f such that $\Gamma(f) = F$.

The Serre-Swan theorem I ($\text{mor}(E_1, E_2) \cong \text{mor}(\Gamma(E_1), \Gamma(E_2))$).

- Injectivity: if $\Gamma(f), \Gamma(g)$ coincide on sections $s : B \rightarrow E$, we want to show that f, g coincide on vectors $y \in E$.
 - 1 *Idea.* Construct a section s such that $s(p(y)) = y$.
 - 2 Construct s locally.
 - 3 use a Urysohn function to extend s to all of E .
- Surjectivity: for $F : \Gamma(E_1) \rightarrow \Gamma(E_2)$, find f such that $\Gamma(f) = F$.
 - 1 *Idea.* $\Gamma(E_1)_x$: the submodule of $\Gamma(E_1)$ of sections vanishing at x .

The Serre-Swan theorem I ($\text{mor}(E_1, E_2) \cong \text{mor}(\Gamma(E_1), \Gamma(E_2))$).

- Injectivity: if $\Gamma(f), \Gamma(g)$ coincide on sections $s : B \rightarrow E$, we want to show that f, g coincide on vectors $y \in E$.
 - 1 *Idea.* Construct a section s such that $s(p(y)) = y$.
 - 2 Construct s locally.
 - 3 use a Urysohn function to extend s to all of E .
- Surjectivity: for $F : \Gamma(E_1) \rightarrow \Gamma(E_2)$, find f such that $\Gamma(f) = F$.
 - 1 *Idea.* $\Gamma(E_1)_x$: the submodule of $\Gamma(E_1)$ of sections vanishing at x .
 - 2 F induces maps $f_x : \Gamma(E_1)/\Gamma(E_1)_x \rightarrow \Gamma(E_2)/\Gamma(E_2)_x$ at each x .

The Serre-Swan theorem I ($\text{mor}(E_1, E_2) \cong \text{mor}(\Gamma(E_1), \Gamma(E_2))$).

- Injectivity: if $\Gamma(f), \Gamma(g)$ coincide on sections $s : B \rightarrow E$, we want to show that f, g coincide on vectors $y \in E$.
 - 1 *Idea.* Construct a section s such that $s(p(y)) = y$.
 - 2 Construct s locally.
 - 3 use a Urysohn function to extend s to all of E .
- Surjectivity: for $F : \Gamma(E_1) \rightarrow \Gamma(E_2)$, find f such that $\Gamma(f) = F$.
 - 1 *Idea.* $\Gamma(E_1)_x$: the submodule of $\Gamma(E_1)$ of sections vanishing at x .
 - 2 F induces maps $f_x : \Gamma(E_1)/\Gamma(E_1)_x \rightarrow \Gamma(E_2)/\Gamma(E_2)_x$ at each x .
 - 3 $\Gamma(E_1)/\Gamma(E_1)_x \cong p_1^{-1}(x)$.

The Serre-Swan theorem II ($\text{Vect}_B \cong \text{Mod}_{C(B)}^{\text{f.g.p.}}$).

- Let $p : E \rightarrow B$ be a vector bundle over a compact Hausdorff > paracompact, normal) space.

The Serre-Swan theorem II ($\mathrm{Vect}_B \cong \mathrm{Mod}_{C(B)}^{\mathrm{f.g.p.}}$).

- Let $p : E \rightarrow B$ be a vector bundle over a compact Hausdorff > paracompact, normal) space.
- (\subseteq) .

The Serre-Swan theorem II ($\text{Vect}_B \cong \text{Mod}_{C(B)}^{\text{f.g.p.}}$).

- Let $p : E \rightarrow B$ be a vector bundle over a compact Hausdorff > paracompact, normal) space.
- (\subseteq) .
 - 1 Compactness $\implies \exists$ surjection $f : B \times K^n \rightarrow E$, so

$$\Gamma(E) \oplus \Gamma(\ker f) = \Gamma(B \times K^n) = C(B)^n.$$

The Serre-Swan theorem II ($\text{Vect}_B \cong \text{Mod}_{C(B)}^{\text{f.g.p.}}$).

- Let $p : E \rightarrow B$ be a vector bundle over a compact Hausdorff > paracompact, normal) space.
- (\subseteq) .

1 Compactness $\implies \exists$ surjection $f : B \times K^n \rightarrow E$, so

$$\Gamma(E) \oplus \Gamma(\ker f) = \Gamma(B \times K^n) = C(B)^n.$$

2 Local basis around each $x \in E \implies s_1, \dots, s_n \in \Gamma(E)$ spans each fiber.

The Serre-Swan theorem II ($\text{Vect}_B \cong \text{Mod}_{C(B)}^{\text{f.g.p.}}$).

- Let $p : E \rightarrow B$ be a vector bundle over a compact Hausdorff > paracompact, normal) space.
- (\subseteq) .

1 Compactness $\implies \exists$ surjection $f : B \times K^n \rightarrow E$, so

$$\Gamma(E) \oplus \Gamma(\ker f) = \Gamma(B \times K^n) = C(B)^n.$$

2 Local basis around each $x \in E \implies s_1, \dots, s_n \in \Gamma(E)$ spans each fiber.

- (\supseteq) .

The Serre-Swan theorem II ($\text{Vect}_B \cong \text{Mod}_{C(B)}^{\text{f.g.p.}}$).

- Let $p : E \rightarrow B$ be a vector bundle over a compact Hausdorff > paracompact, normal) space.
- (\subseteq) .

1 Compactness $\implies \exists$ surjection $f : B \times K^n \rightarrow E$, so

$$\Gamma(E) \oplus \Gamma(\ker f) = \Gamma(B \times K^n) = C(B)^n.$$

2 Local basis around each $x \in E \implies s_1, \dots, s_n \in \Gamma(E)$ spans each fiber.

- (\supseteq) .

1 Take f.g.p M which is a summand of $C(B)^n$.

The Serre-Swan theorem II ($\text{Vect}_B \cong \text{Mod}_{C(B)}^{\text{f.g.p.}}$).

- Let $p : E \rightarrow B$ be a vector bundle over a compact Hausdorff > paracompact, normal) space.
- (\subseteq) .

1 Compactness $\implies \exists$ surjection $f : B \times K^n \rightarrow E$, so

$$\Gamma(E) \oplus \Gamma(\ker f) = \Gamma(B \times K^n) = C(B)^n.$$

2 Local basis around each $x \in E \implies s_1, \dots, s_n \in \Gamma(E)$ spans each fiber.

- (\supseteq) .

1 Take f.g.p M which is a summand of $C(B)^n$.

2 Projection $F : C(B)^n \rightarrow C(B)^n$ onto M is uniquely induced by endomorphism f on $B \times K^n$.

The Serre-Swan theorem II ($\text{Vect}_B \cong \text{Mod}_{C(B)}^{\text{f.g.p.}}$).

- Let $p : E \rightarrow B$ be a vector bundle over a compact Hausdorff > paracompact, normal) space.
- (\subseteq) .

1 Compactness $\implies \exists$ surjection $f : B \times K^n \rightarrow E$, so

$$\Gamma(E) \oplus \Gamma(\ker f) = \Gamma(B \times K^n) = C(B)^n.$$

2 Local basis around each $x \in E \implies s_1, \dots, s_n \in \Gamma(E)$ spans each fiber.

- (\supseteq) .

1 Take f.g.p M which is a summand of $C(B)^n$.

2 Projection $F : C(B)^n \rightarrow C(B)^n$ onto M is uniquely induced by endomorphism f on $B \times K^n$.

3 Show $\text{im } f$ is a subbundle, implying

$$\Gamma(f : B \times K^n \rightarrow \text{im } f) = F : C(B)^n \rightarrow \Gamma(\text{im } f).$$