

# How an Enumerative Combinatorialist Might Solve a PDE

Robbert Liu

University of Toronto

April 24, 2024

# What is enumerative combinatorics

The job of an enumerative combinatorialist is to construct a bridge between the world of discrete geometry and algebra.

# What is enumerative combinatorics

The job of an enumerative combinatorialist is to construct a bridge between the world of discrete geometry and algebra.

$$\begin{array}{ccc} F([n]) & \xrightarrow{|\cdot|} & F(n) \\ \uparrow \cong & & \parallel \\ G([n]) & \xrightarrow{|\cdot|} & G(n) \end{array}$$

Note on Notation:

- $[n] = \{1, \dots, n\}$ .
- $F(X)$  class of structures on a set  $X$ .
- $F(x)$  quantity, func. of  $x$ ,  $|F(x)| = F(x)$ .

# What is enumerative combinatorics

The job of an enumerative combinatorialist is to construct a bridge between the world of discrete geometry and algebra.

$$\begin{array}{ccc} F([n]) & \xrightarrow{|\cdot|} & F(n) \\ \uparrow \cong & & \parallel \\ G([n]) & \xrightarrow{|\cdot|} & G(n) \end{array}$$

Note on Notation:

- $[n] = \{1, \dots, n\}$ .
- $F(X)$  class of structures on a set  $X$ .
- $F(x)$  quantity, func. of  $x$ ,  $|F(x)| = F(x)$ .

Develop a geometric way of treating arithmetic (integral) identities.

# What is enumerative combinatorics

The job of an enumerative combinatorialist is to construct a bridge between the world of discrete geometry and algebra.

$$\begin{array}{ccc} F([n]) & \xrightarrow{|\cdot|} & F(n) \\ \uparrow \cong & & \parallel \\ G([n]) & \xrightarrow{|\cdot|} & G(n) \end{array}$$

Note on Notation:

- $[n] = \{1, \dots, n\}$ .
- $F(X)$  class of structures on a set  $X$ .
- $F(x)$  quantity, func. of  $x$ ,  $|F(x)| = F(x)$ .

Develop a geometric way of treating arithmetic (integral) identities.

i.e. Study algebraic operations on classes of discrete geometric structures equipped with an enumeration homomorphism.



# Counting monotonic paths, the first way

Consider

$$4^n = \sum_{k=0}^n \binom{2k}{k} \binom{2(n-k)}{n-k}$$

Remember:  $\binom{n}{k} = |\{k\text{-subsets of } [n]\}|$ .

# Counting monotonic paths, the first way

Consider

$$4^n = \sum_{k=0}^n \binom{2k}{k} \binom{2(n-k)}{n-k}$$

Remember:  $\binom{n}{k} = |\{k\text{-subsets of } [n]\}|$ .

One way to prove:



# Counting monotonic paths, the first way

Consider

$$4^n = \sum_{k=0}^n \binom{2k}{k} \binom{2(n-k)}{n-k}$$

Remember:  $\binom{n}{k} = |\{k\text{-subsets of } [n]\}|$ .

One way to prove:

- LHS:  $4^n = 2^{2n}$  enumerates all monotonic paths (travelling Up or Right) of length  $2n$

# Counting monotonic paths, the first way

Consider

$$4^n = \sum_{k=0}^n \binom{2k}{k} \binom{2(n-k)}{n-k}$$

Remember:  $\binom{n}{k} = |\{k\text{-subsets of } [n]\}|$ .

One way to prove:

- LHS:  $4^n = 2^{2n}$  enumerates all monotonic paths (travelling Up or Right) of length  $2n$
- RHS: A path, after  $2k$  steps, intersects the diagonal for the last time. (a unique  $k$  exists).

# Counting monotonic paths, the first way

Consider

$$4^n = \sum_{k=0}^n \binom{2k}{k} \binom{2(n-k)}{n-k}$$

Remember:  $\binom{n}{k} = |\{k\text{-subsets of } [n]\}|$ .

One way to prove:

- LHS:  $4^n = 2^{2n}$  enumerates all monotonic paths (travelling Up or Right) of length  $2n$
- RHS: A path, after  $2k$  steps, intersects the diagonal for the last time. (a unique  $k$  exists).
- $\binom{2k}{k}$  counts # choices for initial segment,  $\binom{2(n-k)}{n-k}$  counts # choices for remaining steps.

## The first way (Cont.)





- LHS:  $4^n = 2^{2n}$  enumerates all monotonic paths (travelling Up or Right) of length  $2n$
- RHS: A path, after  $2k$  steps, intersects the diagonal for the last time. (a unique  $k$  exists).
- $\binom{2k}{k}$  counts # choices for initial segment,  $\binom{2(n-k)}{n-k}$  counts # choices for remaining steps.

## The first way (Cont.)

- LHS:  $4^n = 2^{2n}$  enumerates all monotonic paths (travelling Up or Right) of length  $2n$
- RHS: A path, after  $2k$  steps, intersects the diagonal for the last time. (a unique  $k$  exists).
- $\binom{2k}{k}$  counts # choices for initial segment,  $\binom{2(n-k)}{n-k}$  counts # choices for remaining steps.

### Example

$$2^{20} = \sum_{k=0}^{10} \binom{2k}{k} \binom{2(n-k)}{n-k}, \quad k = 5 : \binom{10}{5} \binom{10}{5}$$



# Counting monotonic paths, the second way

Consider

$$(1 - 4x)^{-\frac{1}{2}} = \sum_{k=0}^{\infty} \binom{-\frac{1}{2}}{k} (-4x)^k$$

## Counting monotonic paths, the second way

Consider

$$\begin{aligned}(1 - 4x)^{-\frac{1}{2}} &= \sum_{k=0}^{\infty} \binom{-\frac{1}{2}}{k} (-4x)^k \\ &= \sum_{k=0}^{\infty} \frac{(-\frac{1}{2}) \cdots (-\frac{1}{2} - k + 1)}{k!} (-4)^k x^k\end{aligned}$$

# Counting monotonic paths, the second way

Consider

$$\begin{aligned}(1 - 4x)^{-\frac{1}{2}} &= \sum_{k=0}^{\infty} \binom{-\frac{1}{2}}{k} (-4x)^k \\ &= \sum_{k=0}^{\infty} \frac{(-\frac{1}{2}) \dots (-\frac{1}{2} - k + 1)}{k!} (-4)^k x^k \\ &\dots \\ &= \sum_{k=0}^{\infty} \binom{2k}{k} x^k\end{aligned}$$



# Counting monotonic paths, the second way

Consider

$$\begin{aligned}(1 - 4x)^{-\frac{1}{2}} &= \sum_{k=0}^{\infty} \binom{-\frac{1}{2}}{k} (-4x)^k \\ &= \sum_{k=0}^{\infty} \frac{(-\frac{1}{2}) \dots (-\frac{1}{2} - k + 1)}{k!} (-4)^k x^k \\ &\dots \\ &= \sum_{k=0}^{\infty} \binom{2k}{k} x^k\end{aligned}$$

Squaring both sides (RHS is a convolution)

$$\frac{1}{1 - 4x} = \sum_{n=0}^{\infty} 4^n x^n = \sum_{n=0}^{\infty} \left[ \sum_{k=0}^n \binom{2k}{k} \binom{2(n-k)}{n-k} \right] x^n$$

# Counting monotonic paths, the second way

Consider

$$\begin{aligned}(1 - 4x)^{-\frac{1}{2}} &= \sum_{k=0}^{\infty} \binom{-\frac{1}{2}}{k} (-4x)^k \\ &= \sum_{k=0}^{\infty} \frac{(-\frac{1}{2}) \dots (-\frac{1}{2} - k + 1)}{k!} (-4)^k x^k \\ &\dots \\ &= \sum_{k=0}^{\infty} \binom{2k}{k} x^k\end{aligned}$$

Squaring both sides (RHS is a convolution)

$$\frac{1}{1 - 4x} = \sum_{n=0}^{\infty} 4^n x^n = \sum_{n=0}^{\infty} \left[ \sum_{k=0}^n \binom{2k}{k} \binom{2(n-k)}{n-k} \right] x^n$$

Enumerators destroyed!

# Higher enumerable discrete structures

Wrong scheme:

$$\begin{array}{ccc} F([n]) & \xrightarrow{\quad |\cdot| \quad} & 4^n \\ \updownarrow \cong & & \parallel \\ \bigsqcup G([k]) \times G'([n-k]) & \xrightarrow{\quad |\cdot| \quad} & \sum \binom{2k}{k} \binom{2(n-k)}{n-k} \end{array}$$

# Higher enumerable discrete structures

Wrong scheme:

$$\begin{array}{ccc}
 F([n]) & \xrightarrow{|\cdot|} & 4^n \\
 \updownarrow \cong & & \parallel \\
 \bigsqcup G([k]) \times G'([n-k]) & \xrightarrow{|\cdot|} & \sum \binom{2k}{k} \binom{2(n-k)}{n-k}
 \end{array}$$

Right scheme:

$$\begin{array}{ccc}
 F[X] & \xrightarrow{|\cdot|} & \frac{1}{\sqrt{1-4x}} \\
 \updownarrow \cong & & \parallel \\
 G \cdot G'[X] & \xrightarrow{|\cdot|} & \sum_n \left[ \sum_k \binom{2k}{k} \binom{2(n-k)}{n-k} \right] x^n
 \end{array}$$

# Overview, higher enumerable discrete structures (Cont.)

Goal: Find a model  $\mathbb{S}(X)$  for the algebra of all  $F[X]$ .

# Overview, higher enumerable discrete structures (Cont.)

Goal: Find a model  $\mathbb{S}(X)$  for the algebra of all  $F[X]$ .

- Formal  $+$ ,  $\cdot$ ,  $\circ$ ,  $\frac{d}{dX}$  operations.

# Overview, higher enumerable discrete structures (Cont.)

Goal: Find a model  $\mathbb{S}(X)$  for the algebra of all  $F[X]$ .

- Formal  $+$ ,  $\cdot$ ,  $\circ$ ,  $\frac{d}{dX}$  operations.
- Enumeration  $\mathbb{S}(X) \rightarrow k[[x]]$  must respect operations, i.e. algebra homomorphism.

# Overview, higher enumerable discrete structures (Cont.)

Goal: Find a model  $\mathbb{S}(X)$  for the algebra of all  $F[X]$ .

- Formal  $+$ ,  $\cdot$ ,  $\circ$ ,  $\frac{d}{dX}$  operations.
- Enumeration  $\mathbb{S}(X) \rightarrow k[[x]]$  must respect operations, i.e. algebra homomorphism.
- Hence:  $\mathbb{S}(X)$  behaves like a ring of smooth functions.



# Overview, higher enumerable discrete structures (Cont.)

Goal: Find a model  $\mathbb{S}(X)$  for the algebra of all  $F[X]$ .

- Formal  $+$ ,  $\cdot$ ,  $\circ$ ,  $\frac{d}{dX}$  operations.
- Enumeration  $\mathbb{S}(X) \rightarrow k[[x]]$  must respect operations, i.e. algebra homomorphism.
- Hence:  $\mathbb{S}(X)$  behaves like a ring of smooth functions.
- i.e. Leibniz identity and chain rule.

# Overview, higher enumerable discrete structures (Cont.)

Goal: Find a model  $\mathbb{S}(X)$  for the algebra of all  $F[X]$ .

- Formal  $+$ ,  $\cdot$ ,  $\circ$ ,  $\frac{d}{dX}$  operations.
- Enumeration  $\mathbb{S}(X) \rightarrow k[[x]]$  must respect operations, i.e. algebra homomorphism.
- Hence:  $\mathbb{S}(X)$  behaves like a ring of smooth functions.
- i.e. Leibniz identity and chain rule.
- Generalize to multiple variables, i.e.  $\mathbb{S}(X, Y, Z)$ .

# The algebra of species

## Definition

A (combinatorial) species  $F[X]$  has the following data,

# The algebra of species

## Definition

A (combinatorial) species  $F[X]$  has the following data,

- For every finite set  $U$ , we get a finite set  $F[U]$  of “ $F$ -structures” on the set  $U$ .

# The algebra of species

## Definition

A (combinatorial) species  $F[X]$  has the following data,

- For every finite set  $U$ , we get a finite set  $F[U]$  of “ $F$ -structures” on the set  $U$ .
- For every bijection  $f : U \rightarrow V$ , we get a transport function  $F[U] \rightarrow F[V]$ .

# The algebra of species

## Definition

A (combinatorial) species  $F[X]$  has the following data,

- For every finite set  $U$ , we get a finite set  $F[U]$  of “ $F$ -structures” on the set  $U$ .
- For every bijection  $f : U \rightarrow V$ , we get a transport function  $F[U] \rightarrow F[V]$ .

Using  $f$ , we can relabel the points of a  $F$ -structure on  $U$ , to get a structure on  $V$ .

# The algebra of species

## Definition

A (combinatorial) species  $F[X]$  has the following data,

- For every finite set  $U$ , we get a finite set  $F[U]$  of “ $F$ -structures” on the set  $U$ .
- For every bijection  $f : U \rightarrow V$ , we get a transport function  $F[U] \rightarrow F[V]$ .

Using  $f$ , we can relabel the points of a  $F$ -structure on  $U$ , to get a structure on  $V$ .

## Definition 2

A species is a functor  $F[X]: \text{Core}(\text{FinSet}) \rightarrow \text{Core}(\text{FinSet})$ .

# Some examples



# Some examples

Permutations  $\mathcal{S}[X]$ , Linear orders  $\mathcal{L}[X]$ , Simple graphs  $\mathcal{G}[X]$

# Some examples

Permutations  $\mathcal{S}[X]$ , Linear orders  $\mathcal{L}[X]$ , Simple graphs  $\mathcal{G}[X]$

## Transport function


Consider  $f : \{1, 2, 3, 4, 5\} \rightarrow \{a, b, c, d, e\}$ .

# Some examples

Permutations  $\mathcal{S}[X]$ , Linear orders  $\mathcal{L}[X]$ , Simple graphs  $\mathcal{G}[X]$

## Transport function

Consider  $f : \{1, 2, 3, 4, 5\} \rightarrow \{a, b, c, d, e\}$ .



## More examples

## More examples

- Sets:  $E[U] = *$  for all  $U$ .

## More examples

- Sets:  $E[U] = *$  for all  $U$ .
- Singletons:  $X[*] = *$ , otherwise  $X[U] = \emptyset$ .

## More examples

- Sets:  $E[U] = *$  for all  $U$ .
- Singletons:  $X[*] = *$ , otherwise  $X[U] = \emptyset$ .
- $n$ -sets:  $X^n[U] = *$  if  $|U| = n$ , otherwise  $X^n[X] = \emptyset$ .

## More examples

- Sets:  $E[U] = *$  for all  $U$ .
- Singletons:  $X[*] = *$ , otherwise  $X[U] = \emptyset$ .
- $n$ -sets:  $X^n[U] = *$  if  $|U| = n$ , otherwise  $X^n[X] = \emptyset$ .
- $1[\emptyset] = *$



## More examples

- Sets:  $E[U] = *$  for all  $U$ .
- Singletons:  $X[*] = *$ , otherwise  $X[U] = \emptyset$ .
- $n$ -sets:  $X^n[U] = *$  if  $|U| = n$ , otherwise  $X^n[X] = \emptyset$ .
- $1[\emptyset] = *$
- The 0-species  $0(U) = \emptyset$ .

## More examples

- Sets:  $E[U] = *$  for all  $U$ .
- Singletons:  $X[*] = *$ , otherwise  $X[U] = \emptyset$ .
- $n$ -sets:  $X^n[U] = *$  if  $|U| = n$ , otherwise  $X^n[X] = \emptyset$ .
- $1[\emptyset] = *$
- The 0-species  $0(U) = \emptyset$ .

### Remark (Enumeration homomorphism)

If  $f : U \rightarrow V$  is a bijection, then  $f_F$  is a bijection of  $F$ -structures.  
Thus,  $|F[n]|$  is well-defined.

## More examples

- Sets:  $E[U] = *$  for all  $U$ .
- Singletons:  $X[*] = *$ , otherwise  $X[U] = \emptyset$ .
- $n$ -sets:  $X^n[U] = *$  if  $|U| = n$ , otherwise  $X^n[X] = \emptyset$ .
- $1[\emptyset] = *$
- The 0-species  $0(U) = \emptyset$ .

### Remark (Enumeration homomorphism)

If  $f : U \rightarrow V$  is a bijection, then  $f_F$  is a bijection of  $F$ -structures.  
Thus,  $|F[n]|$  is well-defined. Define:

$$F(x) := \sum_n |F[n]| \frac{x^n}{n!} \quad (1)$$

## More examples

- Sets:  $E[U] = *$  for all  $U$ .
- Singletons:  $X[*] = *$ , otherwise  $X[U] = \emptyset$ .
- $n$ -sets:  $X^n[U] = *$  if  $|U| = n$ , otherwise  $X^n[X] = \emptyset$ .
- $1[\emptyset] = *$
- The 0-species  $0(U) = \emptyset$ .

### Remark (Enumeration homomorphism)

If  $f : U \rightarrow V$  is a bijection, then  $f_F$  is a bijection of  $F$ -structures.  
Thus,  $|F[n]|$  is well-defined. Define:

$$F(x) := \sum_n |F[n]| \frac{x^n}{n!} \quad (1)$$

Immediately,  $E(x) = e^x$ ,  $X^n(x) = \frac{x^n}{n!}$ ,  $1(x) = 1$ ,  $0(x) = 0$ .

# Q&A (Interlude)

# Q&A (Interlude)

Q: Why every finite set?

## Q&A (Interlude)

Q: Why every finite set?

- There is not canonical choice of  $[n]$ .

## Q&A (Interlude)

Q: Why every finite set?

- There is not canonical choice of  $[n]$ .
- $G[\{1, 2, 4\}]$  and  $G[\{1, 2, 3\}]$  embed differently in  $G \in G[4]$ .



# Q&A (Interlude)

Q: Why every finite set?

- There is not canonical choice of  $[n]$ .
- $G[\{1, 2, 4\}]$  and  $G[\{1, 2, 3\}]$  embed differently in  $G \in G[4]$ .

Q: Did we do anything new?

## Q&A (Interlude)

Q: Why every finite set?

- There is not canonical choice of  $[n]$ .
- $G[\{1, 2, 4\}]$  and  $G[\{1, 2, 3\}]$  embed differently in  $G \in G[4]$ .

Q: Did we do anything new?

Yes! The algebra of species is *richer* than the algebra of power series.

## Q&A (Interlude)

Q: Why every finite set?

- There is not canonical choice of  $[n]$ .
- $G[\{1, 2, 4\}]$  and  $G[\{1, 2, 3\}]$  embed differently in  $G \in G[4]$ .

Q: Did we do anything new?

Yes! The algebra of species is *richer* than the algebra of power series.

- $\mathcal{L}(x) = \mathcal{S}(x) = \frac{1}{1-x} = \sum n! \frac{x^n}{n!}.$

## Q&A (Interlude)

Q: Why every finite set?

- There is not canonical choice of  $[n]$ .
- $G[\{1, 2, 4\}]$  and  $G[\{1, 2, 3\}]$  embed differently in  $G \in G[4]$ .

Q: Did we do anything new?

Yes! The algebra of species is *richer* than the algebra of power series.

- $\mathcal{L}(x) = \mathcal{S}(x) = \frac{1}{1-x} = \sum n! \frac{x^n}{n!}$ .
- $\mathcal{L}[X]$  and  $\mathcal{S}[X]$  are NOT isomorphic.

## Q&A (Interlude)

Q: Why every finite set?

- There is not canonical choice of  $[n]$ .
- $G[\{1, 2, 4\}]$  and  $G[\{1, 2, 3\}]$  embed differently in  $G \in G[4]$ .

Q: Did we do anything new?

Yes! The algebra of species is *richer* than the algebra of power series.

- $\mathcal{L}(x) = \mathcal{S}(x) = \frac{1}{1-x} = \sum n! \frac{x^n}{n!}$ .
- $\mathcal{L}[X]$  and  $\mathcal{S}[X]$  are NOT isomorphic.
- Transport functions of  $\sigma : U \rightarrow U$  have different automorphism types, fixed points:

$$\sigma := (12) \in S_5$$

$$\sigma_S(12)(345) = (12)(345).$$

$$\sigma_S(21453) = (12453) \neq (21453).$$

# Operations on species

Let  $F[X]$ ,  $G[X]$  be species.

## Operations on species

Let  $F[X], G[X]$  be species.

- Addition: Define  $(F + G)[U] := F[U] \sqcup G[U]$  and  $f_{F+G} = f_F \sqcup f_G$ .

# Operations on species

Let  $F[X], G[X]$  be species.

- Addition: Define  $(F + G)[U] := F[U] \sqcup G[U]$  and  $f_{F+G} = f_F \sqcup f_G$ .
- An  $(F + G)$ -structure on  $U$  is either an  $F$ -structure or a  $G$ -structure on  $U$ .



# Operations on species

Let  $F[X], G[X]$  be species.

- Addition: Define  $(F + G)[U] := F[U] \sqcup G[U]$  and  $f_{F+G} = f_F \sqcup f_G$ .
- An  $(F + G)$ -structure on  $U$  is either an  $F$ -structure or a  $G$ -structure on  $U$ .
- Multiplication/Convolution: Define  $(F \cdot G)[U] := \sqcup_{U=V \sqcup W} F[V] \times G[W]$ .

# Operations on species

Let  $F[X], G[X]$  be species.

- Addition: Define  $(F + G)[U] := F[U] \sqcup G[U]$  and  $f_{F+G} = f_F \sqcup f_G$ .
- An  $(F + G)$ -structure on  $U$  is either an  $F$ -structure or a  $G$ -structure on  $U$ .
- Multiplication/Convolution: Define  $(F \cdot G)[U] := \sqcup_{U=V \sqcup W} F[V] \times G[W]$ .
- (Example)

## Operations on species

Let  $F[X], G[X]$  be species.

- Addition: Define  $(F + G)[U] := F[U] \sqcup G[U]$  and  $f_{F+G} = f_F \sqcup f_G$ .
- An  $(F + G)$ -structure on  $U$  is either an  $F$ -structure or a  $G$ -structure on  $U$ .
- Multiplication/Convolution: Define  $(F \cdot G)[U] := \sqcup_{U=V \sqcup W} F[V] \times G[W]$ .
- (Example)
- Notice that

$$\begin{aligned}(F \cdot G)(x) &= \sum_n \sum_{[n] \cong [k] \sqcup [k']} |F[k]| \cdot |G[k']| \frac{x^n}{n!} \\&= \sum_n \sum_{k=0}^n \binom{n}{k} |F[k]| \cdot |G[n-k]| \frac{x^n}{n!} \\&= \sum_n \sum_{k=0}^n \frac{|F[k]|}{k!} \cdot \frac{|G[n-k]|}{(n-k)!} x^n = F(x) \cdot G(x)\end{aligned}$$

# Differentiation

# Differentiation

- Recall that  $\frac{d}{dx} \sum F_n \frac{x^n}{n!} = \sum F_{n+1} \frac{x^n}{n!}$ .

# Differentiation

- Recall that  $\frac{d}{dx} \sum F_n \frac{x^n}{n!} = \sum F_{n+1} \frac{x^n}{n!}$ .
- Differentiation: Define  $\frac{d}{dX} F[U] = F[U \sqcup *]$

# Differentiation

- Recall that  $\frac{d}{dx} \sum F_n \frac{x^n}{n!} = \sum F_{n+1} \frac{x^n}{n!}$ .
- Differentiation: Define  $\frac{d}{dX} F[U] = F[U \sqcup *]$
- (Example)

# Differentiation

- Recall that  $\frac{d}{dx} \sum F_n \frac{x^n}{n!} = \sum F_{n+1} \frac{x^n}{n!}$ .
- Differentiation: Define  $\frac{d}{dX} F[U] = F[U \sqcup *]$
- (Example)
- (Sketch of Leibniz identity)



# Composition

# Composition

- If  $G(0) = 0$ , then  $F(G(x)) := \sum F_n \frac{G(x)^n}{n!}$  is well-defined.

# Composition

- If  $G(0) = 0$ , then  $F(G(x)) := \sum F_n \frac{G(x)^n}{n!}$  is well-defined.

## Theorem

The following formula holds:

$$F(G(x)) = \sum_n \sum_{\sum \{d_1, \dots, d_k\} = n} \binom{n}{d_1, \dots, d_k} F_k \prod_{i=1}^k G_{d_i}.$$

# Composition

- If  $G(0) = 0$ , then  $F(G(x)) := \sum F_n \frac{G(x)^n}{n!}$  is well-defined.

## Theorem

The following formula holds:

$$F(G(x)) = \sum_n \sum_{\sum \{d_1, \dots, d_k\} = n} \binom{n}{d_1, \dots, d_k} F_k \prod_{i=1}^k G_{d_i}.$$

- Composition: Define

$$(F \circ G)[U] = \bigsqcup_{D_1 \sqcup \dots \sqcup D_k = U} F[\{D_1, \dots, D_k\}] \times \prod_i^k G[D_i]$$

# Composition

- If  $G(0) = 0$ , then  $F(G(x)) := \sum F_n \frac{G(x)^n}{n!}$  is well-defined.

## Theorem

The following formula holds:

$$F(G(x)) = \sum_n \sum_{\sum \{d_1, \dots, d_k\} = n} \binom{n}{d_1, \dots, d_k} F_k \prod_{i=1}^k G_{d_i}.$$

- Composition: Define

$$(F \circ G)[U] = \bigsqcup_{D_1 \sqcup \dots \sqcup D_k = U} F[\{D_1, \dots, D_k\}] \times \prod_{i=1}^k G[D_i]$$

- An  $(F \circ G)$ -structure on  $U$  is a partition  $U = D_1 \sqcup \dots \sqcup D_k$ , a  $G$  structure on every  $D_i$ , and an  $F$ -structure on  $\{D_1, \dots, D_k\}$

# Composition

- If  $G(0) = 0$ , then  $F(G(x)) := \sum F_n \frac{G(x)^n}{n!}$  is well-defined.

## Theorem

The following formula holds:

$$F(G(x)) = \sum_n \sum_{\sum \{d_1, \dots, d_k\} = n} \binom{n}{d_1, \dots, d_k} F_k \prod_{i=1}^k G_{d_i}.$$

- Composition: Define

$$(F \circ G)[U] = \bigsqcup_{D_1 \sqcup \dots \sqcup D_k = U} F[\{D_1, \dots, D_k\}] \times \prod_{i=1}^k G[D_i]$$

- An  $(F \circ G)$ -structure on  $U$  is a partition  $U = D_1 \sqcup \dots \sqcup D_k$ , a  $G$  structure on every  $D_i$ , and an  $F$ -structure on  $\{D_1, \dots, D_k\}$
- (Example)

# Composition

- If  $G(0) = 0$ , then  $F(G(x)) := \sum F_n \frac{G(x)^n}{n!}$  is well-defined.

## Theorem

The following formula holds:

$$F(G(x)) = \sum_n \sum_{\{d_1, \dots, d_k\} = n} \binom{n}{d_1, \dots, d_k} F_k \prod_{i=1}^k G_{d_i}.$$

- Composition: Define

$$(F \circ G)[U] = \bigsqcup_{D_1 \sqcup \dots \sqcup D_k = U} F[\{D_1, \dots, D_k\}] \times \prod_{i=1}^k G[D_i]$$

- An  $(F \circ G)$ -structure on  $U$  is a partition  $U = D_1 \sqcup \dots \sqcup D_k$ , a  $G$  structure on every  $D_i$ , and an  $F$ -structure on  $\{D_1, \dots, D_k\}$
- (Example)
- (Sketch of chain rule)

# Functional equations



# Functional equations

## Linear orders

$$\mathcal{L} = 1 + X \cdot \mathcal{L}$$

# Functional equations

Linear orders

$$\mathcal{L} = 1 + X \cdot \mathcal{L}$$

Connected species

$$F = E(F^c)$$

Binary rooted trees (Lawvere, Baez)

$$\mathcal{T} = 1 + X\mathcal{T}^2$$

# Multiple variables

## Definition (Multiset)

# Multiple variables

## Definition (Multiset)

- A multiset is a tuple  $(U_1, \dots, U_n)$  of sets.

# Multiple variables

## Definition (Multiset)

- A multiset is a tuple  $(U_1, \dots, U_n)$  of sets.
- A function/bijection of multisets  $(f_k) : (U_1, \dots, U_n) \rightarrow (V_1, \dots, V_n)$  is a tuple of functions/bijections  $f_k : U_k \rightarrow V_k$ .

# Multiple variables

## Definition (Multiset)

- A multiset is a tuple  $(U_1, \dots, U_n)$  of sets.
- A function/bijection of multisets  $(f_k) : (U_1, \dots, U_n) \rightarrow (V_1, \dots, V_n)$  is a tuple of functions/bijections  $f_k : U_k \rightarrow V_k$ .
- A subset of  $(U_1, \dots, U_n)$  is a tuple  $(V_1, \dots, V_n)$  such that  $V_k \subseteq U_k$ .

# Multiple variables

## Definition (Multiset)

- A multiset is a tuple  $(U_1, \dots, U_n)$  of sets.
- A function/bijection of multisets  $(f_k) : (U_1, \dots, U_n) \rightarrow (V_1, \dots, V_n)$  is a tuple of functions/bijections  $f_k : U_k \rightarrow V_k$ .
- A subset of  $(U_1, \dots, U_n)$  is a tuple  $(V_1, \dots, V_n)$  such that  $V_k \subseteq U_k$ .
- A partition of  $(U_1, \dots, U_n)$  is a collection  $\{U_1^i, \dots, U_n^i\}_i$  of subsets of  $(U_1, \dots, U_n)$  such that  $\bigcup_i U_k^i = U_k$ .

# Multiple variables

## Definition (Multiset)

- A multiset is a tuple  $(U_1, \dots, U_n)$  of sets.
- A function/bijection of multisets  $(f_k) : (U_1, \dots, U_n) \rightarrow (V_1, \dots, V_n)$  is a tuple of functions/bijections  $f_k : U_k \rightarrow V_k$ .
- A subset of  $(U_1, \dots, U_n)$  is a tuple  $(V_1, \dots, V_n)$  such that  $V_k \subseteq U_k$ .
- A partition of  $(U_1, \dots, U_n)$  is a collection  $\{U_1^i, \dots, U_n^i\}_i$  of subsets of  $(U_1, \dots, U_n)$  such that  $\bigcup_i U_k^i = U_k$ .

## Definition (Multisort species)

A multi-sort species  $F[X_1, \dots, X_n]$  has the following data,



# Multiple variables

## Definition (Multiset)

- A multiset is a tuple  $(U_1, \dots, U_n)$  of sets.
- A function/bijection of multisets  $(f_k) : (U_1, \dots, U_n) \rightarrow (V_1, \dots, V_n)$  is a tuple of functions/bijections  $f_k : U_k \rightarrow V_k$ .
- A subset of  $(U_1, \dots, U_n)$  is a tuple  $(V_1, \dots, V_n)$  such that  $V_k \subseteq U_k$ .
- A partition of  $(U_1, \dots, U_n)$  is a collection  $\{U_1^i, \dots, U_n^i\}_i$  of subsets of  $(U_1, \dots, U_n)$  such that  $\bigcup_i U_k^i = U_k$ .

## Definition (Multisort species)

A multi-sort species  $F[X_1, \dots, X_n]$  has the following data,

- For every finite multi-set, a finite set  $F[U_1, \dots, U_n]$ .

# Multiple variables

## Definition (Multiset)

- A multiset is a tuple  $(U_1, \dots, U_n)$  of sets.
- A function/bijection of multisets  $(f_k) : (U_1, \dots, U_n) \rightarrow (V_1, \dots, V_n)$  is a tuple of functions/bijections  $f_k : U_k \rightarrow V_k$ .
- A subset of  $(U_1, \dots, U_n)$  is a tuple  $(V_1, \dots, V_n)$  such that  $V_k \subseteq U_k$ .
- A partition of  $(U_1, \dots, U_n)$  is a collection  $\{U_1^i, \dots, U_n^i\}_i$  of subsets of  $(U_1, \dots, U_n)$  such that  $\bigcup_i U_k^i = U_k$ .

## Definition (Multisort species)

A multi-sort species  $F[X_1, \dots, X_n]$  has the following data,

- For every finite multi-set, a finite set  $F[U_1, \dots, U_n]$ .
- For every bijection  $f$ , a transport function  $F[U_1, \dots, U_n] \rightarrow F[V_1, \dots, V_n]$ .

## One final example

Consider the following PDE with  $F(X, Y)$  a 2-sort species.

$$\frac{\partial}{\partial X} F = Y(1 + \frac{\partial}{\partial Y})F$$

## One final example

Consider the following PDE with  $F(X, Y)$  a 2-sort species.

$$\frac{\partial}{\partial X} F = Y(1 + \frac{\partial}{\partial Y})F$$

We observe that this is satisfied by the species  $\text{Surj}(X, Y)$ .

## One final example

Consider the following PDE with  $F(X, Y)$  a 2-sort species.

$$\frac{\partial}{\partial X} F = Y(1 + \frac{\partial}{\partial Y})F$$

We observe that this is satisfied by the species  $\text{Surj}(X, Y)$ .  
From a description of  $\text{Surj}(X, Y)$ , we may guess a (non-unique) solution.

$$\text{Surj}(X, Y) = E((E(X) - 1)Y)$$

## One final example

Consider the following PDE with  $F(X, Y)$  a 2-sort species.

$$\frac{\partial}{\partial X} F = Y(1 + \frac{\partial}{\partial Y})F$$

We observe that this is satisfied by the species  $\text{Surj}(X, Y)$ .  
From a description of  $\text{Surj}(X, Y)$ , we may guess a (non-unique) solution.

$$\text{Surj}(X, Y) = E((E(X) - 1)Y)$$

We get a corresponding generating function  $e^{(e^x - 1)y}$  for the Stirling numbers.

# Uniqueness remarks

# Uniqueness remarks

Implicit Species Theorem (Bergeron, Labelle, Leroux)



# Uniqueness remarks

## Implicit Species Theorem (Bergeron, Labelle, Leroux)

Let  $H[X, Y]$  be a species satisfying  $H(0, 0) = 0$  and  $\frac{\partial H}{\partial Y}(0, 0) = 0$ . Then, there is a computable, unique up to canonical isomorphism, species  $A(X)$ , with  $A(0) = 0$ , satisfying

$$A(X) = H(X, A(X)).$$

# Uniqueness remarks

## Implicit Species Theorem (Bergeron, Labelle, Leroux)

Let  $H[X, Y]$  be a species satisfying  $H(0, 0) = 0$  and  $\frac{\partial H}{\partial Y}(0, 0) = 0$ . Then, there is a computable, unique up to canonical isomorphism, species  $A(X)$ , with  $A(0) = 0$ , satisfying

$$A(X) = H(X, A(X)).$$

This generalizes to  $k$ -systems of equations and  $(n, k)$  variables.

# Uniqueness remarks

## Implicit Species Theorem (Bergeron, Labelle, Leroux)

Let  $H[X, Y]$  be a species satisfying  $H(0, 0) = 0$  and  $\frac{\partial H}{\partial Y}(0, 0) = 0$ . Then, there is a computable, unique up to canonical isomorphism, species  $A(X)$ , with  $A(0) = 0$ , satisfying

$$A(X) = H(X, A(X)).$$

This generalizes to  $k$ -systems of equations and  $(n, k)$  variables.

## Proposition

The system of differential equations:

$$\frac{\partial Y_i}{\partial X} = R_i(Y_1, \dots, Y_k), \quad Y_i(0) = Z_i, \quad i = 1, \dots, k,$$

has a computable, unique up to isomorphism, solution... **for**  
 **$\mathbb{L}$ -species.**

Fin

