

Figure 40. Excess relative risk in assault death rates in year in which each month was $+1^{\circ}\text{C}$ compared with 1980-2009 norm temperatures by month, sex and age group.




Figure 41. Excess relative risk in suicide death rates in year in which each month was +1°C compared with 1980-2009 norm temperatures by month, sex and age group.




Figure 42. Excess relative risk in other unintentional injury death rates in year in which each month was +1°C compared with 1980-2009 norm temperatures by month, sex and age group.

6.5.4 Percentage change of injury deaths

Proportionally, deaths from drownings are predicted to increase more than those of other injury types, by as much 8.3% (7.3%, 9.3%) in men aged 15-24 years (Figure 43); the smallest proportional increase was that of assault and suicide (less than 2% in all age and sex groups). There was a larger percent increase in transport deaths for males than for females, especially in young and middle-ages (e.g., 1.25% (0.90%, 1.60%) for 25-34 year old men versus 0.23% (0.28%, 0.76%) for women of the same age) (Figure 43).

I also found some variation in association between proportional change in deaths for all ages combined and anomalous temperature across months for drownings, with excess risk from drownings exhibiting a distinct peak in summer months (Figure 44). In contrast, I found less variation in proportional change in deaths in the other four types of injuries, with transport, assault and suicide positively associated to a similar degree across all months, with falls consistently negatively associated apart from for females in late summer months (Figure 44).




Figure 43. Percent change in death rates in year in which each month was +1°C compared with 1980-2009 norm temperatures by type of injury, sex and age group summarised across months.




Figure 44. Percent change in death rates in year in which each month was +1°C compared with 1980-2009 norm temperatures by type of injury, sex and month summarised across ages.

6.5.5 Change in injury deaths

I estimated that there would be 941 (831, 1,053) excess injury deaths, equivalent to 0.5% of all injury deaths in 2016, in a year in which each month in each state were +1°C above its long-term norm (Figure 45). Deaths from drowning, transport, assault and suicide would be predicted to increase, partly offset by a decline in deaths from falls in middle and older ages and in winter months (Figure 45). Most excess deaths would be from transport injuries (448) followed by suicide (315). 87% of the excess deaths would occur in males and 13% in females. 80% of all male excess deaths would occur in those aged 15-64 years, who have higher rates of deaths from transport injuries. In those aged 85 years and older, there would be an estimated decline in injury deaths, because deaths from falls are expected to decline in a warmer year.




Figure 45. Additional annual injury deaths for the 2016 US population in year in which each month was +1°C warmer compared with 1980-2009 norm temperatures. The top row shows breakdown by type of injury, sex and age group. The bottom row shows the break down by type of injury, sex and month. Black dots represent net changes in deaths for each set of bars.

6.6 Discussion

While there are no previous studies of how temperature deviations from long-term norm in each month are associated with injury mortality, my results are broadly in agreement with those that have analysed associations with absolute temperature and for specific injury types. A study of suicide in US counties over 37 years (1968-2004) estimated that 1°C higher monthly temperature would lead to a 0.7% rise in suicides,¹⁰⁸ compared to my findings of 0.44-1% in males and 0.39-1.47% in females in different ages. In a study of six French heatwaves during 1971-2003, mortality from unintentional injuries rose by up to 4% during a heatwave period compared to a non-heatwave baseline.¹³³ A study of daily mortality from all injuries in Estonia found a 1.24% increase in mortality when daily maximum temperature went from the 75th to 99th percentile of long-term distribution.¹³²

That anomalously warm temperature influences deaths from drowning, although not previously quantified, is highly plausible because swimming is likely to be more common when monthly temperature is higher. The higher relative and absolute impacts on men compared with women may reflect differences in behaviour. For example, over half of swimming deaths for males occur in natural water, compared to about quarter for females (1999-2010),²⁵¹ which may lead to a larger rise in the former in warmer weather. Similarly, the decline in deaths from falls, which are mostly in older ages, may be because falls in older people are more likely to be due to slipping on ice than in younger ages.²⁵²⁻²⁵⁴

The pathways from anomalous temperature to transport injury are more varied. Firstly, driving performance deteriorates at higher temperatures.²⁵⁵⁻²⁵⁸ Further, alcohol consumption increases during warm temperature anomalies,²⁵⁹ potentially also explaining why teenagers, who are more likely than other age groups to crash while intoxicated,²⁶⁰ experience a larger proportional

rise in deaths from transport than older ages when temperatures are anomalously warm. Lastly, warmer temperatures generally increase road traffic in North America;^{261–264} With more people generally outdoors in warmer weather,²⁶⁵ this could lead to more fatal collisions.

Pathways linking anomalously high temperatures and deaths from assault and suicide are less established. One hypothesis is that, similar to transport, more time spent outdoors in anomalously warm temperatures leads to an increased number of face-to-face interactions, and hence arguments, confrontations, and ultimately assaults.^{266,267} These effects could be compounded by the greater anger levels linked to higher temperatures.^{268,269} Regarding suicide, higher temperature has been hypothesised as associated with higher levels of distress in younger people.²⁷⁰ Nonetheless, links between temperature and mental health requires further investigation,²⁷¹ including whether the relationship varies by sex and age group, as indicated by my results.

The major strength of my study is that I have comprehensively modelled the association of temperature anomaly with injury by type of injury, month, sex and age group. My measure of temperature anomaly internalises long-term historical experience of each state, and is closer to what climate change may bring about than solely examining daily episodes, or average temperature to which people have adapted. To utilise this metric, I integrated two large disparate national datasets on mortality (United States VR data) and meteorology (ERA-Interim), described in Chapter 3, and developed a bespoke Bayesian spatio-temporal model, described in Chapter 5. A limitation of my study is that, like all observation studies, I cannot rule out confounding of results due to other factors, although it is unlikely that such factors will have the same anomalies as temperature, even if their average space and time patterns are the same.

My work highlights how deaths from injuries are not only currently susceptible to temperature anomalies but could also be modified by rising temperatures resulting from climate change, unless countered by social and health system interventions that mitigate these impacts. Though absolute impacts on mortality are modest, some groups, especially men in young to middle-ages, will experience larger impacts. Therefore, a combination of public health interventions that broadly target injuries in these groups – for example targeted messaging for younger males on the risks of transport injury and drowning – and those that trigger in relation to forecasted high temperature periods – for example more targeted blood alcohol level checks – should be a public health priority.

6.7 Summary

Using the mortality and temperature datasets I developed in Chapter 3 and the Bayesian spatio-temporal model I described in Chapter 5, I estimated how anomalous temperatures affect injury deaths in the United States. I calculated that a 1°C anomalously warm year would result in a net increase in injury deaths. The large majority of these additional injury deaths would occur in males of adolescent to middle ages. Four of the five injury groups (transport, drownings, assault and suicide) contribute to the increase in injury deaths under anomalously warm temperatures, with a decline in deaths from falls in older ages.

7 Anomalous temperature and seasonality of cardiorespiratory disease and cancer mortality

7.1 Overview

I applied the Bayesian spatio-temporal model, as described in Chapter 5, to cardiorespiratory disease and cancer mortality data in the United States, as described in Chapter 3. I estimated how anomalous temperatures, defined in Chapter 3, affect deaths from different cardiovascular diseases (ischaemic heart disease (IHD), cerebrovascular disease, other cardiovascular diseases), respiratory diseases (chronic obstructive pulmonary disease (COPD), respiratory infections, other respiratory diseases) and cancers. A total of 61 million deaths from cardiorespiratory disease and cancer deaths in the United States from 1980-2016 were used in this analysis.

In this chapter, I present results of how a 1°C anomalously warm year would change the total number of cardiorespiratory disease and cancer deaths in the United States, by sex and age group. I then present additional results of how the percentage change of cardiorespiratory disease deaths would vary spatially for a 1°C anomalously warm year.

7.2 Introduction

The relationship between daily temperature and cardiorespiratory disease deaths is well-established.^{2-7,120} However, climate change is predicted to modify seasonal and monthly as well as daily patterns of temperature. This has the potential to disrupt long-term adaptation. This was one of the main motivations for building my Bayesian spatio-temporal model to associate monthly death rates with anomalous temperature, as detailed in Chapter 5. There is also limited analysis on how monthly deviations of temperature from long-term norms will

impact deaths in a consistent way across different types of cardiorespiratory disease and cancer deaths for demographic subgroups of the population.

My results from Chapter 6 predicted that there would be an increase in injury deaths, concentrated in younger males, with a 1°C anomalously warm year. In contrast, some previous studies predicted a decrease in cardiorespiratory disease deaths in a warmer climate.^{59,80,81,109} This would be the net effect of increases in cardiorespiratory disease deaths in the summer with larger decreases in the winter. Others argue that this may not be true due to a weakening association of excess winter mortality with temperature,^{272,273} though methods in these studies have been criticised as misinterpreting excess winter mortality.^{58,59} Cardiorespiratory diseases made up 40% of all deaths in the United States in 2016 and therefore remain significant to public health policy and planning. Deaths from cancers also make up a considerable proportion of deaths in the United States, with 23% of all deaths in 2016 being from cancers. Only a few studies have examined an association between temperature and deaths from cancers,^{127,135,138} though slight seasonality in cancer death rates exists for older age groups, shown in Chapter 4. My aim was to evaluate how deaths from cardiorespiratory diseases and cancers may be affected by changes in temperature that could arise as a result of global climate change in the United States.

7.3 Data

Full details of the data processing choices I made can be found in Chapter 3. Here, I used VR data on all 60,967,769 cardiorespiratory disease and cancer deaths in the contiguous United States from 1980 to 2016 along with population records to calculate monthly death rates for each type of cardiorespiratory disease and cancer death, state, sex and age group combination. Of cardiorespiratory disease and cancer deaths, 41,105,009 (67.4%) were from cardiorespiratory diseases and 19,862,760 (35.6%) were from cancers. For each death, I

mapped each ICD-9 and ICD-10 code to cardiorespiratory diseases and cancers (Table 11). I separated the cardiorespiratory disease deaths into three cardiovascular (IHD, cerebrovascular disease, other cardiovascular diseases) and three respiratory (COPD, respiratory infections, other respiratory diseases) diseases. Other cardiovascular diseases included rheumatic, hypertensive and inflammatory heart diseases. Other respiratory diseases included asthma and bronchiectasis. I did not further subdivide cancer deaths as death outcomes are not generally considered as sensitive to temperature changes.^{127,135,138} The remainder of deaths not included in analysis in Chapters 6 and 7 were from a heterogeneous group of causes other than cancers, cardiorespiratory diseases and injuries, described in more detail in Chapter 3. The causes of death that led to deaths in this group varied by sex and age group. Therefore, I did not include them in the analysis. I obtained monthly population-weighted temperature anomalies from ERA-Interim by a process described in Chapter 3.²¹⁵

7.4 Statistical methods

I analysed the association of monthly death rates with anomalous temperature using a Bayesian spatio-temporal model, fully described in Chapter 5 and described in brief in Section 6.4.

Chronic disease	ICD-9	ICD-10
Cardiorespiratory diseases	381-382, 390-519	H65-H66, I00-J99
Cardiovascular diseases	390-459	I00-I99
Ischaemic heart disease (IHD)	410-414	I20-I25
Cerebrovascular disease	430-438	I60-I69
Other cardiovascular diseases	390-409, 415-429, 439-459	I00-I19, I26-I59, I70-I99
Respiratory diseases	381-382, 460-519	H65-H66, J00-J99
Chronic obstructive pulmonary disease (COPD)	490-492, 495-496	J40-J44
Respiratory infections	381-382, 460-466, 480-487	H65-H66, J00-J6, J9-J18, J20-J22
Other respiratory diseases	467-479, 488-489, 493-494, 497-519	J7-J8, J19, J23-J39, J45-J99
Cancers	140-239	C00-C99, D00-D48

Table 11. Cardiorespiratory disease and cancer groups used in the analysis with ICD-9 and ICD-10 codes.

7.5 Results

7.5.1 Summary of cardiorespiratory disease and cancer deaths

From 1980 to 2016, 20,070,797 boys and men and 21,034,212 girls and women died from cardiorespiratory diseases in the contiguous United States. These deaths accounted for 46.3% and 49.9% of all male and female deaths respectively (Figure 2 in Chapter 3). 77.1% of male cardiorespiratory disease deaths and 88.7% of female cardiorespiratory disease deaths were in those aged 65 years and older (Figure 46). For females, 42.7% of cardiorespiratory disease deaths were in those aged 85 years and older (Figure 46).

IHD accounted for 45.3% of cardiorespiratory disease deaths in males and 38.7% in females (Figure 46). Men 35 years and older and women 45 years and older died from IHD more than any other cardiorespiratory disease. Below 35 years of age for men and 45 years of age for women, other cardiovascular diseases were the largest single cause of cardiorespiratory disease death in both males and females (Figure 46). As I presented in Chapter 4, there has been a persistent seasonality of mortality in cardiorespiratory disease deaths over time for most age groups in both sexes (Figure 17 and Figure 18). For each of the six types of cardiorespiratory disease death, there were more deaths in the winter than in the summer (Figure 47).

There were 10,428,202 cancer deaths for boys and men and 9,434,558 for girls and women in the contiguous United States during 1980-2016 (Figure 2 in Chapter 3). There were slightly more deaths from cancers in winter than in summer (Figure 47) as we found for those 55 years and older in Chapter 4 (Figure 15 and Figure 16).




Figure 46. Number of deaths by cardiorespiratory disease and cancer, sex and age group in the contiguous United States for 1980-2016.




Figure 47. Number of deaths by cardiorespiratory disease and cancer, month, sex and age group in the contiguous United States for 1980-2016.

7.5.2 Trends in cardiorespiratory disease and cancer mortality

There was a decline in age-standardised death rates of four out of six types of cardiorespiratory disease (IHD, cerebrovascular disease, other cardiovascular diseases and respiratory infections) for both men and women from 1980 to 2016. IHD death rate declined more than other cardiorespiratory diseases from 1980 to 2016, by over 50% for both men and women (Figure 48). COPD death rates increased in males from the 1980s until the late 1990s, and steadily decreased since then. In contrast, COPD death rates in females have increased since the 1980s (Figure 48). Age-standardised death rates from other respiratory diseases also increased over the time period (Figure 48). There is a discontinuity in age-standardised death rates in other cardiovascular diseases due to the change from the ICD-9 to ICD-10 coding in the United States in 1999, when some deaths in this group were assigned to IHD (Figure 48). Death rates from cancers have decreased over time (Figure 48).




Figure 48. National age-standardised death rates from 1980 to 2016, by cardiorespiratory disease and cancer, month and sex.

7.5.3 Excess risk of cardiorespiratory disease and cancer deaths

I used the resultant risk estimates (Figure 49-Figure 55) to calculate additional deaths if each month in each state were +1°C above its long-term norm. I found no consistent association between deaths for cancers and anomalous temperature (Figure 55). I therefore did not include the results in further calculations.




Figure 49. Excess relative risk in IHD death rates in year in which each month was +1°C compared with 1980-2009 norm temperatures by month, sex and age group.




Figure 50. Excess relative risk in cerebrovascular disease death rates in year in which each month was +1°C compared with 1980-2009 norm temperatures by month, sex and age group. 163




Figure 51. Excess relative risk in other cardiovascular disease death rates in year in which each month was $+1^{\circ}\text{C}$ compared with 1980-2009 norm temperatures by month, sex and age group. 164




Figure 52. Excess relative risk in COPD death rates in year in which each month was +1°C compared with 1980-2009 norm temperatures by month, sex and age group. 165




Figure 53. Excess relative risk in respiratory infection death rates in year in which each month was +1°C compared with 1980-2009 norm temperatures by month, sex and age group. 166




Figure 54. Excess relative risk in other respiratory disease death rates in year in which each month was +1°C compared with 1980-2009 norm temperatures by month, sex and age group. 167




Figure 55. Excess relative risk in cancer death rates in year in which each month was $+1^{\circ}\text{C}$ compared with 1980-2009 norm temperatures by month, sex and age group. 168

7.5.4 Percentage change of cardiorespiratory disease deaths

Proportionally, deaths from IHD are predicted to show the most consistent decrease in older age groups (Figure 56). Proportional reductions in deaths for those 55 years and older were greater than 0.5% in males and 0.2% in females in a year in which each month in each state were +1°C above its long-term norm (Figure 56). Younger age groups generally possess a large range of uncertainty in all types of cardiorespiratory disease death due to the small numbers of deaths (Figure 56). The largest single age group decrease was that of respiratory infections in males 85 and older, (1.0% (0.7%, 1.3%)) (Figure 56). With few exceptions (e.g., those aged 15-24 years in deaths from other respiratory diseases) the scale and sign of the percent changes were similar in both males and females across ages and cardiorespiratory diseases (Figure 56).

There were variations in association between proportional change in deaths for all ages combined across months (Figure 57). The main exceptions were for COPD in females and other respiratory diseases in both sexes (Figure 57), though low absolute numbers of other respiratory disease deaths led to the large uncertainty in the estimated rate ratios (Figure 57). The largest single monthly increase was for other cardiorespiratory diseases for females in July, 0.7% (0.4%,1.0%), with the largest single decrease for IHD for males in December, 1.1% (0.9%, 1.2%) (Figure 57).




Figure 56. Percent change in death rates in year in which each month was +1°C compared with 1980-2009 norm temperatures by cardiorespiratory disease, sex and age group summarised across months.




Figure 57. Percent change in death rates in year in which each month was +1°C compared with 1980-2009 norm temperatures by cardiorespiratory disease, sex and month summarised across age.

7.5.5 Change in cardiorespiratory disease deaths

I estimated that there would be 4,369 (4,024, 4,706) fewer cardiorespiratory disease deaths, equivalent to 0.4% of all cardiorespiratory disease deaths in 2016, in a year in which each month in each state were +1°C above its long-term norm (Figure 58). Deaths from all cardiovascular diseases (IHD, cerebrovascular disease and other cardiovascular diseases) and respiratory diseases (COPD, respiratory infections and other respiratory diseases) would be predicted to decrease (Figure 58). The biggest reduction of deaths would be from IHD, with 1,883 (1,706, 2,058) fewer deaths, followed by other cardiovascular diseases, with 1,039 (835, 1,243) fewer deaths. 81% of the total reduction in deaths would be from cardiovascular diseases, with the rest in respiratory diseases (Figure 58). 4,236 deaths (97%) of overall reduction would come from those aged 55 years and older. 55% of the reduction in deaths would occur in males and 45% in females (Figure 58). There would be decreases in all months across all cardiorespiratory diseases, with the exception of the summer months (June, July and August). The greatest single increase would be in July for other cardiovascular diseases for both males (76 (41, 112)) and females (103 (62, 144)). The biggest reductions would occur in the coldest months (November, December, January and February) (Figure 58).




Figure 58. Additional annual cardiorespiratory disease deaths for the 2016 US population in year in which each month was +1°C compared with 1980-2009 norm temperatures. The top row shows breakdown by cardiorespiratory disease, sex and age group. The bottom row shows the break down by cardiorespiratory disease, sex and month. Black dots represent net changes in deaths for each set of bars.

7.5.6 Percentage change of sub-national cardiorespiratory disease deaths

I also ran a model where each state was given a separate temperature anomaly coefficient to examine spatial variation in percentage change in deaths if each month in each state were +1°C above its long-term norm, described by Equation 9 in Chapter 5. In this model, I used all cardiorespiratory diseases together, as models did not converge or give sensible results when breaking down deaths by type of cardiorespiratory disease in addition to state, sex and age group.

Percentage change in death rates in a year in which each month in each state were +1°C above its long-term norm varied by state and month in both males and females (Figure 59 and Figure 60), though in some months more than others. There were similar associations between males and females in the same state and month. In January, an illustrative month for the impacts of anomalous temperature in winter months, there were decreases in percentage change in death rates across the entire contiguous United States for both males and females (Figure 59 and Figure 60). These decreases ranged from 1.5% (1.2%, 1.9%) (Louisiana) to 0.2% (0.1%, 0.4%) (New Mexico) in males (Figure 59), and 1.3% (1.0%, 1.7%) (Washington) to 0.1% (0.3%, 0.5%) (Arizona) in females (Figure 60). In July, an illustrative month for the impact of anomalous temperature in summer months, there were increases in 47 out of the 49 spatial units for males (Figure 59) and all spatial units for females (Figure 60). The biggest increase in July for males was in New Mexico, at 1.1% (0.5%, 1.7%) (Figure 59) and in Utah, at 1.2% (0.6%, 1.7%) for females (Figure 60). In May and September, late spring and early autumn respectively, there were mixed positive and negative associations (Figure 59 and Figure 60). In these months, there were negative associations with anomalous temperature in coastal regions. In contrast, there were positive associations in the West North Central and Southwest regions of the United States, though it should be noted that the low population of states in this

region contributed to large uncertainties in these estimates. There were similar associations in June and August across the United States in both males and females (Figure 59 and Figure 60). Overall, the Southeast region showed the greatest negative associations with anomalous temperature across all months for both sexes.




Figure 59. Percent change in cardiorespiratory disease death rates for males in year in which each month was +1°C compared with 1980-2009 norm temperatures by state and month summarised across age groups.




Figure 60. Percent change in cardiorespiratory disease death rates for females in year in which each month was +1°C compared with 1980-2009 norm temperatures by state and month summarised across age groups.

7.6 Discussion

My work here shows that there would be an expected net decrease from cardiorespiratory disease deaths in a year in which each month in each state were $+1^{\circ}\text{C}$ above its long-term norm (Figure 58). This reduction would be strongly concentrated in older ages, be shared almost evenly between males and females and occur in all but summer months (Figure 58).

Most previous studies have examined how deaths from all-causes^{2,3,101,103,109,111–117,5,118,6,7,87,88,98–100} and cardiorespiratory disease deaths^{3,5,121–127,6,7,87,88,115–117,120} are associated with daily, not monthly, temperature. Nevertheless, some limited comparisons can be made with other studies.^{103,117,122} It should be noted that for summer months one study examined effect of extreme high temperatures¹¹⁷ and the other the effect of a 5°C increase in temperatures.¹²² A study of winter months looked at decreases below 18°C .¹⁰³

Illustrative of summer months, there was an increase of IHD deaths in July of 0.4% (0.2%, 0.6%) for males and 0.5% (0.3%, 0.8%) in females for a 1°C anomalously warm year (Figure 57). Previous studies which studied association of IHD deaths in summer months with daily temperature found slightly higher (1.7% to 2.5%) increases of risk for all deaths from both sexes.^{117,122} My cerebrovascular disease findings in July show a 0.1% (-0.3%, 0.4%) increase in males and 0.5% (0.2%, 0.8%) in females (Figure 57). Another study of daily temperature found higher risk (2.5%) than in my findings. For COPD, I found a negative association of percentage change across most months for both sexes (Figure 57). A previous study found a large daily association for summer COPD deaths (4.3%), though this was for extreme high temperatures.¹¹⁷ This is in contrast to my study, which examined the association of mild deviations from long-term norm temperatures on mortality.

Representing winter months, my IHD findings in January show a 0.9% (0.8%, 1.1%) decrease in males and 0.6% (0.4%, 0.7%) in females for a 1°C anomalously warm year (Figure 57). This result is consistent with a previous study of cold months in Europe.¹⁰³

There are plausible explanations for my results. Cardiovascular deaths are higher in the winter than the summer (Figure 47). This is partially explained by blood pressure being inversely associated with temperature.¹²⁹ Cold temperature in the winter therefore results in generally higher blood pressure, placing strain on the heart and damaging vessel walls.¹²⁹ The lower norm temperatures in winter months can also lead to increased blood clotting and thrombosis.^{103,129} This increases the risk of IHD and cerebrovascular disease and ultimately death from these causes.¹⁰³ Warm temperatures in winter months would therefore reduce the risk of dying from these causes (Figure 58). That warmer temperatures in summer months increase risk of death in cardiovascular diseases is thought to be due to the reduction of plasma volume by release of platelets and red blood cells into the blood stream and the loss of water and salt through sweating.¹³¹ This puts strain on the circulatory system by increasing sticking of platelets to artery walls which leads to blockages and sudden decrease in blood pressure and can lead to death.¹³¹ Further increases away from long-term norms increase strain on the circulatory system and risk of death from cardiovascular diseases. This explains why more deaths are expected from cardiovascular diseases in anomalously warm summer months (Figure 58).

Respiratory disease deaths are also higher in winter than at other times in the year in the United States (Figure 47). Colder temperatures are associated with suppression of the immune system, which facilitates respiratory infections taking hold in humans.¹⁰³ Warmer anomalies in winter months can therefore reduce risks of respiratory infections, which explains the reduction in excess risk for older groups (Figure 53). Cold air can also cause airways to go into spasm and trigger coughing, wheezing, shortness of breath and tightness in the chest.²⁷⁴ Mould spores are more common in colder air, known to cause coughing and sneezing, which exacerbate existing

respiratory conditions.²⁷⁵ Warmer anomalies in winter would reduce these risk factors. In contrast, warmer anomalous temperatures in summer months are associated with airway inflammation, which can lead to death from respiratory complications.¹³⁰

I found no consistent association between deaths from cancers and a year in which each month in each state were +1°C above its long-term norm (Figure 55). This agrees with limited previous studies,^{127,135,138} with the exception of an old study which advised caution in its interpretation as it did not account for geographical differences.¹³⁵ Though I found that cancer deaths were weakly seasonal in males 55 years and in females 65 years and older (Figure 15 and Figure 16 in Chapter 1), this may be for reasons other than temperature variation. I note this though I found a non-significant increase for respiratory infections in summer months and in no other respiratory disease (Figure 57).

The major strength of my anomalous temperature study is that, when combined with work in Chapter 6, I comprehensively modelled the association of temperature anomaly with a comprehensive set of causes of death, by month, sex and age group. I did this using the two large datasets I processed in Chapter 3 and the Bayesian spatio-temporal model I developed in Chapter 5. The temperature anomaly metric I also described in Chapter 3 (Figure 8) factors in long-term local experience when examining temperature-mortality association. This has enabled me to make an estimate of the change in deaths with a 1°C anomaly increase, which is within the range of anomaly size experienced by many states (Figure 9 in Chapter 3). My study design has also enabled me to show that the reduction in deaths would be greatest in older groups in the winter months, and would be almost evenly shared between males and females (Figure 58). A limitation of my study is that I cannot rule out confounding of results due to other factors that may be correlated with temperature anomalies.

My work adds to the body of evidence that suggests that a warming world will reduce winter deaths.^{59,80,81,109} While increases in average temperatures would increase deaths in the summer, this would be outweighed by the reductions in winter (Figure 58). My work goes further than previous work by breaking down changes in deaths by cause of death, month, age group and sex (Figure 58). The predicted reduction of 0.4% of all cardiorespiratory disease deaths is slight and concentrated in older age groups. This demonstrates that the burden from cardiorespiratory disease deaths will remain large and a significant public health concern under climate change.

7.7 Summary

Using the mortality and temperature datasets I developed in Chapter 3 and the Bayesian spatio-temporal model I described in Chapter 5, I estimated how anomalous monthly temperatures affect cardiorespiratory disease and cancer deaths in the United States. I calculated that a 1°C anomalously warm year would result in a net decrease in cardiorespiratory disease deaths. The large majority of this reduction would be concentrated in older age groups. The reduction would be across all cardiorespiratory diseases, with the largest in ischaemic heart disease. Reductions would be shared almost evenly between males and females. Reductions would occur across the year, apart from the summer months, when deaths would increase slightly. I found no association between anomalous temperature and cancer deaths.

8 Discussion

8.1 Comparison with published literature

8.1.1 Seasonal dynamics of mortality

Prior to my work, it had been established that total mortality from the entire population in the United States are higher in the winter than in the summer.^{57,60,70} This was known to be mainly due to higher winter death rates from influenza and cardiorespiratory diseases in older age groups.^{71–74} Previous studies in the United States have also identified seasonality for some causes -of death.^{54,57,60–62,70,224} No previous study made a consistent national and subnational analysis of seasonality in the United States by cause of death, sex, age group, over time and space, nor how it varies across climate regions. In Chapter 4 of this thesis, I used wavelet and centre of gravity analyses, which allowed me to identify seasonality of mortality and how it has changed over time. This work uncovered the distinct behaviours across causes of death, and by sex and age group.

Where studies were comparable, my results agreed with previous work. I demonstrated that the strong seasonality in older age groups of cardiorespiratory diseases and weak seasonality of cancer deaths, found in some previous studies,^{60,61,70,224} have endured over the last four decades. I also extended previous work. For example, I showed that there have been variations in when injury deaths peaked and how the seasonal differences in injury deaths have changed over time in relation to age group and sex, not previously demonstrated. I also showed how changes in all-cause mortality seasonality by age group and sex can be explained by changes in particular causes of death, such as how substantial decline in seasonal mortality differences in boys under five was due to the decreasing seasonality of death from cardiorespiratory diseases in that age group.

8.1.2 Anomalous temperature and mortality

A large proportion of previous studies of the temperature-mortality relationship focused on single day or multi-day episodes and temperature.^{3–7} Other studies either associated monthly or seasonal average temperatures with mortality,^{108,109} or accounted for seasonal impacts by stratifying daily data by month.^{81,114} Previous studies largely examined all-cause,^{2,3,101,103,109,111–117,5–7,87,88,98–100} or cardiorespiratory disease deaths.^{3,5,121–126,6,7,87,88,115–117,120} In the United States, studies ranged from a single spatial unit,^{86,87,109,122,137,168,169} to hundreds of independently-analysed units across the country.^{81,101,155,161,170,114,118,121,124,127,136,144,146} Some work used Bayesian methods to conduct meta-analyses of relative risks for temperature and mortality.^{172,173}

To my knowledge, no previous study made an analysis of how temperature deviations from long-term norm in each month are associated with mortality by cause of death, sex and age group. In my work on anomalous temperature and mortality in Chapters 6 and 7, I examined deviations from long-term norms to analyse relationships using a Bayesian spatio-temporal model by cause of death, month, sex, age group and in the case of cardiorespiratory disease deaths (Chapter 7), by state.

Though previous studies did not examine anomalous temperature, some of my results can be compared with other studies of temperature. For example, my results for injury deaths, described in Chapter 6, align well with a previous study of the association between suicides and temperature in the entire contiguous United States and Mexico.¹⁰⁸ However, my work went further in demonstrating that while suicides are predicted to increase with rising anomalous temperature, the greatest absolute impact would be on younger men. With cardiorespiratory disease deaths, described in Chapter 7, I showed that there would be a net decrease expected with mild positive deviations from long-term norm temperatures. This would be concentrated

in the oldest age groups and would be shared almost evenly between males and females. My results agreed with previous studies which predicted a decrease in absolute number of deaths for cardiorespiratory disease deaths.^{59,80,81,109} My study went further by showing that this would be concentrated in older age groups, and percentage change in cardiorespiratory disease deaths would be nearly homogeneous across space in the warmest and coldest months (Chapter 7). I also showed in Chapter 7 that there was no association with cancer and anomalous temperature across any month, sex and age group combination.

8.2 Strengths and limitations

In my thesis, I processed two large datasets on death rates from VR records and monthly anomalous temperature from ERA-Interim weather reanalysis, described in Chapter 3. My work here is the first comprehensive study of the United States to use a mutually exclusive and collectively exhaustive set of causes of death for both examining dynamics of seasonality of mortality, described in Chapter 4, as well as association between anomalous temperature and mortality, described in Chapters 6 and 7.

The strengths of my dynamics of seasonality study, described in Chapter 4, were its innovative methods of characterizing seasonality of mortality dynamically over space and time, by cause of death, sex and age group and; using wavelet and centre of gravity analyses; and using ERA-Interim data to compare the association between seasonality of death rates and regional temperature.

I developed a Bayesian spatio-temporal model, described in Chapter 5, to borrow strength across spatial and temporal units. Using this model, I estimated excess risk of mortality for injuries, described in Chapter 6, and cardiorespiratory disease and cancer deaths, described in Chapter 7. I was able to go further than previous studies by examining changes in deaths due to anomalous temperature by cause of death, sex and age group.

A limitation of my results in Chapters 4 ,6 and 7 is that I did not investigate seasonality of mortality by socioeconomic characteristics which may help with understanding its determinants and planning responses. Whether temperature anomaly within a month is spread or concentrated, as described in Chapters 6 and 7, may also have different implications to health as single-day or multi-day heat and cold events are known to be a threat to human lives. However, additional extreme temperature measures, which I described in Section 5.4.11, did not add any further insight to my analysis. Like all observation studies, confounding from other factors cannot be ruled out. However, it is unlikely that such factors will have the same anomalies as temperature, even if their average space and time patterns are the same. Future climate change may also produce different types and patterns of anomalies from the ones we can study historically.

8.3 Public health and policy implications

Continued higher death counts in winter for cardiorespiratory diseases in older age groups, shown in Chapter 4, indicate that improving interventions during the winter months will remain a priority. These may include better insulation of homes, winter heat provision and influenza vaccinations. Social interventions will remain important in improving and maintaining health in older age, such as regular visits to older people or keeping them connected over the internet to other people with newer technologies. This effort is necessary irrespective of the geography and local climate in the United States, since I also showed in Chapter 4 that seasonality is present and similarly timed throughout the country. The relative difference in seasonality of mortality has decreased for children under five, which may help to demonstrate the effectiveness of efforts to reduce peaks of cardiorespiratory diseases in the winter. The continued presence of cancer death seasonality in older groups may be due to co-morbidity in winter months, causing already frail sufferers to die.

There would be increases in deaths from injuries (transport, drownings, assault, suicides) in younger males, during periods of elevated anomalous temperature, as shown in Chapter 6. My analysis demonstrated that a 1°C anomalously warm year would result in 0.5% overall increase in injury deaths, concentrated in younger men. This may have implications on interventions targeted at younger age groups during periods of elevated temperature. These could include targeted messaging for younger males on the dangers of driving and swimming triggered when higher-than-normal temperatures are forecast. The concentration of these deaths in younger age groups, particularly younger men, also means that additional injury deaths from anomalous temperatures could result in loss to the economy through loss of otherwise healthy working years. The estimated reduction in cardiorespiratory disease deaths (0.4% overall decrease in a 1°C anomalously warm year), concentrated in older age groups, demonstrate that cardiorespiratory disease deaths would remain an ongoing concern under climate change, since climate change would not reduce cardiorespiratory disease deaths by a significant amount. Policy makers should continue to aim for further reductions in such chronic diseases which, along with cancers, remain the deadliest threat to most people in the United States.

8.4 Future work

My work in this thesis is translatable to other countries with reliable vital registration data. Though many countries are still lacking appropriate high quality mortality data,¹⁴ the analyses in Chapters 4, 6 and 7 could be replicated as data becomes available. The algorithm I developed for gridded weather data, described in Chapter 3, could be used anywhere in the world. Comparing differences between communities in different countries could serve as evidence to measure success of potential adaption measures enacted in one country but not another.

In the United States itself, other research on anomalous temperature and mortality could examine urban and rural differences. Similar work has been carried out in the United

Kingdom.^{120,143} Further work could examine how socioeconomic factors explain differences by state in excess risk per 1°C temperature anomaly by state, such as in my analysis of cardiorespiratory disease deaths in Chapter 7. By combining with climate model projections of future deviations of temperature from current long-term norms, my work here could be used to help further understand the implications of climate change on mortality by cause of death, sex and age group. A recent study on all-cause mortality for several cities in the United States highlighted the differences in attributable heat mortality between meeting the Paris Agreement's temperature goals and current trajectory of about 3°C increase above preindustrial levels.¹¹⁹ Further work based on my thesis could expand this tranche of work on climate change and human mortality to multiple causes of death, by sex and age group and across the entire contiguous United States, using the benefits of Bayesian spatio-temporal modelling, such as the 'borrowing of strength' across different spatial and temporal subunits.

8.5 Conclusions

Seasonality of mortality persists for deaths in older age groups in the United States, though the timing and degree to which this occurs varies by cause of death, sex and age group. Seasonal differences in deaths among children and young adults have largely disappeared. There are impacts, though slight, in how injury and cardiorespiratory disease death rates would change under anomalous temperature. Injury deaths in young men, particularly for transport, drowning, assault and suicide, are positively associated with anomalous warm temperature. In contrast, deaths from falls in older age groups were negatively associated with anomalous warm temperature. Anomalous warm temperature is also associated with decreases in cardiorespiratory disease deaths. This decrease would be concentrated in older age groups and shared almost equally between males and females. There is no association between cancer deaths and anomalous temperatures. My research shows that in the United States, continued efforts are required to further reduce peaks in mortality in older age groups through targeted

interventions. Under climate change, while estimated modest increases in injury mortality in younger males is concerning, the burden of cardiorespiratory disease and cancer deaths will remain a high public health priority due to the respective small and no reduction in deaths.

Works cited

- 1 CDC. Estimates of influenza vaccination coverage among adults—United States, 2017–18 flu season. FluVaxView. 2018.
- 2 Gasparrini A, Guo Y, Hashizume M, *et al.* Mortality risk attributable to high and low ambient temperature: a multicountry observational study. *Lancet* 2015; **386**: 369–75.
- 3 Basu R. High ambient temperature and mortality: a review of epidemiologic studies from 2001 to 2008. *Environ Heal A Glob Access Sci Source* 2009; **8**: 40.
- 4 Ye X, Wolff R, Yu W, Vaneckova P, Pan X, Tong S. Ambient temperature and morbidity: a review of epidemiological evidence. *Environ Health Perspect* 2012; **120**: 19–28.
- 5 Basu R, Samet JM. Relation between elevated ambient temperature and mortality: a review of the epidemiologic evidence. *Epidemiol Rev* 2002; **24**: 190–202.
- 6 Kovats RS, Hajat S. Heat stress and public health: a critical review. *Annu Rev Public Health* 2008. DOI:10.1146/annurev.publhealth.29.020907.090843.
- 7 Song X, Wang S, Hu Y, *et al.* Impact of ambient temperature on morbidity and mortality: an overview of reviews. *Sci Total Environ* 2017; **586**: 241–54.
- 8 WHO. COP24 special report: health and climate change. 2018.
- 9 UNFCCC. Paris Agreement. 2015.
- 10 Watts N, Amann M, Arnell N, *et al.* The 2018 report of the Lancet Countdown on health and climate change: shaping health of nations for centuries to come. *Lancet* 2018; **6736**: 1–4.
- 11 WHO. World health statistics 2016: monitoring health for the SDGs. 2016.
- 12 Great Britain. Weekly return of births and deaths in London and in other great towns. *Gen Regist Off* 1876.
- 13 National Research Council (US) Committee on National Statistics. Vital statistics: summary of a workshop. 2009 DOI:10.17226/12714.
- 14 WHO. WHO methods and data sources for country-level cause of death 2000-2016. 2018 DOI:ISBN 9789241564854.
- 15 Omran AR. A century of epidemiologic transition in the United States. *Prev Med (Baltim)* 1977. DOI:10.1016/0091-7435(77)90003-2.
- 16 Ma J, Ward EM, Siegel RL, Jemal A. Temporal trends in mortality in the United States, 1969-2013. *JAMA - J Am Med Assoc* 2015. DOI:10.1001/jama.2015.12319.
- 17 Murphy SL, Xu J, Kochanek KD, Arias E. Mortality in the United States, 2017. *NCHS Data Brief* 2018.
- 18 WHO. WHO methods and data sources for life tables 1990-2016. 2018. http://www.who.int/healthinfo/statistics/LT_method.pdf.
- 19 Oeppen J, Vaupel JW. Broken limits to life expectancy. *Science (80-)* 2002. DOI:10.4054/DemRes.2011.24.11.
- 20 Riley JC. Rising life expectancy: a global history. 2015 DOI:10.1017/CBO9781316036495.
- 21 Ezzati M, Friedman AB, Kulkarni SC, Murray CJL. The reversal of fortunes: trends in county mortality and cross-county mortality disparities in the United States. *PLoS Med*

2008. DOI:10.1371/journal.pmed.0050066.
- 22 Chetty R, Stepner M, Abraham S, *et al.* The association between income and life expectancy in the United States, 2001-2014. *JAMA - J Am Med Assoc* 2016. DOI:10.1001/jama.2016.4226.
 - 23 Murray CJL, Kulkarni SC, Michaud C, *et al.* Eight Americas: investigating mortality disparities across races, counties, and race-counties in the United States. *PLoS Med* 2006. DOI:10.1371/journal.pmed.0030260.
 - 24 NCHS. Health, United States, 2017: with special feature on mortality. 2017.
 - 25 Kontis V, Bennett JE, Mathers CD, Li G, Foreman K, Ezzati M. Future life expectancy in 35 industrialised countries: projections with a Bayesian model ensemble. *Lancet* 2017. DOI:10.1016/S0140-6736(16)32381-9.
 - 26 Foreman KJ, Marquez N, Dolgert A, *et al.* Forecasting life expectancy, years of life lost, and all-cause and cause-specific mortality for 250 causes of death: reference and alternative scenarios for 2016–40 for 195 countries and territories. *Lancet* 2018. DOI:10.1016/S0140-6736(18)31694-5.
 - 27 Olshansky SJ, Passaro DJ, Hershow RC, *et al.* A potential decline in life expectancy in the United States in the 21st Century. *N Engl J Med* 2005; **60**: 450–2.
 - 28 Costello A, Abbas M, Allen A, *et al.* Managing the health effects of climate change. *Lancet* 2009; **373**: 1693–733.
 - 29 Pope CA, Ezzati M, Dockery DW. Fine-particulate air pollution and life expectancy in the United States. *N Engl J Med* 2009. DOI:10.1056/nejmsa0805646.
 - 30 Wallace JM, Hobbs P V. Atmospheric science: An introductory survey. 2006 DOI:10.1016/C2009-0-00034-8.
 - 31 India Meteorological Department. History of meteorological services in India. Gov. India. 2018. http://www.imd.gov.in/pages/about_history.php (accessed April 2, 2019).
 - 32 Middleton WEK. A history of the thermometer. Johns Hopkins, Baltimore, 1966.
 - 33 Middleton WEK. A brief history of the barometer. *J R Astron Soc Canada* 1944; **28**.
 - 34 Manley G. Central England temperatures: monthly means 1659 to 1973. *Q J R Meteorol Soc* 1974. DOI:10.1002/qj.49710042511.
 - 35 Vose RS, Applequist S, Squires M, *et al.* Improved historical temperature and precipitation time series for U.S. climate divisions. *J Appl Meteorol Climatol* 2014. DOI:10.1175/JAMC-D-13-0248.1.
 - 36 NOAA. NOAA weather stations. [arcgis.com](https://www.arcgis.com/home/webmap/viewer.html?webmap=fa40207019de41bc9a37eal5235ae0c). 2014. <https://www.arcgis.com/home/webmap/viewer.html?webmap=fa40207019de41bc9a37eal5235ae0c> (accessed Jan 8, 2019).
 - 37 Karl TR, Koss WJ. Regional and national monthly, seasonal, and annual temperature weighted by area, 1895-1983. *Natl Clim Data Cent* 1984; : 38.
 - 38 Parks RM, Bennett JE, Foreman KJ, Toumi R, Ezzati M. National and regional seasonal dynamics of all-cause and cause-specific mortality in the USA from 1980 to 2016. *Elife* 2018; **7**. DOI:10.7554/eLife.35500.
 - 39 Fall S, Watts A, Nielsen-Gammon J, *et al.* Analysis of the impacts of station exposure on the US historical climatology network temperatures and temperature trends. *J Geophys Res Atmos* 2011. DOI:10.1029/2010JD015146.
 - 40 Karl TR, Melillo JT, Peterson TC. Global climate change impacts in the United States. Cambridge University Press, 2009.

- 41 Menne MJ, Williams CN, Palecki MA. On the reliability of the U.S. surface temperature record. *J Geophys Res Atmos* 2010. DOI:10.1029/2009JD013094.
- 42 Menne MJ, Williams CN, Vose RS. The U.S. historical climatology network monthly temperature data, version 2. *Bull Am Meteorol Soc* 2009. DOI:10.1175/2008BAMS2613.1.
- 43 Menne MJ, Williams CN. Homogenization of temperature series via pairwise comparisons. *J Clim* 2009. DOI:10.1175/2008JCLI2263.1.
- 44 Sun L, Kunkel KE, Stevens LE, Buddenberg A, Dobson JG, Easterling DR. Regional surface climate conditions in CMIP3 and CMIP5 for the United States: differences, similarities, and implications for the US national climate assessment. *NOAA Tech Rep NESDIS 144* 2015; : 111-p.
- 45 Xu Z, FitzGerald G, Guo Y, Jalaludin B, Tong S. Impact of heatwave on mortality under different heatwave definitions: a systematic review and meta-analysis. *Environ Int* 2016. DOI:10.1016/j.envint.2016.02.007.
- 46 Habeeb D, Vargo J, Stone B. Rising heat wave trends in large US cities. *Nat Hazards* 2015. DOI:10.1007/s11069-014-1563-z.
- 47 US global change research program. Heat waves. 2018. <https://www.globalchange.gov/browse/indicators/us-heat-waves> (accessed May 20, 2019).
- 48 Jones B, O'Neill BC, McDaniel L, McGinnis S, Mearns LO, Tebaldi C. Future population exposure to US heat extremes. *Nat Clim Chang* 2015; **5**: 652–5.
- 49 Mora C, Dousset B, Caldwell IR, *et al.* Global risk of deadly heat. *Nat Clim Chang* 2017; **7**: 501–6.
- 50 Rogot E, Fabsitz R, Feinleib M. Daily variation In USA mortality. *Am J Epidemiol* 1976. DOI:10.1093/oxfordjournals.aje.a112218.
- 51 Miller G. 'Airs, waters, and places' in history. *J Hist Med Allied Sci* 1962. DOI:10.1093/jhmas/XVII.1.129.
- 52 Shaw BD. Seasons of death: aspects of mortality in Imperial Rome. *J Rom Stud* 2006. DOI:10.2307/300425.
- 53 Healy JD. Excess winter mortality in Europe: a cross country analysis identifying key risk factors. *J Epidemiol Community Health* 2003; **57**: 784–9.
- 54 Reichert TA, Simonsen L, Sharma A, Pardo SA, Fedson DS, Miller MA. Influenza and the winter increase in mortality in the United States, 1959-1999. *Am J Epidemiol* 2004. DOI:10.1093/aje/kwh227.
- 55 Brown G, Fearn V, Wells C. Exploratory analysis of seasonal mortality in England and Wales, 1998 to 2007. *Heal Stat Q* 2010. DOI:10.1057/hsq.2010.21.
- 56 ONS. Excess winter mortality in England and Wales. Stat. Bull. 2019. <https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/deaths/bulletins/excesswintermortalityinenglandandwales/previousReleases>.
- 57 Rau R. Seasonality in human mortality. A demographic approach. *Wirtschafts- und Sozialwissenschaftlichen Fak* 2004; **PhD**: 361.
- 58 Hajat S, Kovats S. A note of caution about the excess winter deaths measure. *Nat Clim Chang* 2014.
- 59 Hajat S, Gasparrini A. The excess winter deaths measure: why its use is misleading for public health understanding of cold-related health impacts. *Epidemiology* 2016.

DOI:10.1097/EDE.0000000000000479.

- 60 Rosenwaike I. Seasonal variation of deaths in the United States, 1951–1960. *J Am Stat Assoc* 1966; **61**: 706–19.
- 61 Feinstein CA. Seasonality of deaths in the US by age and cause. *Demogr Res* 2002; **6**: 469–86.
- 62 Kalkstein AJ. Regional similarities in seasonal mortality across the United States: an examination of 28 metropolitan statistical areas. *PLoS One* 2013; **8**. DOI:10.1371/journal.pone.0063971.
- 63 Trudeau R. Monthly and daily patterns of death. *Stat Canada* 1997; **9**.
- 64 Gemmell I, McLoone P, Boddy FA, Dickinson GJ, Watt GCM. Seasonal variation in mortality in Scotland. *Int J Epidemiol* 2000. DOI:10.1093/ije/29.2.274.
- 65 Marti-Soler H, Gonseth S, Gubelmann C, *et al*. Seasonal variation of overall and cardiovascular mortality: a study in 19 countries from different geographic locations. *PLoS One* 2014. DOI:10.1371/journal.pone.0113500.
- 66 Laake K, Sverre JM. Winter excess mortality: a comparison between Norway and England plus Wales. *Age Ageing* 1996. DOI:10.1093/ageing/25.5.343.
- 67 Mackenbach JP, Kunst AE, Looman CWN. Seasonal variation in mortality in the Netherlands. *J Epidemiol Community Health* 1992; **46**: 261–5.
- 68 Green MS, Harari G, Kristal-Boneh E. Excess winter mortality from ischaemic heart disease and stroke during colder and warmer years in Israel: an evaluation and review of the role of environmental temperature. *Eur J Public Health* 2007. DOI:10.1093/eurpub/4.1.3.
- 69 Nakaji S, Parodi S, Fontana V, *et al*. Seasonal changes in mortality rates from main causes of death in Japan. *Eur J Epidemiol* 2004. DOI:10.1007/s10654-004-4695-8.
- 70 Rau R, Bohk-Ewald C, Muszyńska MM, Vaupel JW. Visualizing mortality dynamics in the Lexis diagram. 2018 DOI:10.1007/978-3-319-64820-0.
- 71 Dushoff J, Plotkin JB, Viboud C, Earn DJD, Simonsen L. Mortality due to influenza in the United States - an annualized regression approach using multiple-cause mortality data. *Am J Epidemiol* 2006. DOI:10.1093/aje/kwj024.
- 72 WHO. Influenza (seasonal) fact sheet. 2018. [https://www.who.int/news-room/fact-sheets/detail/influenza-\(seasonal\)](https://www.who.int/news-room/fact-sheets/detail/influenza-(seasonal)) (accessed May 20, 2019).
- 73 Nguyen JL, Yang W, Ito K, Matte TD, Shaman J, Kinney PL. Seasonal influenza infections and cardiovascular disease mortality. *JAMA Cardiol* 2016. DOI:10.1001/jamacardio.2016.0433.
- 74 Mercer JB. Cold - An underrated risk factor for health. *Environ. Res.* 2003. DOI:10.1016/S0013-9351(02)00009-9.
- 75 Simonsen L, Reichert TA, Viboud C, Blackwelder WC, Taylor RJ, Miller MA. Impact of influenza vaccination on seasonal mortality in the US elderly population. *Arch Intern Med* 2005; **165**: 265–72.
- 76 Simonsen L, Taylor RJ, Viboud C, Miller MA, Jackson LA. Mortality benefits of influenza vaccination in elderly people: an ongoing controversy. *Lancet Infect Dis* 2007. DOI:10.1016/S1473-3099(07)70236-0.
- 77 Rolfes MA, Flannery B, Chung J, *et al*. Effects of influenza vaccination in the United States during the 2017–2018 influenza Season. *Clin Infect Dis* 2019. DOI:10.1093/cid/ciz075.

- 78 Wu NC, Zost SJ, Thompson AJ, *et al.* A structural explanation for the low effectiveness of the seasonal influenza H3N2 vaccine. *PLoS Pathog* 2017. DOI:10.1371/journal.ppat.1006682.
- 79 Paules CI, Sullivan SG, Subbarao K, Fauci AS. Chasing seasonal influenza — the need for a universal influenza vaccine. *N Engl J Med* 2017. DOI:10.1056/nejmp1714916.
- 80 Ballester J, Rodó X, Robine J, Herrmann FR. European seasonal mortality and influenza incidence due to winter temperature variability. *Nat Clim Chang* 2016; **6**: 6–10.
- 81 Kinney PL, Schwartz J, Pascal M, *et al.* Winter season mortality: will climate warming bring benefits? *Environ Res Lett* 2015; **10**. DOI:10.1088/1748-9326/10/6/064016.
- 82 Taylor NAS. Human heat adaptation. *Compr Physiol* 2014. DOI:10.1002/cphy.c130022.
- 83 IPCC. Climate Change 2014 – impacts, adaptation and vulnerability: part B: regional aspects: working group II contribution to the IPCC Fifth Assessment Report. Cambridge University Press, 2014 DOI:doi:10.1017/CBO9781107415386.
- 84 Pfister C, Brázdil R, Glaser R, *et al.* Documentary evidence on climate in sixteenth-century Europe. *Clim Change* 1999. DOI:10.1023/A:1005540707792.
- 85 Luterbacher J, Dietrich D, Xoplaki E, Grosjean M, Wanner H. European seasonal and annual temperature variability, trends, and extremes since 1500. *Science* (80-) 2004. DOI:10.1126/science.1093877.
- 86 Semenza JC, Rubin CH, Falter KH, *et al.* Heat-related deaths during the July 1995 heat wave in Chicago. *N Engl J Med* 1996. DOI:10.1056/NEJM199607113350203.
- 87 Kaiser R, Le Tertre A, Schwartz J, Gotway CA, Daley WR, Rubin CH. The effect of the 1995 heat wave in Chicago on all-cause and cause-specific mortality. *Am J Public Health* 2007. DOI:10.2105/AJPH.2006.100081.
- 88 Garcia-Herrera R, Díaz J, Trigo RM, Luterbacher J, Fischer EM. A review of the European summer heat wave of 2003. *Crit Rev Environ Sci Technol* 2010. DOI:10.1080/10643380802238137.
- 89 Parsons K. Human thermal environments: the effects of hot, moderate, and cold environments on human health, comfort, and performance. CRC Press, 2014 DOI:10.1016/0003-6870(94)90062-0.
- 90 Bouchama A, Knochel JP. Heat stroke. *N Engl J Med* 2002; **346**. DOI:10.1002/cphy.c140017.
- 91 Kilbourne EM. Heat waves and hot environments. Oxford University Press, 1997.
- 92 UNISDR. Sendai Framework for Disaster Risk Reduction 2015 - 2030. *Third World Conf Disaster Risk Reduction, Sendai, Japan, 14-18 March 2015* 2015; : 1–25.
- 93 DHS. Extreme heat. ready.gov. 2019. <https://www.ready.gov/heat> (accessed May 8, 2019).
- 94 Borden KA, Cutter SL. Spatial patterns of natural hazards mortality in the United States. *Int J Health Geogr* 2008. DOI:10.1186/1476-072X-7-64.
- 95 NOAA. Weather fatalities 2018. 2018. <https://www.nws.noaa.gov/om/hazstats.shtml> (accessed May 20, 2019).
- 96 Shapiro E. 5 natural disasters that devastated the US in 2018. abc News. 2018. <https://abcnews.go.com/US/natural-disasters-devastated-us-2018/story?id=59367683> (accessed May 20, 2019).
- 97 EPA. Climate change indicators: heat-related deaths. 2017. <https://www.epa.gov/climate-indicators/climate-change-indicators-heat-related-deaths>

(accessed May 20, 2019).

- 98 Anderson BG, Bell ML. Weather-related mortality: how heat, cold, and heat waves affect mortality in the United States. *Epidemiology* 2009; **20**: 205.
- 99 Gasparrini A, Armstrong B. The impact of heat waves on mortality. *Epidemiol* 2011. DOI:10.1097/EDE.0b013e3181fdcd99.
- 100 Hajat S, Armstrong B, Baccini M, *et al*. Impact of high temperatures on mortality: is there an added heat wave effect? *Epidemiology* 2006. DOI:10.1097/01.ede.0000239688.70829.63.
- 101 Guo Y, Gasparrini A, Armstrong B, *et al*. Heat wave and mortality: a multicountry, multicomunity study. *Env Heal Perspect* 2017; : 1–11.
- 102 Barnett AG, Hajat S, Gasparrini A, Rocklöv J. Cold and heat waves in the United States. *Environ Res* 2012. DOI:10.1016/j.envres.2011.12.010.
- 103 The Eurowinter Group. Cold exposure and winter mortality from ischaemic heart disease, cerebrovascular disease, respiratory disease, and all causes in warm and cold regions of Europe. *Lancet* 1997; **349**: 1341–6.
- 104 Karl TR, Meehl G a, Miller CD, Hassol SJ, Waple A, Murray W. Weather and climate extremes in a changing climate regions of focus: North America, Hawaii, Caribbean, and U.S. Pacific Islands. 2008.
- 105 IPCC. Climate Change 2014 Synthesis Report. 2014.
- 106 Armstrong B. Models for the relationship between ambient temperature and daily mortality. *Epidemiology* 2006. DOI:10.1097/01.ede.0000239732.50999.8f.
- 107 Gasparrini A. Distributed lag linear and non-linear models in R: the package dlnm. *J Stat Softw* 2011.
- 108 Burke M, González F, Baylis P, *et al*. Higher temperatures increase suicide rates in the United States and Mexico. *Nat Clim Chang* 2018. DOI:10.1038/s41558-018-0222-x.
- 109 Shi L, Kloog I, Zanobetti A, Liu P, Schwartz JD. Impacts of temperature and its variability on mortality in New England. *Nat Clim Chang* 2015; **5**: 988–91.
- 110 Larsen U. The effects of monthly temperature fluctuations on mortality in the United States from 1921 to 1985. *Int J Biometeorol* 1990. DOI:10.1007/bf01048710.
- 111 Carmona R, Díaz J, Mirón IJ, Ortiz C, Luna MY, Linares C. Mortality attributable to extreme temperatures in Spain: a comparative analysis by city. *Environ Int* 2016. DOI:10.1016/j.envint.2016.02.018.
- 112 Le Tertre A, Lefranc AA, Eilstein D, *et al*. Impact of the 2003 heatwave on all-cause mortality in 9 French cities. *Epidemiology* 2006; **17**: 75–9.
- 113 Sheridan SC, Lee CC, Allen MJ. The mortality response to absolute and relative temperature extremes. *Environ Res public Heal* 2019.
- 114 Gasparrini A, Guo Y, Hashizume M, *et al*. Changes in susceptibility to heat during the summer: a multicountry analysis. *Am J Epidemiol* 2016; **183**: 1027–36.
- 115 Baccini M, Biggeri A, Accetta G, *et al*. Heat effects on mortality in 15 European cities. *Epidemiology* 2008. DOI:10.1097/EDE.0b013e318176bfd.
- 116 Stafoggia M, Forastiere F, Agostini D, *et al*. Vulnerability to heat-related mortality: a multicity, population-based, case-crossover analysis. *Epidemiology* 2006. DOI:10.1097/01.ede.0000208477.36665.34.
- 117 Gasparrini A, Armstrong B, Kovats S, Wilkinson P. The effect of high temperatures on

- cause-specific mortality in England and Wales. *Occup Environ Med* 2012. DOI:10.1136/oem.2010.059782.
- 118 Zanobetti A, Schwartz J. Temperature and mortality in nine US cities. *Epidemiology* 2008. DOI:10.1097/EDE.0b013e31816d652d.
 - 119 Eunice Lo Y, Mitchell D, Gasparri A, *et al.* Increasing mitigation ambition to meet the Paris Agreement's temperature goal avoids substantial heat-related mortality in U.S. cities. *Sci Adv* 2019.
 - 120 Bennett JE, Blangiardo M, Fecht D, Elliott P, Ezzati M. Vulnerability to the mortality effects of warm temperature in the districts of England and Wales. *Nat Clim Chang* 2014; **4**: 269–73.
 - 121 Basu R, Dominici F, Samet JM. Temperature and mortality among the elderly in the United States: a comparison of epidemiologic methods. *Epidemiology* 2005; **16**: 58–66.
 - 122 Basu R, Ostro BD. A multicounty analysis identifying the populations vulnerable to mortality associated with high ambient temperature in California. *Am J Epidemiol* 2008. DOI:10.1093/aje/kwn170.
 - 123 Ishigami A, Hajat S, Kovats RS, *et al.* An ecological time-series study of heat-related mortality in three European cities. *Environ Heal A Glob Access Sci Source* 2008. DOI:10.1186/1476-069X-7-5.
 - 124 Braga ALF, Zanobetti A, Schwartz J. The effect of weather on respiratory and cardiovascular deaths in 12 U.S. cities. *Environ Health Perspect* 2002; **110**: 859–63.
 - 125 Kolb S, Radon K, Valois MF, Héguy L, Goldberg MS. The short-term influence of weather on daily mortality in congestive heart failure. *Arch Environ Occup Heal* 2007. DOI:10.3200/AEOH.62.4.169-176.
 - 126 Kunst AE, Looman CWN, Mackenbach JP. Outdoor air temperature and mortality in the Netherlands: a time-series analysis. *Am J Epidemiol* 1993. DOI:10.1093/oxfordjournals.aje.a116680.
 - 127 Curriero FC, Heiner KS, Samet JM, Zeger SL, Strug L, Patz JA. Temperature and mortality in 11 cities of the eastern United States. *Am J Epidemiol* 2002; **155**: 80–7.
 - 128 Dilaveris P, Synetos A, Giannopoulos G, Gialafos E, Pantazis A, Stefanadis C. Climate impacts on myocardial infarction deaths in the Athens territory: the CLIMATE study. *Heart* 2006. DOI:10.1136/hrt.2006.091884.
 - 129 Wilmschurst P. Temperature and cardiovascular mortality. *BMJ* 1994. DOI:10.1136/bmj.309.6961.1029.
 - 130 Michelozzi P, Accetta G, De Sario M, *et al.* High temperature and hospitalizations for cardiovascular and respiratory causes in 12 European cities. *Am J Respir Crit Care Med* 2009. DOI:10.1164/rccm.200802-217OC.
 - 131 Keatinge WR, Coleshaw SRK, Easton JC, Cotter F, Mattock MB, Chelliah R. Increased platelet and red cell counts, blood viscosity, and plasma cholesterol levels during heat stress, and mortality from coronary and cerebral thrombosis. *Am J Med* 1986. DOI:10.1016/0002-9343(86)90348-7.
 - 132 Orru H, Åström DO. Increases in external cause mortality due to high and low temperatures: evidence from northeastern Europe. *Int J Biometeorol* 2017. DOI:10.1007/s00484-016-1270-4.
 - 133 Rey G, Jouglé E, Fouillet A, *et al.* The impact of major heat waves on all-cause and cause-specific mortality in France from 1971 to 2003. *Int Arch Occup Environ Health* 2007. DOI:10.1007/s00420-007-0173-4.

- 134 Xiang J, Bi P, Pisaniello D, Hansen A. Health impacts of workplace heat exposure: an epidemiological review. *Ind Health* 2014; **52**: 91–101.
- 135 Newell GR, Waggoner DE. Cancer mortality and environmental temperature in the United States. *Lancet* 1970. DOI:10.1016/S0140-6736(70)90988-8.
- 136 Medina-Ramón M, Zanobetti A, Cavanagh DP, Schwartz J. Extreme temperatures and mortality: assessing effect modification by personal characteristics and specific cause of death in a multi-city case-only analysis. *Environ Health Perspect* 2006. DOI:10.1289/ehp.9074.
- 137 Schwartz J. Who is sensitive to extremes of temperature? A case-only analysis. *Epidemiology* 2005. DOI:10.1097/01.ede.0000147114.25957.71.
- 138 Yan YY. The influence of weather on human mortality in Hong Kong. *Soc Sci Med* 2000. DOI:10.1016/S0277-9536(99)00301-9.
- 139 Glaser J, Lemery J, Rajagopalan B, *et al*. Climate change and the emergent epidemic of CKD from heat stress in rural communities: the case for heat stress nephropathy. *Clin J Am Soc Nephrol* 2016. DOI:10.2215/CJN.13841215.
- 140 Herath C, Jayasumana C, De Silva PMCS, De Silva PHC, Siribaddana S, De Broe ME. Kidney diseases in agricultural communities: a case against heat-stress nephropathy. *Kidney Int. Reports*. 2018. DOI:10.1016/j.ekir.2017.10.006.
- 141 Pearce N, Caplin B. Let's take the heat out of the CKDu debate: more evidence is needed. *Occup Environ Med* 2019. DOI:10.1136/oemed-2018-105427.
- 142 Burse RL. Sex differences in human thermoregulatory response to heat and cold stress. *Hum Factors J Hum Factors Ergon Soc* 1979. DOI:10.1177/001872087912210606.
- 143 Hajat S, Kovats RS, Lachowycz K. Heat-related and cold-related deaths in England and Wales: who is at risk? *Occup Environ Med* 2007. DOI:10.1136/oem.2006.029017.
- 144 Zanobetti A, O'Neill MS, Gronlund CJ, Schwartz JD. Summer temperature variability and long-term survival among elderly people with chronic disease. *Proc Natl Acad Sci* 2012. DOI:10.1073/pnas.1113070109.
- 145 Díaz J, Linares C, Tobías A. Impact of extreme temperatures on daily mortality in Madrid (Spain) among the 45-64 age-group. *Int J Biometeorol* 2006. DOI:10.1007/s00484-006-0033-z.
- 146 O'Neill MS, Zanobetti A, Schwartz J. Modifiers of the temperature and mortality association in seven US cities. *Am J Epidemiol* 2003. DOI:10.1093/aje/kwg096.
- 147 Bell ML, O'Neill MS, Ranjit N, Borja-Aburto VH, Cifuentes LA, Gouveia NC. Vulnerability to heat-related mortality in Latin America: a case-crossover study in São Paulo, Brazil, Santiago, Chile and Mexico City, Mexico. *Int J Epidemiol* 2008. DOI:10.1093/ije/dyn094.
- 148 Rainham DGC, Smoyer-Tomic KE. The role of air pollution in the relationship between a heat stress index and human mortality in Toronto. *Environ Res* 2003. DOI:10.1016/S0013-9351(03)00060-4.
- 149 Vaneckova P, Beggs PJ, de Dear RJ, McCracken KWJ. Effect of temperature on mortality during the six warmer months in Sydney, Australia, between 1993 and 2004. *Environ Res* 2008. DOI:10.1016/j.envres.2008.07.015.
- 150 El-Zein A, Tewtel-Salem M, Nehme G. A time-series analysis of mortality and air temperature in Greater Beirut. *Sci Total Environ* 2004. DOI:10.1016/j.scitotenv.2004.02.027.

- 151 Hajat S, Vardoulakis S, Heaviside C, Eggen B. Climate change effects on human health: projections of temperature-related mortality for the UK during the 2020s, 2050s and 2080s. *J Epidemiol Community Health* 2014. DOI:10.1136/jech-2013-202449.
- 152 Li T, Horton RM, Bader DA, *et al.* Aging will amplify the heat-related mortality risk under a changing climate: projection for the elderly in Beijing, China. *Sci Rep* 2016. DOI:10.1038/srep28161.
- 153 O'Neill MS, Hajat S, Zanobetti A, Ramirez-Aguilar M, Schwartz J. Impact of control for air pollution and respiratory epidemics on the estimated associations of temperature and daily mortality. *Int J Biometeorol* 2005. DOI:10.1007/s00484-005-0269-z.
- 154 Gouveia N, Hajat S, Armstrong B. Socioeconomic differentials in the temperature-mortality relationship in São Paulo, Brazil. *Int J Epidemiol* 2003. DOI:10.1093/ije/dyg077.
- 155 O'Neill MS, Zanobetti A, Schwartz J. Disparities by race in heat-related mortality in four US cities: the role of air conditioning prevalence. *J Urban Heal* 2005. DOI:10.1093/jurban/jti043.
- 156 Xu Y, Dadvand P, Barrera-Gómez J, *et al.* Differences on the effect of heat waves on mortality by sociodemographic and urban landscape characteristics. *J Epidemiol Community Health* 2013. DOI:10.1136/jech-2012-201899.
- 157 Rey G, Fouillet A, Bessemoulin P, *et al.* Heat exposure and socio-economic vulnerability as synergistic factors in heat-wave-related mortality. *Eur J Epidemiol* 2009. DOI:10.1007/s10654-009-9374-3.
- 158 Barnett AG, Tong S, Clements ACA. What measure of temperature is the best predictor of mortality? *Environ Res* 2010. DOI:10.1016/j.envres.2010.05.006.
- 159 Fanger P. Thermal comfort: Analysis and applications in environmental engineering. McGraw-Hill, New York, 1972 DOI:10.1016/s0003-6870(72)80074-7.
- 160 Buckley JP, Samet JM, Richardson DB. Does air pollution confound studies of temperature? *Epidemiology* 2014. DOI:10.1097/EDE.0000000000000051.
- 161 Ferreira Braga AL, Zanobetti A, Schwartz J. The time course of weather-related deaths. *Epidemiology* 2001; **12**: 662–7.
- 162 Saha M V., Davis RE, Hondula DM. Mortality displacement as a function of heat event strength in 7 US cities. *Am J Epidemiol* 2014. DOI:10.1093/aje/kwt264.
- 163 Armstrong B, Gasparrini A, Hajat S. Estimating mortality displacement during and after heat waves. *Am J Epidemiol* 2014. DOI:10.1093/aje/kwu083.
- 164 Dominici F. Time-series analysis of air pollution and mortality: a statistical review. *Res Rep Health Eff Inst* 2004.
- 165 Armstrong B, Gasparrini A, Tobias A. Conditional Poisson models: a flexible alternative to conditional logistic case cross-over analysis. *BMC Med Res Methodol* 2014. DOI:10.1186/1471-2288-14-122.
- 166 Maclure M. The case-crossover design: a method for studying transient effects on the risk of acute events. *Am J Epidemiol* 1991. DOI:10.1093/aje/kwx105.
- 167 Elliott P, Wakefield J, Best N, Briggs D. Spatial epidemiology: methods and applications. Oxford University Press, 2001 DOI:10.1093/ije/30.5.1204.
- 168 Brooke Anderson G, Bell ML. Heat waves in the United States: mortality risk during heat waves and effect modification by heat wave characteristics in 43 U.S. communities. *Environ Health Perspect* 2011. DOI:10.1289/ehp.1002313.

- 169 Madrigano J, Jack D, Anderson GB, Bell ML, Kinney PL. Temperature, ozone, and mortality in urban and non-urban counties in the northeastern United States. *Environ Heal A Glob Access Sci Source* 2015. DOI:10.1186/1476-069X-14-3.
- 170 Bobb JF, Dominici F, Peng RD. A Bayesian model averaging approach for estimating the relative risk of mortality associated with heat waves in 105 U.S. cities. *Biometrics* 2011. DOI:10.1111/j.1541-0420.2011.01583.x.
- 171 Aykroyd RG. Bayesian modeling of temperature-related mortality with latent functional relationships. *Commun. Stat. - Theory Methods*. 2018. DOI:10.1080/03610926.2017.1421223.
- 172 Baccini M, Kosatsky T, Analitis A, *et al*. Impact of heat on mortality in 15 European cities: attributable deaths under different weather scenarios. *J Epidemiol Community Heal* 2011; **65**: 64–70.
- 173 Yu W, Mengersen K, Wang X, *et al*. Daily average temperature and mortality among the elderly: a meta-analysis and systematic review of epidemiological evidence. *Int J Biometeorol* 2012. DOI:10.1007/s00484-011-0497-3.
- 174 Thomas A, Best N, Way R, *et al*. WinBUGS User Manual. *Components* 2003. DOI:http://www.mrc-bsu.cam.ac.uk/wp-content/uploads/manual14.pdf.
- 175 Depaoli S, Clifton JP, Cobb PR. Just Another Gibbs Sampler (JAGS): flexible software for MCMC implementation. *J Educ Behav Stat* 2016. DOI:10.3102/1076998616664876.
- 176 Stan Development Team. Stan: a C++ library for probability and sampling. 2014.
- 177 Rue H, Martino S, Chopin N. Approximate Bayesian inference for latent Gaussian models by using integrated nested Laplace approximations. *J R Stat Soc Ser B Stat Methodol* 2009. DOI:10.1111/j.1467-9868.2008.00700.x.
- 178 Kristensen K, Nielsen A, Berg CW, Skaug H, Bell BM. TMB: automatic differentiation and Laplace approximation. *J Stat Softw* 2016.
- 179 Mathers CD, Fat DM, Inoue M, Rao C, Lopez AD. Counting the dead and what they died from: an assessment of the global status of cause of death data. *Bull World Health Organ* 2005.
- 180 Boerma JT, Stansfield SK. Health statistics now: are we making the right investments? *Lancet*. 2007. DOI:10.1016/S0140-6736(07)60364-X.
- 181 Central statistics office. 2010 Census of population and housing. 2012.
- 182 Lind D. The citizenship question on the 2020 census, explained. Vox.com. 2018. <https://www.vox.com/policy-and-politics/2018/3/28/17168048/census-citizenship-2020-immigrants-count-trump-lawsuit> (accessed May 11, 2019).
- 183 Thomson-Deveaux A. The citizenship question could cost California And Texas a seat in congress. FiveThirtyEight. 2019. <https://fivethirtyeight.com/features/the-citizenship-question-could-cost-california-and-texas-a-seat-in-congress/> (accessed June 17, 2019).
- 184 United States Census Bureau. Methodology for the intercensal population and housing unit estimates: 2000 to 2010. 2012 <https://www2.census.gov/programs-surveys/popest/technical-documentation/methodology/intercensal/2000-2010-intercensal-estimates-methodology.pdf>.
- 185 Ingram DD, Parker JD, Schenker N, *et al*. United States Census 2000 population with bridged race categories. *Vital Health Stat* 2 2003; : 1–55.
- 186 United States Census Bureau. County intercensal tables 1980-1990. 2016. <https://www.census.gov/data/tables/time-series/demo/popest/1980s-county.html>.

- 187 Fowler T, Southgate RJ, Waite T, *et al.* Excess winter deaths in Europe: a multi-country descriptive analysis. *Eur J Public Health* 2015; **25**: 339–45.
- 188 Campbell A. Excess winter mortality in England and Wales: 2016 to 2017 (provisional) and 2015 to 2016 (final). *Stat Bull Off Natl Stat* 2017.
- 189 Ahmad OB, Boschi-Pinto C, Lopez AD, Murray CJ, Lozano R, Inoue M. Age standardization of rates: a new WHO standard. *WHO* 2000. DOI:10.1161/hypertensionaha.114.04394.
- 190 Messite J, Stellman SD. Accuracy of death certificate completion: the need for formalized physician training. *J Am Med Assoc* 1996. DOI:10.1001/jama.275.10.794.
- 191 Glasser JH. The quality and utility of death certificate data. *Am J Public Health* 1981. DOI:10.2105/AJPH.71.3.231.
- 192 Lu TH, Tsau SM, Wu TC. The Automated Classification of Medical Entities (ACME) system objectively assessed the appropriateness of underlying cause-of-death certification and assignment. *J Clin Epidemiol* 2005. DOI:10.1016/j.jclinepi.2005.03.017.
- 193 WHO. International Classification of Diseases 11th Revision. 2018. <https://icd.who.int/en/>.
- 194 WHO. ICD-10 International Statistical Classification of Diseases and Related Health Problems. 2011 DOI:10.1016/j.jclinepi.2009.09.002.
- 195 CDC. International Classification of Diseases, tenth revision, clinical modification (ICD-10-CM). 2018. <https://www.cdc.gov/nchs/icd/icd10cm.htm> (accessed April 12, 2019).
- 196 WHO. International Statistical Classification of Diseases and Related Health Problems 10th Revision. 2003. <http://apps.who.int/classifications/apps/icd/icd10online2003/fr-icd.htm> (accessed June 23, 2019).
- 197 Anderson RN, Miniño AM, Hoyert DL, Rosenberg HM. Comparability of cause of death between ICD-9 and ICD-10: preliminary estimates. *Natl Vital Stat Rep* 2001.
- 198 Foreman KJ, Lozano R, Lopez AD, Murray CJL. Modeling causes of death: an integrated approach using CODEm. *Popul Health Metr* 2012. DOI:10.1186/1478-7954-10-1.
- 199 Anderson RN, Miniño A, Hoyert D, Rosenberg H. Influence of rules from the tenth revision of the International Classification of Diseases on US cancer mortality trends. *J Natl Cancer Inst* 2003. DOI:10.1093/jnci/djg116.
- 200 WHO. History of the development of the ICD. WHO. 2019. <https://www.who.int/classifications/icd/en/HistoryOfICD.pdf> (accessed March 31, 2019).
- 201 D’Amico M, Agozzino E, Biagino A, Simonetti A, Marinelli P. Ill-defined and multiple causes on death certificates - a study of misclassification in mortality statistics. *Eur J Epidemiol* 1999. DOI:10.1023/A:1007570405888.
- 202 Lu TH, Anderson RN, Kawachi I. Trends in frequency of reporting improper diabetes-related cause-of-death statements on death certificates, 1985-2005: an algorithm to identify incorrect causal sequences. *Am J Epidemiol* 2010. DOI:10.1093/aje/kwq057.
- 203 McEwen LN, Karter AJ, Curb JD, Marrero DG, Crosson JC, Herman WH. Temporal trends in recording of diabetes on death certificates: results from Translating Research into Action for Diabetes (TRIAD). *Diabetes Care* 2011. DOI:10.2337/dc10-2312.

- 204 Naghavi M, Abajobir AA, Abbafati C, *et al.* Global, regional, and national age-sex specific mortality for 264 causes of death, 1980-2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet* 2017. DOI:10.1016/S0140-6736(17)32152-9.
- 205 Elo IT, Preston SH. Estimating African-American mortality from inaccurate data. *Demography* 2007. DOI:10.2307/2061751.
- 206 Perry M, Hollis D. The generation of monthly gridded datasets for a range of climatic variables over the UK. *Int J Climatol* 2005. DOI:10.1002/joc.1161.
- 207 NCDC. Climatology of the U.S. No. 81: monthly station normals of temperature, precipitation, and heating and cooling degree days 1971-2000. 2002.
- 208 African Ministerial Conference on Meteorology. Integrated African strategy. 2013 www.wmo.int/amcomet.
- 209 Compo GP, Whitaker JS, Sardeshmukh PD, *et al.* The twentieth century reanalysis project. *Q. J. R. Meteorol. Soc.* 2011. DOI:10.1002/qj.776.
- 210 Bengtsson L, Shukla J. Integration of space and in situ observations to study global climate change. *Bull Am Meteorol Soc* 2002. DOI:10.1175/1520-0477(1988)069<1130:iosais>2.0.co;2.
- 211 Trenberth KE, Olson JG. An evaluation and intercomparison of global analyses from the National Meteorological Center and the European Centre for Medium Range Weather Forecasts. *Bull Am Meteorol Soc* 2002. DOI:10.1175/1520-0477(1988)069<1047:aeaio>2.0.co;2.
- 212 Kalnay E, Kanamitsu M, Kistler R, *et al.* The NCEP/NCAR 40-year reanalysis project. *Bull Am Meteorol Soc* 1996. DOI:10.1175/1520-0477(1996)077<0437:TNYRP>2.0.CO;2.
- 213 Lorenc AC, Rawlins F. Why does 4D-Var beat 3D-Var? *Q J R Meteorol Soc* 2006. DOI:10.1256/qj.05.85.
- 214 Uppala SM, Kållberg PW, Simmons AJ, *et al.* The ERA-40 re-analysis. *Q. J. R. Meteorol. Soc.* 2005. DOI:10.1256/qj.04.176.
- 215 Dee DP, Uppala SM, Simmons AJ, *et al.* The ERA-Interim reanalysis: configuration and performance of the data assimilation system. *Q J R Meteorol Soc* 2011; **137**: 553–97.
- 216 Hoffmann L, Günther G, Li D, *et al.* From ERA-Interim to ERA5: the considerable impact of ECMWF's next-generation reanalysis on Lagrangian transport simulations. *Atmos Chem Phys* 2019. DOI:10.5194/acp-19-3097-2019.
- 217 Bennett JE, Tamura-Wicks H, Parks RM, *et al.* National and county life expectancy loss from particulate matter pollution in the USA (in press). *PLOS Med* 2019.
- 218 Daly C, Taylor G, Gibson W. The Prism approach to mapping precipitation and temperature. *10th AMS Conf Appl Climatol* 1997.
- 219 McKee CM. Deaths in winter: Can Britain learn from Europe? *Eur J Epidemiol* 1989; **5**: 178–82.
- 220 Langford IH, Bentham G. The potential effects of climate change on winter mortality in England and Wales. *Int J Biometeorol* 1995; **38**: 141–7.
- 221 Martens WJ. Climate change, thermal stress and mortality changes. *Soc Sci Med* 1998; **46**: 331–44.
- 222 Davis RE, Knappenberger PC, Michaels PJ, Novicoff WM. Seasonality of climate-

- human mortality relationships in US cities and impacts of climate change. *Clim Res* 2004; **26**: 61–76.
- 223 Medina-Ramón M, Schwartz J. Temperature, temperature extremes, and mortality: a study of acclimatisation and effect modification in 50 US cities. *Occup Environ Med* 2007; **64**: 827–33.
 - 224 Seretakis D. Changing seasonality of mortality from coronary heart disease. *JAMA J Am Med Assoc* 1997; **278**: 1012.
 - 225 Carson C, Hajat S, Armstrong B, Wilkinson P. Declining vulnerability to temperature-related mortality in London over the 20th century. *Am J Epidemiol* 2006; **164**: 77–84.
 - 226 Sheridan SC, Kalkstein AJ, Kalkstein LS. Trends in heat-related mortality in the United States, 1975–2004. *Nat Hazards* 2009; **50**: 145–60.
 - 227 Bobb JF, Peng RD, Bell ML, Dominici F. Heat-related mortality and adaptation to heat in the United States. *Environ Health Perspect* 2014; **122**: 811–6.
 - 228 Torrence C, Compo GP. A practical guide to wavelet analysis. *Bull Am Meteorol Soc* 1998. DOI:10.1175/1520-0477(1998)079<0061:APGTWA>2.0.CO;2.
 - 229 Hubbard BB. The world according to wavelets: the story of a mathematical technique in the making. A K Peters/CRC Press, 1998.
 - 230 Moy CM, Seltzer GO, Rodbell DT, Anderson DM. Variability of El Niño/Southern Oscillation activity at millennial timescales during the Holocene epoch. *Nature* 2002; **420**: 162–5.
 - 231 Grenfell BT, Bjørnstad ON, Kappey J. Travelling waves and spatial hierarchies in measles epidemics. *Nature* 2001; **414**: 716–23.
 - 232 Cazelles B, Chavez M, Berteaux D, *et al.* Wavelet analysis of ecological time series. *Oecologia*. 2008; **156**: 287–304.
 - 233 Fisher NI. Statistical analysis of circular data. Cambridge University Press, 1995.
 - 234 National Highway Traffic Safety Administration. Trend and pattern analysis of highway crash fatality by month and day. *Natl Cent Stat Anal* 2005.
 - 235 MacDorman MF, Gregory E. Fetal and perinatal mortality: United States, 2013. *Natl Vital Stat Reports* 2015; **64**: 1–24.
 - 236 Public Health England. The Cold Weather Plan for England. *Public Heal Engl* 2017.
 - 237 Lerchl A. Changes in the seasonality of mortality in Germany from 1946 to 1995: the role of temperature. *Int J Biometeorol* 1998; **42**: 84–8.
 - 238 Morris ME. Smart-Home Technologies to Assist Older People to Live Well at Home. *J Aging Sci* 2013; **01**. DOI:10.4172/2329-8847.1000101.
 - 239 Huang C, Barnett AG, Wang X, Vaneckova P, Fitzgerald G, Tong S. Projecting future heat-related mortality under climate change scenarios: a systematic review. *Environ Health Perspect* 2011. DOI:10.1289/ehp.1103456.
 - 240 Gasparrini A, Guo Y, Sera F, *et al.* Projections of temperature-related excess mortality under climate change scenarios. *Lancet Planet Heal* 2017. DOI:10.1016/S2542-5196(17)30156-0.
 - 241 IPCC. IPCC special report on the impacts of global warming of 1.5 °C - Summary for policy makers. 2018. <http://www.ipcc.ch/report/sr15/>.
 - 242 Rue H, Held L. Gaussian Markov random fields. Theory and applications. 2005 DOI:10.1007/s00184-007-0162-3.

- 243 Besag J. Spatial interaction and the statistical analysis of lattice systems. *J R Stat Soc Ser B (Statistical Methodol)* 1974. DOI:10.2307/2984812.
- 244 Nguyen JL, Schwartz J, Dockery DW. The relationship between indoor and outdoor temperature, apparent temperature, relative humidity, and absolute humidity. *Indoor Air* 2014; **24**: 103–12.
- 245 Spiegelhalter DJ, Best NG, Carlin BP, Van Der Linde A. Bayesian measures of model complexity and fit. *J R Stat Soc Ser B Stat Methodol* 2002; **64**: 583–616.
- 246 R Core Team. R: a language and environment for statistical computing. *R Found Stat Comput* 2012. DOI:ISBN 3-900051-07-0.
- 247 McMichael AJ, Woodruff RE, Hales S. Climate change and human health: present and future risks. *Lancet* 2006. DOI:10.1016/S0140-6736(06)68079-3.
- 248 Smith KR, Woodward A, Campbell-Lendrum D, *et al.* Human health: impacts, adaptation, and co-benefits. In: *Climate Change 2014 Impacts, Adaptation and Vulnerability: Part A: Global and Sectoral Aspects*. 2015. DOI:10.1017/CBO9781107415379.016.
- 249 Haines A, Ebi K. The imperative for climate action to protect health. *N Engl J Med* 2019; : 263–73.
- 250 Lozano R, Naghavi M, Foreman K, *et al.* Global and regional mortality from 235 causes of death for 20 age groups in 1990 and 2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 2012. DOI:10.1016/S0140-6736(12)61728-0.
- 251 Xu J. Unintentional drowning deaths in the United States, 1999-2010. *NCHS Data Brief* 2014.
- 252 Ambrose AF, Paul G, Hausdorff JM. Risk factors for falls among older adults: a review of the literature. *Maturitas* 2013. DOI:10.1016/j.maturitas.2013.02.009.
- 253 Bobb JF, Ho KKL, Yeh RW, *et al.* Time-course of cause-specific hospital admissions during snowstorms: an analysis of electronic medical records from major hospitals in Boston, Massachusetts. *Am J Epidemiol* 2017; **185**: 283–94.
- 254 Kelsey JL, Berry SD, Procter-Gray E, *et al.* Indoor and outdoor falls in older adults are different: the maintenance of balance, independent living, intellect, and zest in the elderly of Boston study. *J Am Geriatr Soc* 2010. DOI:10.1111/j.1532-5415.2010.03062.x.
- 255 Daanen HAM, Van De Vliert E, Huang X. Driving performance in cold, warm, and thermoneutral environments. *Appl Ergon* 2003. DOI:10.1016/S0003-6870(03)00055-3.
- 256 Zlatoper TJ. Determinants of motor vehicle deaths in the United States: a cross-sectional analysis. *Accid Anal Prev* 1991. DOI:10.1016/0001-4575(91)90062-A.
- 257 Mackie RR, Hanlon JF. O. A study of the combined effects of extended driving and heat stress on driver arousal and performance. In: *Symposium on relationships among theory, physiological correlates, and operational performance*. 1976. DOI:10.1007/978-1-4684-2529-1_25.
- 258 Wyon DP, Wyon I, Norin F. Effects of moderate heat stress on driver vigilance in a moving vehicle. *Ergonomics* 1996. DOI:10.1080/00140139608964434.
- 259 Opinium. Brits drink more alcohol in warmer weather. *Opinium.co.uk*. 2018. <https://www.opinium.co.uk/brits-drink-more-alcohol-in-warmer-weather/> (accessed Jan 10, 2019).
- 260 Voas RB, Torres P, Romano E, Lacey JH. Alcohol-related risk of driver fatalities: an

- update using 2007 data. *J Stud Alcohol Drugs* 2012. DOI:10.15288/jsad.2012.73.341.
- 261 Datla S, Sahu P, Roh H-J, Sharma S. A comprehensive analysis of the association of highway traffic with winter weather conditions. *Procedia - Soc Behav Sci* 2013. DOI:10.1016/j.sbspro.2013.11.143.
 - 262 Roh H-J, Sahu PK, Sharma S, Datla S, Mehran B. Statistical investigations of snowfall and temperature interaction with passenger car and truck traffic on primary highways in Canada. *J Cold Reg Eng* 2016. DOI:10.1061/(ASCE)CR.1943-5495.0000099.
 - 263 Roh H-J, Datla S, Sharma S. Effect of snow, temperature and their interaction on highway truck traffic. *J Transp Techn* 2013. DOI:10.4236/jtts.2013.31003.
 - 264 Roh HJ, Sharma S, Sahu PK. Modeling snow and cold effects for classified highway traffic volumes. *KSCE J Civ Eng* 2016. DOI:10.1007/s12205-015-0236-0.
 - 265 Graff Zivin J, Neidell M. Temperature and the allocation of time: implications for climate change. *J Labor Econ* 2014. DOI:10.1086/671766.
 - 266 Glaeser EL, Sacerdote B, Scheinkman JA. Crime and social interactions. *Q J Econ* 1996. DOI:10.2307/2946686.
 - 267 Rotton J, Cohn EG. Global warming and U.S. crime rates: an application of routine activity theory. *Environ Behav* 2003. DOI:10.1177/0013916503255565.
 - 268 Anderson CA. Temperature and aggression: ubiquitous effects of heat on occurrence of human violence. *Psychol Bull* 1989. DOI:10.1037/0033-2909.106.1.74.
 - 269 Baron RA, Bell PA. Aggression and heat: the influence of ambient temperature, negative affect, and a cooling drink on physical aggression. *J Pers Soc Psychol* 1976. DOI:10.1037/0022-3514.33.3.245.
 - 270 Majeed H, Lee J. The impact of climate change on youth depression and mental health. *Lancet Planet Heal* 2017; **1**: e94–5.
 - 271 Berry HL, Waite TD, Dear KBG, Capon AG, Murray V. The case for systems thinking about climate change and mental health. *Nat Clim Chang* 2018. DOI:10.1038/s41558-018-0102-4.
 - 272 Staddon PL, Montgomery HE, Depledge MH. Climate warming will not decrease winter mortality. *Nat Clim Chang* 2014. DOI:10.1038/nclimate2121.
 - 273 Ebi KL, Mills D. Winter mortality in a warming climate: a reassessment. *Wiley Interdiscip Rev Clim Chang* 2013; **4**: 203–12.
 - 274 Asthma UK. Weather. 2019. <https://www.asthma.org.uk/advice/triggers/weather/> (accessed June 4, 2019).
 - 275 Asthma UK. Moulds and fungi. 2019. <https://www.asthma.org.uk/advice/triggers/moulds-and-fungi/> (accessed June 4, 2019).