

**Anomalously warm temperatures are associated with increased injury deaths**

Robbie M Parks<sup>1,2</sup>, James E Bennett<sup>1,2,3</sup>, Helen Tamura-Wicks<sup>1,2</sup>, Vasilis Kontis<sup>1,2</sup>, Ralf Toumi<sup>4</sup>, Goodarz Danaei<sup>5</sup>, Majid Ezzati<sup>1,2,3,6\*</sup>

<sup>1</sup>MRC-PHE Centre for Environment and Health, Imperial College London, London, United Kingdom

<sup>2</sup>Department of Epidemiology and Biostatistics, School of Public Health, Imperial College London, London, United Kingdom

<sup>3</sup>Abdul Latif Jameel Institute for Disease and Emergency Analytics, Imperial College London, London, United Kingdom

<sup>4</sup>Space and Atmospheric Physics, Imperial College London, London, United Kingdom

<sup>5</sup>Harvard T.H. Chan School of Public Health, Boston, Massachusetts, USA

<sup>6</sup>WHO Collaborating Centre on NCD Surveillance and Epidemiology, Imperial College London, London, United Kingdom

\* Correspondence to: [majid.ezzati@imperial.ac.uk](mailto:majid.ezzati@imperial.ac.uk)

Temperatures which deviate from long-term local norm affect human health, and are projected to become more frequent as the global climate changes.<sup>1</sup> There is limited data on how such anomalies affect deaths from injuries. Here, we used data on mortality and temperature over 38 years (1980-2017) in the entire contiguous USA and formulated a Bayesian spatio-temporal model to quantify how anomalous temperatures, defined as deviations of monthly temperature from the local average monthly temperature over the entire analysis period, affect deaths from unintentional (transport, falls and drownings) and intentional (assault and suicide) injuries, by age group and sex. We found that a 1.5°C anomalously warm year, as envisioned under the Paris Climate Agreement,<sup>2</sup> would be associated with an estimated 1,601 (95% credible interval 1,430-1,776) additional injury deaths in the contiguous USA. 84% of these additional deaths would occur in males, mostly in adolescent to middle ages. These deaths would comprise of increases in deaths from drownings, transport, assault and suicide, offset partly by a decline in deaths from falls in older ages. The findings demonstrate the need for targeted interventions against injuries during periods of anomalously high temperatures, especially as these episodes increase with global climate change.

Anomalously warm and cold weather events are an important public health concern in today's world, and one of the key drivers for seeking adaptation measures against anthropogenic climate change.<sup>3-5</sup> Current assessments of the health effects of weather and climate, and by extension of global climate change, largely focus on parasitic and infectious diseases and cardiorespiratory and other chronic diseases.<sup>3-8</sup> Less research has been conducted on injuries,<sup>9-13</sup> especially in a consistent way across injury types and demographic subgroups of the population. There are two reasons to investigate a potential role for temperature anomalies on injury mortality: First death rates from injuries vary seasonally and the seasonality varies by

age group,<sup>14,15</sup> which motivates investigating whether temperature may play a role in their pathogenesis. Second, there are plausible behavioural and physiological pathways for a relationship between temperature and injury – for example changes in driving patterns and performance,<sup>12,16–24</sup> alcohol drinking,<sup>13</sup> and levels of anger<sup>25–27</sup> – which motivates testing whether injury deaths are affected by temperature anomalies. Our aim was to evaluate how deaths from various injuries in the USA may be affected by anomalously warm and temperatures that occur today and are expected to become increasingly common as a result of global climate change.<sup>1</sup>

We used vital registration data on all injury deaths in the contiguous USA (i.e., excluding Alaska and Hawaii) from 1980 to 2017, with information on sex, age at death, underlying cause of death and county and state of residence. From 1980 to 2017, 4,145,963 boys and men and 1,825,817 girls and women died from an injury in the contiguous USA, accounting for 9.3% and 4.2% of all male and female deaths respectively. 95.7% of male injury deaths and 94% of female injury deaths were in those aged 15 years and older, and over half (52.3%) of male injury deaths were in those aged 15–44 years (Figure 1). In contrast with males, there was less of an age gradient in females after 15 years of age.

Injuries from transport, falls, drownings, assault, and suicide accounted for 78.6% of injury deaths in males and 71.8% in females. The remainder were from a heterogeneous group of “other injuries” (Figure 1), within which the composition of injuries that led to death varied by sex and age group. Transport was the leading injury cause of death in women younger than 75 years and men younger than 35 years. Between 35 and 74 years of age, more men died of suicide than any other injury. Above 75 years of age, falls were the largest cause of death in both men and women.

68

69 There was a decline in age-standardised death rates of three out of five major injuries (transport,  
70 drownings and assault) from 1980 to 2017, although assault deaths have more recently (since  
71 2014) increased (Figure 2). In contrast, age-standardised death rates from falls increased over  
72 time while those from suicide initially decreased followed by an increase to surpass 1980  
73 levels. The largest overall decline over time was for transport deaths in both sexes and for  
74 deaths from drownings in men, which declined by over 50% from 1980 to 2017. Age-  
75 standardised death rates for transport injuries and drownings peaked in summer months but  
76 deaths from other major injuries did not have clear seasonal patterns.

77

78 We defined a measure of anomalous temperature for each county and month, which represents  
79 the deviation from the county's average temperature in that month over the entire analysis  
80 period (Extended Data Figure 1). County-level anomalies were aggregated to state level with  
81 use of population weights. Average size of anomaly over the study period (1980-2017), a  
82 measure of how variable temperatures are around their state-month long-term average, ranged  
83 from 0.4°C for Florida in September, to 3.4°C for North Dakota in February (Extended Data  
84 Figure 2). The average size of anomaly had a median value of 1.2°C across all states and  
85 months (Extended Data Figure 2). Temperature anomalies were largest in January and  
86 December and smallest in August and September. They were larger in northern and central  
87 states than in southern and coastal ones.

88

89 We analysed the association of monthly injury death rates with anomalous temperature using  
90 a Bayesian spatio-temporal model, described in detail in Methods. We used the resultant risk  
91 estimates and the age-sex-specific death rates from each injury in 2017, to estimate additional  
92 deaths if each month in each state were +1.5°C above its long-term average as envisioned under

the Paris Climate Agreement. We present additional results, based on a +2°C, which is the upper bound of the Paris Climate Agreement as Extended Data. Based on this analysis, there would be an estimated 1,601 (95% credible interval 1,430-1,776) excess injury deaths, equivalent to 0.75% of all injury deaths in 2017, in a year in which each month in each state was +1.5°C warmer than its long-term average (Figure 3). The number of excess injury deaths would increase to 2,135 (95% credible interval 1,906-2,368), equivalent to 1.0% of all injury deaths in 2017, in each year in which each month in each state was +2°C warmer than its long-term average (Extended Data Figure 3).

Deaths from drowning, transport, assault and suicide would increase, partly offset by a decline in deaths from falls in middle and older ages and in winter months (Figure 3). Most excess deaths would be from transport injuries (739; 650-814 in the +1.5°C warmer scenario) followed closely by suicide (540; 445-631). 84% of the excess deaths would occur in males and 16% in females. 92% of all male excess deaths would occur in those aged 15-64 years, who have higher rates of deaths from transport and suicide. In those aged 85 years and older, there would be an estimated decline in injury deaths, because deaths from falls are expected to decline in a warmer year.

Proportionally, deaths from drownings are estimated to increase more than those of other injury types, by as much 13.7% (12.5, 15.2) for a +1.5°C anomaly in men aged 15-24 years (Figure 4). The smallest proportional increase was that of assault and suicide (less than 3% in all age and sex groups). There was a larger percent increase in transport deaths for males than for females, especially in young and middle-ages (e.g., 2.0% (1.6, 2.6) for 25-34 year old men versus 0.5% (-0.3, 1.4) for women of the same age) (Figure 4).

That anomalously warm temperature influences deaths from drowning, although not previously quantified, is highly plausible because swimming is likely to be more common when temperature is higher. The higher relative and absolute impacts on men compared with women may reflect differences in their behaviours. For example, over half of swimming deaths for males occur in natural water, compared to about one quarter for females.<sup>28</sup> The former may rise more in warmer weather. Similarly, deaths from falls declined more in older ages because falls in the elderly are more likely to be due to slipping on ice than those in younger people.<sup>29–</sup>

<sup>31</sup>

The pathways from anomalous temperature to transport injury are more varied. Firstly, driving performance deteriorates at higher temperatures.<sup>20–23</sup> Further, alcohol consumption increases in warm temperatures,<sup>13</sup> which also provides an explanation for why teenagers, who are more likely than other age groups to crash while intoxicated,<sup>32</sup> experience a larger proportional rise in deaths from transport when temperatures are anomalously warm than older adults. Lastly, warmer temperatures generally increase road traffic in North America;<sup>12,16–19,24</sup> coupled with more people outdoors in warmer weather,<sup>33</sup> this increase could lead to more fatal collisions.

Pathways linking anomalously high temperatures and deaths from assault and suicide are less established. One hypothesis is that, more time spent outdoors in anomalously warmer temperatures leads to an increased number of face-to-face interactions, and hence arguments, confrontations, and ultimately assaults.<sup>34,35</sup> These effects could be compounded by the greater anger levels linked to higher temperatures.<sup>25–27</sup> However, further research on the association of temperature and assault, and the factors mediating it, is needed.<sup>36</sup> Regarding suicide, it has been hypothesised that higher temperature is associated with higher levels of distress in younger people.<sup>37</sup> Nonetheless, the mechanisms for the links between temperature and mental health

143 requires further investigation, including whether the relationship varies by age and sex, as  
144 indicated by our results. Future research should also investigate the extent to which the  
145 increased risk of injury death as a result of anomalous temperature depends on community  
146 characteristics such as poverty and deprivation, social connectivity and cohesion, quality of  
147 roads and housing, public transportation options, emergency response, and social services.

148  
149 Our work highlights how deaths from injuries are currently susceptible to temperature  
150 anomalies and could also be modified by rising temperatures resulting from climate change,  
151 unless countered by social infrastructure and health system interventions that mitigate these  
152 impacts. Though absolute impacts on mortality are modest, some groups, especially men in  
153 young to middle-ages, experience larger impacts. Therefore, a combination of public health  
154 interventions that broadly target injuries in these groups – for example targeted messaging for  
155 younger males on the risks of transport injury and drowning – and those that trigger in relation  
156 to forecasted high temperature periods – for example additional targeted blood alcohol level  
157 checks – should be a public health priority.

## 158    **References**

- 159    1.    IPCC. *Climate change 2013: the physical science basis. Working group I contribution*  
160        *to the fifth assessment report of the intergovernmental panel on climate change.*  
161        (Cambridge, UK and New York, USA: Cambridge University Press, 2013).
- 162    2.    United Nations. *Paris Agreement*. (2015).
- 163    3.    McMichael, A. J., Woodruff, R. E. & Hales, S. Climate change and human health:  
164        present and future risks. *Lancet* (2006). doi:10.1016/S0140-6736(06)68079-3
- 165    4.    Smith, K. R. *et al.* Human health: impacts, adaptation, and co-benefits. in *Climate*  
166        *Change 2014 Impacts, Adaptation and Vulnerability: Part A: Global and Sectoral*  
167        *Aspects* (2015). doi:10.1017/CBO9781107415379.016
- 168    5.    Haines, A. & Ebi, K. The imperative for climate action to protect health. *N. Engl. J.*  
169        *Med.* 263–273 (2019).
- 170    6.    Watts, N. *et al.* The 2018 report of the Lancet Countdown on health and climate  
171        change: shaping health of nations for centuries to come. *Lancet* **6736**, 1–4 (2018).
- 172    7.    Huang, C. *et al.* Projecting future heat-related mortality under climate change  
173        scenarios: a systematic review. *Environ. Health Perspect.* (2011).  
174        doi:10.1289/ehp.1103456
- 175    8.    Gasparrini, A. *et al.* Projections of temperature-related excess mortality under climate  
176        change scenarios. *Lancet Planet. Heal.* (2017). doi:10.1016/S2542-5196(17)30156-0
- 177    9.    Burke, M. *et al.* Higher temperatures increase suicide rates in the United States and  
178        Mexico. *Nat. Clim. Chang.* (2018). doi:10.1038/s41558-018-0222-x
- 179    10.    Rey, G. *et al.* The impact of major heat waves on all-cause and cause-specific  
180        mortality in France from 1971 to 2003. *Int. Arch. Occup. Environ. Health* (2007).  
181        doi:10.1007/s00420-007-0173-4
- 182    11.    Orru, H. & Åström, D. O. Increases in external cause mortality due to high and low  
183        temperatures: evidence from northeastern Europe. *Int. J. Biometeorol.* (2017).  
184        doi:10.1007/s00484-016-1270-4
- 185    12.    Robertson, L. S. Reversal of the road death trend in the U.S. in 2015–2016: An  
186        examination of the climate and economic hypotheses. *J. Transp. Heal.* (2018).  
187        doi:10.1016/j.jth.2018.04.005
- 188    13.    Opinium. Brits drink more alcohol in warmer weather. *Opinium.co.uk* (2018).  
189        Available at: [https://www.opinium.co.uk/brits-drink-more-alcohol-in-warmer-](https://www.opinium.co.uk/brits-drink-more-alcohol-in-warmer-weather/)  
190        [weather/](https://www.opinium.co.uk/brits-drink-more-alcohol-in-warmer-weather/). (Accessed: 10th January 2019)
- 191    14.    Parks, R. M., Bennett, J. E., Foreman, K. J., Toumi, R. & Ezzati, M. National and  
192        regional seasonal dynamics of all-cause and cause-specific mortality in the USA from  
193        1980 to 2016. *Elife* **7**, (2018).
- 194    15.    Rau, R. Seasonality in human mortality. A demographic approach. *Wirtschafts- und*  
195        *Sozialwissenschaftlichen Fak. PhD*, 361 (2004).
- 196    16.    Datla, S., Sahu, P., Roh, H.-J. & Sharma, S. A comprehensive analysis of the  
197        association of highway traffic with winter weather conditions. *Procedia - Soc. Behav.*  
198        *Sci.* (2013). doi:10.1016/j.sbspro.2013.11.143
- 199    17.    Roh, H.-J., Sahu, P. K., Sharma, S., Datla, S. & Mehran, B. Statistical investigations of  
200        snowfall and temperature interaction with passenger car and truck traffic on primary  
201        highways in Canada. *J. Cold Reg. Eng.* (2016). doi:10.1061/(ASCE)CR.1943-  
202        5495.0000099
- 203    18.    Roh, H.-J., Datla, S. & Sharma, S. Effect of snow, temperature and their interaction on  
204        highway truck traffic. *J. Transp. Techn.* (2013). doi:10.4236/jtts.2013.31003
- 205    19.    Roh, H. J., Sharma, S. & Sahu, P. K. Modeling snow and cold effects for classified  
206        highway traffic volumes. *KSCE J. Civ. Eng.* (2016). doi:10.1007/s12205-015-0236-0



20. Daanen, H. A. M., Van De Vliert, E. & Huang, X. Driving performance in cold, warm, and thermoneutral environments. *Appl. Ergon.* (2003). doi:10.1016/S0003-6870(03)00055-3
21. Zlatoper, T. J. Determinants of motor vehicle deaths in the United States: a cross-sectional analysis. *Accid. Anal. Prev.* (1991). doi:10.1016/0001-4575(91)90062-A
22. Mackie, R. R. & Hanlon, J. F. . O. A study of the combined effects of extended driving and heat stress on driver arousal and performance. in *Symposium on relationships among theory, physiological correlates, and operational performance* (1976). doi:10.1007/978-1-4684-2529-1\_25
23. Wyon, D. P., Wyon, I. & Norin, F. Effects of moderate heat stress on driver vigilance in a moving vehicle. *Ergonomics* (1996). doi:10.1080/00140139608964434
24. Robertson, L. Climate change, weather and road deaths. *Inj. Prev.* (2018). doi:10.1136/injuryprev-2017-042419
25. Anderson, C. A. Temperature and aggression: ubiquitous effects of heat on occurrence of human violence. *Psychol. Bull.* (1989). doi:10.1037/0033-2909.106.1.74
26. Baron, R. A. & Bell, P. A. Aggression and heat: the influence of ambient temperature, negative affect, and a cooling drink on physical aggression. *J. Pers. Soc. Psychol.* (1976). doi:10.1037/0022-3514.33.3.245
27. Talaei, A., Hedjazi, A., Rezaei Ardani, A., Fayyazi Bordbar, M. R. & Talaei, A. The relationship between meteorological conditions and homicide, suicide, rage, and psychiatric hospitalization. *J. Forensic Sci.* (2014). doi:10.1111/1556-4029.12471
28. Xu, J. Unintentional drowning deaths in the United States, 1999-2010. *NCHS Data Brief* (2014).
29. Ambrose, A. F., Paul, G. & Hausdorff, J. M. Risk factors for falls among older adults: a review of the literature. *Maturitas* (2013). doi:10.1016/j.maturitas.2013.02.009
30. Bobb, J. F. *et al.* Time-course of cause-specific hospital admissions during snowstorms: an analysis of electronic medical records from major hospitals in Boston, Massachusetts. *Am. J. Epidemiol.* **185**, 283–294 (2017).
31. Kelsey, J. L. *et al.* Indoor and outdoor falls in older adults are different: the maintenance of balance, independent living, intellect, and zest in the elderly of Boston study. *J. Am. Geriatr. Soc.* (2010). doi:10.1111/j.1532-5415.2010.03062.x
32. Voas, R. B., Torres, P., Romano, E. & Lacey, J. H. Alcohol-related risk of driver fatalities: an update using 2007 data. *J. Stud. Alcohol Drugs* (2012). doi:10.15288/jsad.2012.73.341
33. Graff Zivin, J. & Neidell, M. Temperature and the allocation of time: implications for climate change. *J. Labor Econ.* (2014). doi:10.1086/671766
34. Glaeser, E. L., Sacerdote, B. & Scheinkman, J. A. Crime and social interactions. *Q. J. Econ.* (1996). doi:10.2307/2946686
35. Rotton, J. & Cohn, E. G. Global warming and U.S. crime rates: an application of routine activity theory. *Environ. Behav.* (2003). doi:10.1177/0013916503255565
36. Mach, K. J. *et al.* Climate as a risk factor for armed conflict. *Nature* (2019). doi:10.1038/s41586-019-1300-6
37. Majeed, H. & Lee, J. The impact of climate change on youth depression and mental health. *Lancet Planet. Heal.* **1**, e94–e95 (2017).

## Methods

### *Data sources*

We used data on deaths by sex, age, underlying cause of death and state of residence in the contiguous USA from 1980 to 2017 through the National Center for Health Statistics (NCHS) ([https://www.cdc.gov/nchs/nvss/dvs\\_data\\_release.htm](https://www.cdc.gov/nchs/nvss/dvs_data_release.htm)) and on population from the NCHS bridged-race dataset for 1990 to 2017 ([https://www.cdc.gov/nchs/nvss/bridged\\_race.htm](https://www.cdc.gov/nchs/nvss/bridged_race.htm)) and from the US Census Bureau prior to 1990 (<https://www.census.gov/data/tables/time-series/demo/popest/1980s-county.html>). We did not include Alaska and Hawaii, (which together made up 0.5% of the US population in 2017) because their climates and environment are distinct from other states due to their substantial physical distance. We calculated monthly population counts through linear interpolation, assigning each yearly count to July.

The underlying cause of death was coded according to the international classification of diseases (ICD) system (9<sup>th</sup> revision from 1980 to 1998 and 10<sup>th</sup> revision thereafter). The 6 million injury deaths fell into six categories: transport, falls, drownings, assault, suicide and an aggregate set of other injuries (Extended Data Table 1). We report the results of all of these categories except other injuries (1,402,941 deaths or 23% of total injury deaths during 1980-2017), because the composition of this aggregate group varies by sex, age group, state and time.

We obtained data on temperature from ERA5, which uses data from global in-situ and satellite measurements to generate a worldwide meteorological dataset, with full space and time coverage over our analysis period.<sup>38</sup> We used gridded four-times-daily estimates at a resolution of 30 km to generate monthly temperatures by county.

### *Anomalous temperature metric*

With few exceptions,<sup>9,39</sup> current climate change risk assessments extrapolate from associations of daily mortality with daily temperature.<sup>7,8,40–42</sup> Climate change, however, will fundamentally modify weather, including seasonal weather patterns, compared to long-term averages, and hence can disrupt existing forms of adaptation. To mimic the conditions that may arise with global climate change, we developed methodology to examine how deviations from long-term average temperature may impact injury death rates.

We first defined a measure of anomalous temperature for each county and month, which represents the deviation from the average temperature of the county in that month over the entire analysis period (Extended Data Figure 1). To calculate the magnitude of temperature anomaly, we first calculated average temperatures for each month in each county over the entire 38 years of analysis. We subtracted these long-term average temperatures from respective monthly temperature values to generate a temperature anomaly time series for each month and year in each county (Extended Data Figure 1). The temperature anomaly metric measures the extent that temperature experienced in a specific month, year and county is warmer or cooler than the long-term average to which the population has acclimatised. These values can be different for different months in the same county, and different counties in the same month. Further, a county with higher, but more stable, temperature in a specific month has smaller anomalies than one with lower but more inter-annually variable temperature. County-level anomalies were aggregated to state level with use of population weights for analysing their associations with mortality.

### *Statistical methods*

We analysed the association of monthly injury death rates with anomalous temperature using a Bayesian spatio-temporal model, which leveraged variations over space and time to infer

associations. We modelled the number of deaths in each month in each year as following a Poisson distribution:

$$deaths_{state-time} \sim \text{Poisson}(death\ rate_{state-time} \cdot population_{state-time})$$

with log-transformed death rates modelled as a sum of components that depend on location (state) of death, month of year, overall time (in months) and temperature anomaly:

$$\begin{aligned} \log(death\ rate_{state-time}) = & \\ & \alpha_0 + \beta_0 \cdot time + \\ & \alpha_{state} + \beta_{state} \cdot time + \\ & \alpha_{month} + \beta_{month} \cdot time + \\ & \zeta_{state-month} + \\ & \psi_{state-month} \cdot time + \\ & \nu_{time} + \\ & \gamma_{month} \cdot Anomaly_{state-time} + \\ & \epsilon_{state-time} \end{aligned}$$

The model contained terms that represent the national level and trend in mortality, with  $\alpha_0$  as the common intercept and  $\beta_0$  the common slope with overall time. Death rates also vary by month, which may be partly related to temperature and partly due to other monthly factors; monthly variations tend to be smooth across adjacent months.<sup>14</sup> Therefore, we allowed each month of the year to systematically have a different mortality level and trend, with  $\alpha_{month}$  the month-specific intercept and  $\beta_{month}$  the month-specific slope with overall time. We used a first-order random walk prior for the monthly random intercepts and slopes, widely used to characterise smoothly varying trends.<sup>43</sup> The random walk had a cyclic structure, so that December was adjacent to January.

We also included state random intercepts and slopes for death rates, with  $\alpha_{state}$  as the state-specific intercept and  $\beta_{state}$  the state-specific slope with overall time. These terms measure deviations of each state from national values, and allow variation in level and trend in mortality by state. We modelled the state-level random intercepts and slopes using the Besag, York, and Mollie (BYM) spatial model,<sup>44</sup> which includes both spatially-structured random effects with

an intrinsic Conditional Autoregressive (ICAR) prior and spatially-unstructured independent and identically distributed (IID) Gaussian random effects. The extent to which information is shared between neighbouring states depends on the uncertainty of death rates in a state and the empirical similarity of death rates in neighbouring states. We also included state-month interactions for intercepts and slopes ( $\zeta_{state-month}$  and  $\psi_{state-month}$ ), to allow variation in mortality levels and trends in a particular state for different months and vice-versa. These state-month interactions were modelled as IID and therefore were of Type I space-time interactions.<sup>45</sup> Non-linear change over overall time (in months) was captured by a first-order random walk,  $v_{time}$ .<sup>43</sup> In order to ensure identifiability each set of random walk terms or state random effects was constrained to sum to zero.

Finally, we included a term that relates log-transformed death rate to the above-defined state-month temperature anomaly,  $\gamma_{month} \cdot Anomaly_{state-time}$ . The coefficients of  $\gamma_{month}$  represent the logarithm of the monthly death rate ratio per 1°C increase in anomaly. There was a separate coefficient for each month which means that an anomaly of the same magnitude could have different associations with injury mortality in different months. As with the month-specific intercepts and trends, we used a cyclic first-order random walk to smooth the coefficient of the temperature anomaly across months. An over-dispersion term ( $\epsilon_{state-time}$ ) captured the variation unaccounted for by other terms in the model, modelled as  $N(0, \sigma_{\epsilon}^2)$ . We used weakly informative priors so that parameter estimation was driven by the data. As in previous analyses,<sup>46,47</sup> hyper-priors were defined on the logarithm of the precisions of the random effects, in other words on  $\log(1/\sigma^2)$ . These were modelled as  $\log\text{Gamma}(\theta, \delta)$  distributions with shape  $\theta = 1$  and rate  $\delta = 0.001$ . The same hyper-priors were used for all precision parameters of the random effects in the model. For the common slope, we used  $N(0, 1000)$  and for the common intercept a flat prior.

In addition to representing the spatial (across states) and temporal (across months and years) patterns of mortality, the intercept terms ( $\alpha_{month}$ ,  $\alpha_{state}$ ,  $\zeta_{state-month}$ ) in our statistical model implicitly adjust for unobserved factors that influence mortality at the state, month and state-month level; the slope terms ( $\beta_{month}$ ,  $\beta_{state}$ ,  $\psi_{state-month}$ ) do so for changes in these factors over time.<sup>46</sup> This means that the only confounding factors would be those that have the same state-month anomaly as temperature.

We fitted the models using integrated nested Laplace approximation (INLA), using the R-INLA software, which offers orders of computational efficiency improvement in Bayesian inference compared to traditional MCMC.<sup>48</sup> The uncertainty in our results were obtained from 5000 draws from the posterior marginal of each month's excess relative risk. The reported 95% credible intervals are the 2.5<sup>th</sup> to 97.5<sup>th</sup> percentiles of the sampled values.

Analyses were done separately by injury type, because different injuries can have differing associations with anomalously warm and cold temperature. Analyses were also done separately by sex and age group (0-4 years, 10-year age groups from 5 to 84 years, and 85+ years) because injury death rates vary by age group and sex (Figure 1 and Extended Data Table 2), as might their associations with temperature. We used the resultant risk estimates and the age-sex-specific death rates from each injury in 2017, to calculate additional deaths if each month in each state were +1.5°C above its long-term average, not only realistic in our lifetimes under current projections of global climate change but an agreed upper bound chosen under the Paris Climate Agreement.<sup>2,49</sup> +1.5°C is also within the range of anomaly size experienced by some states (Extended Data Figure 2). We did similar calculations for +2°C, which is the upper bound of the Paris Climate Agreement, and present these as Extended Data. For these

calculations, we multiplied the actual death counts for each month, sex, state and age group in 2017 by the corresponding excess relative risk, which was calculated as the exponential of the coefficient of the temperature anomaly term from the above analysis.

### *Sensitivity analyses*

We conducted sensitivity analyses to assess how much our results might depend on the temperature metric used to generate anomalous temperature. First, instead of building our monthly temperature anomalies based on daily mean temperatures, we used daily maxima and minima. These measures were strongly correlated to those generated from daily means (Extended Data Table 3), and therefore we did not run models using these alternatives.

Second, together with temperature anomaly based on daily mean temperatures, we also included a second measure of anomaly in the model. The additional measures were related to more extreme anomalous situations which may be relevant if the impacts on injuries are related to more extreme temperatures and how frequent they are in each month:

- temperature anomaly calculated based on 90<sup>th</sup> percentile (°C) of daily mean temperatures within a month, compared to the average of 90<sup>th</sup> percentiles for each state and month
- number of days in a month above the long-term 90<sup>th</sup> percentile of average temperature for each state and month (adjusted for length of month)
- number of 3+ day episodes above the long-term 90<sup>th</sup> percentile of average temperature for each state and month (adjusted for length of month)

The correlations among these variables and anomaly based on mean were between 0.60 and 0.89 (Extended Data Table 4). The estimated rate ratios of temperature anomaly based on daily means (i.e., the anomaly measure used in the main analysis) were robust to the addition of

alternative measures of anomaly, while the coefficients of the additional measures were generally not significant and with large credible intervals. Therefore, we did not include the alternative additional measures of extreme anomalous temperature in the main analysis.

### *Comparison with prior studies*

While there are no previous studies of how deviations of monthly temperature from long-term average are associated with injury mortality, our results are broadly in agreement with those that have analysed associations with absolute temperature and for specific injury types. A study of suicide in US counties over 37 years (1968-2004) estimated that 1°C higher monthly temperature would lead to a 0.7% rise in suicides,<sup>9</sup> compared to our findings of 0.7-1.5% in males and 0.5-2.9% in females in different ages for a +1.5°C anomaly. A cross-sectional analysis in 100 US counties found that a 1°C higher temperature would lead to a 1.3% increase in death rates from road traffic injuries,<sup>24</sup> compared to our finding of 0.6-3.1% in males and 0.5-2.0% in females for a +1.5°C anomaly. In a study of six French heatwaves during 1971-2003, mortality from unintentional injuries rose by up to 4% during a heatwave period compared to a non-heatwave baseline.<sup>10</sup> A study of daily mortality from all injuries from Estonia found a 1.24% increase in mortality when daily maximum temperature went from the 75<sup>th</sup> to 99<sup>th</sup> percentile of long-term distribution.<sup>11</sup>

### *Strengths and limitations*

The major strength of our study is that we have comprehensively modelled the association of temperature anomaly with injury by type of injury, month, age group and sex. Our measure of temperature anomaly internalises long-term historical experience of each state, and is closer to what climate change may bring about than solely examining daily episodes, or average temperature to which people have adapted. To utilise this metric, we integrated two large



434 disparate national datasets on mortality (US vital statistics) and meteorology (ERA5), and  
435 developed a bespoke Bayesian spatio-temporal model. A limitation of our study is that, like all  
436 observation studies, we cannot rule out confounding of results due to other factors. As  
437 described above, our statistical model by design adjusts for factors related to month, state and  
438 state-month that are either invariant over time or that change linearly. Rather, the confounding  
439 factors would be those with anomalies that are similar to those of monthly temperature in each  
440 state, such as air pollution. However, to our knowledge, there is currently no evidence of an  
441 association between air pollution and injury mortality. We analysed the associations between  
442 anomalous temperature and injury mortality at the state level because the small number of  
443 events and computational demands made county-level analyses unfeasible. Analyses at finer  
444 spatial resolution, such as county or district,<sup>50</sup> would be ideal because the impacts of  
445 anomalously warm and cold temperature on deaths from injuries may depend on  
446 socioeconomic (e.g., poverty; social connectivity and cohesion; availability of guns),  
447 environmental (e.g., availability of swimming pools; distance to bodies of water), infrastructure  
448 (e.g., quality and safety of roads; public transportation options), and health and social services  
449 (e.g., counselling and mental health services; emergency response). We used categories of  
450 injuries that are relevant for public health purposes and for designing and implementing  
451 interventions. It may be possible to further split each category. For example, 92% of all  
452 transport injuries in males and 96% in females are from road traffic injuries, with the remainder  
453 being classified as other transport injuries (Extended Data Figure 5). Similarly, suicides can be  
454 classified based on the means of suicide. To the extent that these sub-categories are relevant  
455 for interventions, they can be separately analysed in future studies. Finally, as with any  
456 Bayesian model, choices of prior distributions and hyper-parameters are necessary. There are  
457 alternatives to the priors we used. For example, our weakly informative gamma priors could  
458 have been replaced with penalised complexity priors<sup>51</sup> or uniform priors on the standard

459 deviation scale.<sup>52</sup> We tested a limited number of alternatives and found that our results were  
460 robust to such specifications.

461

462

## References

38. European Centre for Medium-Range Weather Forecasts (ECMWF). ERA5. *Reanalysis datasets* (2019).
39. Shi, L., Kloog, I., Zanobetti, A., Liu, P. & Schwartz, J. D. Impacts of temperature and its variability on mortality in New England. *Nat. Clim. Chang.* **5**, 988–991 (2015).
40. Gasparrini, A. *et al.* Mortality risk attributable to high and low ambient temperature: a multicountry observational study. *Lancet* **386**, 369–375 (2015).
41. Ye, X. *et al.* Ambient temperature and morbidity: a review of epidemiological evidence. *Environ. Health Perspect.* **120**, 19–28 (2012).
42. Basu, R. High ambient temperature and mortality: a review of epidemiologic studies from 2001 to 2008. *Environ. Heal. A Glob. Access Sci. Source* **8**, 40 (2009).
43. Rue, H. & Held, L. *Gaussian Markov random fields. Theory and applications.* Chapman & Hall (2005). doi:10.1007/s00184-007-0162-3
44. Besag, J., York, J. & Mollié, A. Bayesian image restoration, with two applications in spatial statistics. *Ann. Inst. Stat. Math.* (1991). doi:10.1007/BF00116466
45. Knorr-Held, L. Bayesian modelling of inseparable space-time variation in disease risk. in *Statistics in Medicine* (2000). doi:10.1002/1097-0258(20000915/30)19:17/18<2555::AID-SIM587>3.0.CO;2-#
46. Bennett, J. E. *et al.* National and county life expectancy loss from particulate matter pollution in the USA. *PLOS Med.* (2019).
47. Kontis, V. *et al.* Future life expectancy in 35 industrialised countries: projections with a Bayesian model ensemble. *Lancet* (2017). doi:10.1016/S0140-6736(16)32381-9
48. Rue, H., Martino, S. & Chopin, N. Approximate Bayesian inference for latent Gaussian models by using integrated nested Laplace approximations. *J. R. Stat. Soc. Ser. B Stat. Methodol.* (2009). doi:10.1111/j.1467-9868.2008.00700.x
49. IPCC. *IPCC special report on the impacts of global warming of 1.5 °C - Summary for policy makers.* (2018).
50. Bennett, J. E., Blangiardo, M., Fecht, D., Elliott, P. & Ezzati, M. Vulnerability to the mortality effects of warm temperature in the districts of England and Wales. *Nat. Clim. Chang.* **4**, 269–273 (2014).
51. Simpson, D., Rue, H., Riebler, A., Martins, T. G. & Sørbye, S. H. Penalising model component complexity: a principled, practical approach to constructing priors. *Stat. Sci.* (2017). doi:10.1214/16-STS576
52. Gelman, A. Prior distribution for variance parameters in hierarchical models. *Bayesian Anal.* (2006). doi:10.1214/06-BA117A

## **Data availability**

ERA5 temperature data are downloadable from <https://www.ecmwf.int/en/forecasts/datasets/reanalysis-datasets/era5>. Vital statistics files with geographical information can be requested through submission of a proposal to NCHS (<https://www.cdc.gov/nchs/nvss/nvss-restricted-data.htm>).

## **Code availability**

The computer code for the Bayesian model used in this work will be available at [www.globalenvhealth.org/code-data-download](http://www.globalenvhealth.org/code-data-download) upon publication of the paper.

## **Extended Data**

This file contains Extended Data Figure 1, Extended Data Figure 2, Extended Data Figure 3, Extended Data Figure 4, Extended Data Figure 5, Extended Data Table 1, Extended Data Table 2, Extended Data Table 3 and Extended Data Table 4.

## **Acknowledgements**

Robbie Parks is supported by a Wellcome Trust ISSF Studentship. The development of statistical methods is supported by grants from the Wellcome Trust (grant 209376/Z/17/Z). Work on the US mortality data is supported by a grant from US Environmental Protection Agency, as part of the Center for Clean Air Climate Solution (CACES) (assistance agreement number R835873). This paper has not been formally reviewed by EPA. The views expressed in this document are solely those of authors and do not necessarily reflect those of the Agency. EPA does not endorse any products or commercial services mentioned in this publication. We thank Marta Blangiardo, Seth Flaxman and Christopher Paciorek for discussions on statistical

model, and Kavi Bhalla, Howie Frumkin, Andy Haines and Tord Kjellstrom for suggestions of relevant studies.

#### **Author contributions**

All authors contributed to study concept and interpretation of results. RP, GD and ME collated and organised temperature and mortality files. RP, JEB, VK and ME developed statistical model, which was implemented by RP, JEB and VK. RP performed the analysis, with input from other authors. RP and ME wrote the first draft of the paper; other authors contributed to revising and finalizing the paper.

#### **Competing interests statement**

ME reports a charitable grant from AstraZeneca Young Health Programme, and personal fees from Prudential, Scor, and Third Bridge, all outside the submitted work; all other authors declare no competing interests.

537 **Figure 1.** Number of injury deaths, by type of unintentional (transport, falls, drownings, and  
538 other) and intentional (assault and suicide) injury, by sex and age group in the contiguous USA  
539 for 1980-2017.

540 **Figure 2.** National age-standardised death rates from 1980 to 2017, by type of injury, sex and  
541 month.

542 **Figure 3.** Additional annual injury deaths for the 2017 US population in year in which each  
543 month was +1.5°C warmer compared with 1980-2017 average temperatures. The top row  
544 shows breakdown by type of injury, sex and age group. The bottom row shows the break down  
545 by type of injury, sex and month. Black dots represent net changes in deaths for each set of  
546 bars. See Extended Data Figure 3 for results for scenario of 2°C warmer.



547 **Figure 4.** Percent change in death rates in year in which each month was +1.5°C compared  
548 with 1980-2017 average temperatures by type of injury, sex and (A) age group or (B) month.  
549 See Extended Data Figure 4 for scenario of 2°C warmer.