



SUGGESTED FORMULA

Betamethasone Dipropionate 0.05%/ Miconazole 2% and Mupirocin 1% Nipple Ointment

Version number: 1.0

Volume or quantity: 100 gm

Betmethasone Dipropionate, Micronized, USP (BE181)	0.05gm
Miconazole, USP (M1188)	2gm
Mupirocin 2% Ointment (Commerical)	50gm
Aquaphor™	45gm

^{*}The breast should be cleaned with distilled or purified water prior to feeding the infant

SUGGESTED COMPOUNDING PROCEDURES

Weigh all ingredients

- 1. Calculate the required quantity of each ingredient for the total amount to be prepared
- 2. Accurately weigh and/or measure each ingredient
- 3. Levigate micronazole and betamethasone dipropionate powders with a small quantity of the Aquaphor™
- 4. Geometrically, add the remaining Aquaphor™
- 5. Add the mupirocin ointment to the Aquaphor™ mixture by geometric dilution and mix well
- 6. Package and label
- 7. Suggested Quality assessments:
 - a. color
 - b. texture
 - c. container
 - d. Label auxiliary labels, storage, BUD, compounded medication

Store in light resistant air-tight container, at Controlled room temperature

No claims are made as to the safety or efficacy of this preparation. This formulation is provided solely at the unsolicited request of the pharmacist.

Beyond-Use Dates of preparations are conservative estimates from reference books, peer-reviewed literature, and intended duration of therapy, formulation from commercially available products, organoleptic observations and current USP guidelines. Compounders may have stability studies performed by a reputable laboratory if they wish to extend the Beyond-Use Date. It is recommended that you follow USP <795> recommendations for potency testing.

Beyond-Use Date estimated to be 6 months per Compounding Today Formula #907 Precautions should be taken to prevent cross-contamination and exposure of ingredients to the compounder and contamination of the preparation by the compounder. Wear appropriate personal protective equipment. Use safety enclosures (hoods) when weighing and mixing. 5/18 JD