



SUGGESTED FORMULA

Dexamethasone Elixir / Diphenhydramine Elixir / Nystatin Suspension Oral Suspension

Version number: 1.0

Volume: 100mL

Dexamethasone Elixir	20mL
Diphenhydramine 12.5mg/5mL Elixir Oral	100mL
Nystatin 100,000u/mL Oral Suspension	60mL

Diphenhydramine Elixir container alcohol, from 5% to 14% depending upon the product utilized which may cause the preparation to be irritating to open lesions or oral wounds. Bulk powders are recommended for this formula. Dexamethasone elixir: 0.5mg/5mL; Diphenhydramine elixir: 2.5mg/mL and Nystatin 100,000 units/mL. The concentration of components may be altered by the clinician and patient's needs.

SUGGESTED COMPOUNDING PROCEDURES

1. Calculate the required quantity of each ingredient for the total amount to be prepared
2. Accurately weigh and/or measure each ingredient
3. Add the Dexamethasone and Diphenhydramine elixirs to a calibrated container and mix
4. Add the Nystatin Suspension to the elixirs mixture and shake well
5. Package and label as to final concentration of each component
6. Suggested quality assessments:
 - a. Clarity
 - b. Color
 - c. Label (s) - auxiliary labels, storage, BUD

***Do not drink alcoholic beverages with this medicine**

***Shake well**

***May cause drowsiness**

***Keep out of the reach of children**

Store in air-tight container.

Store at room temperature away from excessive heat

No claims are made as to the safety or efficacy of this preparation. This formulation is provided solely at the unsolicited request of the pharmacist.

Beyond-Use Dates of preparations are conservative estimates by the formulator using reference books, peer-reviewed literature, and intended duration of therapy, formulation from commercially available products, organoleptic observations and current USP guidelines. Compounders may have stability studies performed by a reputable laboratory if they wish to extend the Beyond-Use Date.

It is recommended that you follow USP <795> recommendations for potency testing.

Beyond Use Date should be assigned based on the current USP <795> Standards

Precautions should be taken to prevent cross-contamination and exposure of ingredients to the compounder and contamination of the preparation by the compounder. Wear appropriate protective equipment. Use safety enclosures (hoods) when weighing and mixing.

Although much attention has been paid to ensure the accuracy of the formulation contained here, Spectrum Pharmacy Products accepts no liability for the loss or damage arising from reliance on the information. Compounding pharmacists using this formula take full responsibility for the formulations and hold Spectrum Pharmacy Products and Spectrum Chemical Mfg Corp. and its officers, directors and employees harmless for any claim arising from use of or reliance on information contained therein.

10/19 JD