



## SUGGESTED FORMULA

\_\_\_\_\_

Flecainide Acetate 20mg/mL in Cherry Syrup

Version number: 1.0 Volume: 100mL

Flecainide Acetate 2gm

Cherry Syrup (C1223) Q.S. 100mL

\_\_\_\_\_

## **SUGGESTEDCOMPOUNDINGPROCEDURES**

- 1. Check calculations and weigh all ingredients.
- 2. Place Flecainide powder into a mortar
- 3. Add approximately 2mL of cherry syrup and mix to a uniform paste
- 4. Add additional cherry syrup in approximately geometric portions almost to volume, mixing thoroughly after each addition
- 5. Transfer the contents of the mortar to a previously calibrated bottle
- 6. Add sufficient cherry syrup to bring to volume
- 7. Shake well and Label
- 8. Suggested Quality Assessments–follow pharmacy SOPs:
  - a. Weight to Volume calculation
  - b. Color
  - c. Clarity
  - d. Label auxiliary labels, storage, BUD, compounded medication

Store in air tight amber plastic containers refrigerated.

No claims are made as to the safety or efficacy of this preparation. This formulation is provided solely at the unsolicited request of the pharmacist.

Beyond-Use Dates of preparations are conservative estimates by the formulator using reference books, peer-reviewed literature, intended duration of therapy, formulation from commercially available products, organoleptic observations and current USP guidelines. Compounders may have stability studies performed by a reputable laboratory if they wish to extend the Beyond-Use Date. It is recommended that you follow USP <795> recommendations for potency testing.

Beyond-Use Date is estimated to be 60 days when stored Refrigerated per CompoundingToday.com Formula#142

Precautions should be taken to prevent cross-contamination and exposure of ingredients to the compounder and contamination of the preparation by the compounder. Wear appropriate protective equipment. Use safety enclosures (hoods) when weighing and mixing.

9/18 JD