



SUGGESTED FORMULA

Triamcinolone 1mg/mL Mouthwash

Version number: 1.0 Volume: 100mL

Triamcinolone Acetonide, USP 100mg Menthol Crystals, USP 50mg Ethanol, USP 10mL Propylene Glycol, USP 30mL Glycerin, USP 20mL Sodium Saccharin, USP 100mg Sodium Metabisulfite, NF 20mg Disodium EDTA, USP 100mg Purified Water, USP 5mL Liquid flavor of choice (Optional) 1.6mL Sorbitol 70%, USP Q.S 100 mL

SUGGESTED COMPOUNDING PROCEDURES

- 1. Add Ethanol, Propylene Glycol, Glycerin and Purified Water to appropriate beaker with a stir
- 2. Add Menthol Crystals, Sodium Saccharin, Sodium Metabisulfite and Disodium EDTA to beaker and stir to dissolve.
- 3. Add Triamcinolone Acetonide to step 2 and mix until completely dissolved.
- 4. Bring to final volume with Sorbitol 70% and mix until uniform
- 5. Suggested quality assessments:
 - a. Clarity
 - b. Color
 - c. Label (s)- auxiliary labels (external use), storage, BUD

Store in air-tight container.

Store at Controlled Room Temperature

No claims are made as to the safety or efficacy of this preparation. This formulation is provided solely at the unsolicited request of the pharmacist.

Beyond-Use Dates of preparations are conservative estimates by the formulator using reference books, peer-reviewed literature, intended duration of therapy, formulation from commercially available products, organoleptic observations and current USP guidelines. Compounders may have stability studies performed by a reputable laboratory if they wish to extend the Beyond-Use Date.

It is recommended that you follow USP <795> recommendations for potency testing.

Beyond-Use Date is estimated to be <u>180 Days</u> per *Trissel's Stability of Compounded Formulations 5th edition*

Precautions should be taken to prevent cross-contamination and exposure of ingredients to the compounder and contamination of the preparation by the compounder. Wear appropriate protective equipment. Use safety enclosures (hoods) when weighing and mixing.