

# Insert clever, witty title here...

Ian D. Roberts,<sup>★</sup> Laura C. Parker

*Department of Physics and Astronomy, McMaster University, Hamilton ON L8S 4M1, Canada*

Accepted XXX. Received YYY; in original form ZZZ

## ABSTRACT

**Key words:** galaxies: clusters: general – galaxies: evolution – galaxies: groups: – galaxies: statistics

## 1 INTRODUCTION

## 2 DATA

### 2.1 Yang group catalogue

### 2.2 Field catalogue

### 2.3 Group dynamics

### 2.4 Matched data set

To ensure a fair comparison between galaxies in different environments (ie. field galaxies, galaxies in G groups, and galaxies in NG groups) we match our sample of G group galaxies and NG group galaxies by stellar mass, redshift, and halo mass. Additionally, we then match our sample of field galaxies by stellar mass and redshift ensuring that all of our galaxy samples are matched according to important galaxy properties. This is especially important when trying to elucidate information on the effect of group dynamics on galaxy SF and morphological properties for two main reasons:

First, stellar mass, redshift, and halo mass have all been shown to influence galaxy SF and morphology (REF); whereas the impact of group dynamics has been more difficult to pin down (REF) which is perhaps suggestive of a more modest role. Therefore, if one hopes to identify trends in galaxy SF and morphology with group dynamics it is crucial to properly control for these other effects.

Second, standard statistical normality tests, such as the AD test, are inherently biased in identifying non-Gaussian distributions when sample size is large. This is a result of the statistical power of the test increasing with sample size which subsequently allows the detection of more and more subtle departures from normality. These subtle departures from normality may not be physically relevant (in principle, no group is truly Gaussian anyways) and what really matters is whether galaxies in groups which show large departures from normality have different properties than galaxies in groups which show smaller departures from normality.

Since group richness tends scales with halo mass, in the absence of any matching procedure, a sample of NG groups will be biased towards large halo masses compared to a similar sample of G groups – even though many high halo mass NG groups may have been identified on the basis of very small departures from normality. Ensuring that our G and NG samples have very similar halo mass distributions allows us to make a fairer comparison between the two samples.

## 3 RESULTS

### 3.1 Infalling region

### 3.2 Virialized region

## 4 DISCUSSION

## 5 SUMMARY & CONCLUSIONS

## ACKNOWLEDGMENTS

IDR thanks the Ontario Graduate Scholarship program for funding. LCP thanks the National Science and Engineering Research Council of Canada for funding. The authors thank F. Evans for matching together the various SDSS catalogues used in this research. We thank X. Yang et al. for making their SDSS DR7 group catalogue publicly available, L. Simard et al. for the publication of their SDSS DR7 morphology catalogue, J. Brinchmann et al. for publication of their SDSS SFRs, and the NYU-VAGC team for the publication of their SDSS DR7 catalogue. This research would not have been possible without access to these public catalogues.

Funding for the SDSS has been provided by the Alfred P. Sloan Foundation, the Participating Institutions, the National Science Foundation, the U.S. Department of Energy, the National Aeronautics and Space Administration, the Japanese Monbukagakusho, the Max Planck Society, and the Higher Education Funding Council for England. The SDSS Web Site is <http://www.sdss.org/>.

The SDSS is managed by the Astrophysical Research Consortium for the Participating Institutions. The Participating Institutions are the American Museum of Natu-

<sup>★</sup> E-mail: [roberid@mcmaster.ca](mailto:roberid@mcmaster.ca)




Figure 1.




Figure 2.

This paper has been typeset from a  $\text{\LaTeX}$  file prepared by the author.

ral History, Astrophysical Institute Potsdam, University of Basel, University of Cambridge, Case Western Reserve University, University of Chicago, Drexel University, Fermilab, the Institute for Advanced Study, the Japan Participation Group, Johns Hopkins University, the Joint Institute for Nuclear Astrophysics, the Kavli Institute for Particle Astrophysics and Cosmology, the Korean Scientist Group, the Chinese Academy of Sciences (LAMOST), Los Alamos National Laboratory, the Max-Planck-Institute for Astronomy (MPIA), the Max-Planck-Institute for Astrophysics (MPA), New Mexico State University, Ohio State University, University of Pittsburgh, University of Portsmouth, Princeton University, the United States Naval Observatory, and the University of Washington.




Figure 3.




Figure 4.




Figure 5.