

Hochschule RheinMain - University of Applied Sciences

University of Applied Sciences
Department of
Design, Computer Science, Media
Prof. Dr. D. Richter

Transilvania University Brasov
Faculty of
Mathematics and Informatics
Dr. I. Plajer

ISIP 2019
International School on Image Processing
July 15th - 26th, 2019




detlef.richter@HS-RM.de

Prof. Dr. D. Richter Department [Design>Computer Science>Media] 10

Hochschule RheinMain - University of Applied Sciences

University of Applied Sciences
Department of
Design, Computer Science, Media
Prof. Dr. D. Richter

Lectures in Image Processing

Chapter 7

Hough Transform

(Part 1 : HT for Lines)

Prof. Dr. D. Richter Department [Design>Computer Science>Media] 10

Hochschule RheinMain - University of Applied Sciences

Notice to Copyright

All information in this paper and part thereof
is only for private use
in connection with the lecture
of Prof. Dr. D. Richter
from Hochschule RheinMain - University of Applied Sciences

Reproduction or publication and part thereof
is not allowed
without written permission of the author.

Prof. Dr. D. Richter Department [Design>Computer Science>Media] 10

Hochschule RheinMain - University of Applied Sciences

7. Hough Transform

- Graphical data processing : Data → Image
- Digital image manipulation : Image → Image
- Digital image processing : Image → Data

Prof. Dr. D. Richter Department [Design>Computer Science>Media] 10

Hochschule RheinMain - University of Applied Sciences



7.0 Hough Transform

- 7.1 Hough-Transform for Lines (2 parameters)
- 7.2 Hough-Transform for Circles (3 parameters)
- 7.3 Hough-Transform for Parabolas (4 parameters)
- 7.4 Analysis of Hough matrices
- 7.5 Hough-Transform for Lines, Sophisticated Applications

Prof. Dr. D. Richter Department [Design>Computer Science>Media] 10

Hochschule RheinMain - University of Applied Sciences

7.0 Hough Transform, Goal of Application (1)

The following equations are to be found:

line 1: $y = m_0 \cdot x + n_0$ or $x \cdot \cos \alpha_0 + y \cdot \sin \alpha_0 = r_0$
 lines 2 to 4 as well with afterwards known m_0, n_0 or α_0 and r_0 .
 circle 1: $(x - x_m)^2 + (y - y_m)^2 = r^2$
 circle 2 as well with afterwards known x_m, y_m , and r .

Prof. Dr. D. Richter Department [Design>Computer Science>Media] 10

7.0 Hough Transform, Goal of Application (2)



Equations of analytical defined shapes.

- orientation of the parabola (1 parameter),
- the focal length $f = p/2$ or width $w = 2p$ (1 parameter),
- vertex (2 parameters).

Use a-priori knowledge to reduce the number of parameters

Prof. Dr. D. Richter

Department [Design>Computer Science>Media]

10

7.0 Hough Transform, Goal of Application (2)



Focal length $f = p/2$ or width $w = 2p$ (1 parameter):

$$y^2 = 2 p x$$

Vertex (2 parameters):

$$(y - y_0)^2 = 2 p (x - x_0)$$

Orientation of the parabola α (1 parameter):


$$\begin{pmatrix} y' \\ x' \end{pmatrix} = R(\alpha) * \begin{pmatrix} y \\ x \end{pmatrix}$$

Prof. Dr. D. Richter

Department [Design>Computer Science>Media]

10

7.1 Hough Transform for lines



Equation of straight line :

$$y = m_0 * x + n_0$$

x, y variable; m_0, n_0 const.

m_0 : slope

n_0 : interception of y-axis

Consider point P_1 :


$$y_1 = m_0 * x_1 + n_0$$

Prof. Dr. D. Richter

Department [Design>Computer Science>Media]

10

7.1 Hough Transform for lines



The values of n depend on the slopes m for all lines passing point P_1 of a straight line g_1 :

$$g_1 : n = -x_1 * m + y_1,$$

n, m : variable; x_1, y_1 : const.

$-x_1$: slope;
 y_1 : intersection of n -axis


Line g_1 in the m, n -diagram represents point P_1 of x, y -diagram

Prof. Dr. D. Richter

Department [Design>Computer Science>Media]

10

7.1 Hough Transform for lines




Only one white point P_1 is known. Calculate all potential lines passing P_1 . Register all pairs (m, n) in a matrix.

Prof. Dr. D. Richter

Department [Design>Computer Science>Media]

10

7.1 Hough Transform for lines



Consider point P_2 :

$$y_2 = m_0 * x_2 + n_0$$

Representation of point P_2 within n, m -diagram :

$$g_2 : n = -x_2 * m + y_2,$$


n, m : variable; x_2, y_2 : const.

Prof. Dr. D. Richter

Department [Design>Computer Science>Media]

10

7.1 Hough Transform for lines




The intersecting point (m_0, n_0) of the straight lines g_1 and g_2 represents that line within the x, y -diagram, which passes the points P_1 and P_2 as well.

Same consideration holds for all points P_i on line $y = m_0 * x + n_0$
→ straight lines in n, m -diagram passing the point m_0, n_0

Prof. Dr. D. Richter

Department [Design>Computer Science>Media] 10

7.1 Hough Transform for lines



Define m, n -matrix

Find white pixels $P(x_i, y_i)$ in the image assumed to lie on lines.

Calculate all values $n, n = -x_i * m + y_i$, for given x_i, y_i with independent variable m [m : integer].

Increment according matrix elements m, n for all combinations m, n of possible lines passing point $P_i(x_i, y_i)$.
 m, n : integer values

Prof. Dr. D. Richter

Department [Design>Computer Science>Media] 10

7.1 Hough Transform for lines

What happens with lines parallel to image columns?

In this case : $m = \pm \infty$ und $n = \pm \infty$


∞ : not computable in computers.

Look for another analytical equation for straight lines in which infinite values don't occur.

Prof. Dr. D. Richter

Department [Design>Computer Science>Media] 10


7.1 Hough Transform for lines, Hesse Equation



Prof. Dr. D. Richter

Department [Design>Computer Science>Media] 10

7.1.1 Hough Transform for lines, Hesse Equation



Hesse equation of straight lines :

$$g_1 : x * \cos \alpha_0 + y * \sin \alpha_0 = r_0$$

r_0 : perpendicular from origin to g_1

α_0 : angle between x -axis and perpendicular (math. counterclockwise)


Consider pixel $P_1(x_1, y_1)$ lying upon g_1 :

$$x_1 * \cos \alpha_0 + y_1 * \sin \alpha_0 = r_0$$

Prof. Dr. D. Richter

Department [Design>Computer Science>Media] 10

7.1.1 Hough Transform for lines, Hesse Equation



Consider 512^2 -image.

Consider $x_1 * \cos \alpha + y_1 * \sin \alpha = r$ with α and r as variables







All possible lines passing P_1 are calculated within a range of $-90^\circ \leq \alpha < +90^\circ$

Limiting α to an interval of $\pm 90^\circ$ results in positive and negative radii.

Maximal radius : $\pm 512 * \sqrt{2}$

Prof. Dr. D. Richter

Department [Design>Computer Science>Media] 10



7.1.2 Hough Transform, Definition of Matrices

- Define Hough-Matrix for two quadrants (e.g. angles α : $-90^\circ \leq \alpha < +90^\circ$ or $0^\circ \leq \alpha < 180^\circ$) and allow positive and negative radii.
- Define Hough-Matrix for three quadrants (in this case for angles α : $-90^\circ \leq \alpha < +180^\circ$) and exclude negative radii.
- Define Hough-Matrix for four quadrants and allow positive radii with half of maximum size.

7.1.2 Hough Transform, Definition of Matrices

Memory space needed:

- 2 quadrants : $180 * 960 * 2 = 345\ 600$
- 3 quadrants : $270 * 960 * 1 = 259\ 200$
- 4 quadrants : $360 * 960 * \frac{1}{2} = 172\ 800$


Assumed to have an angle resolution of 1° and radius resolution of 1 Pixel.

7.1.2 Hough Transform, Definition of Matrices


- Resolution of r and α has to be defined according to the specified problem (e.g. α in steps of 1° , 2° , 3° or 5° , radius resolution might be smaller than, equal or larger than one pixel).
- The better the resolution, the more time consuming the calculation.
- Or : Who wants to see details, needs more time to look at.

7.1.5 Hough Transform, Application


Hough matrix of two quadrants, positive and negative radii, absolute and logarithmic presentation.
How could the original image look like?



7.1.5 Hough Transform, Application




7.1.5 Hough Transform, Application



- Chess board like calibrating pattern (see chapter 6 : Radial symmetric distortion by lenses)
- High-pass filtered image

7.1.5 Hough Transform, Application




Hough matrix with 4 quadrants
(no results for quadrant 3)

- Representation of Hough-Matrix
- 10 vertical equidistant lines (plus 1 line at image boundary with larger distance) ($\alpha = 0^\circ$ and 360°)
- 7 equidistant horizontal lines ($\alpha = 90^\circ$)
- Define r, α
- Calculate corresponding intersection points

Prof. Dr. D. Richter

Department [Design>Computer Science>Media] 10


7.1.5 Hough Transform, Application



Prof. Dr. D. Richter

Department [Design>Computer Science>Media] 10


7.1.5 Hough Transform, Application



Prof. Dr. D. Richter 270° resp. 90°

Department [Design>Computer Science>Media] 10

7.1.5 Hough Transform, Application



Prof. Dr. D. Richter 270° resp. 90°

Department [Design>Computer Science>Media] 10