



# High Performance Computing with C++ Final Report

ROBERT SCHÜTZE

4140641

[schuetze.r@web.de](mailto:schuetze.r@web.de)

June 9, 2021

# Contents

### Derivation of LJ-Force

Suppose we have a list of  $N$  atoms with the position vectors  $\vec{r}_k := [x_k, y_k, z_k]^T$ . The vector between a pair of atoms is defined as  $\vec{r}_{ij} := \vec{r}_j - \vec{r}_i$ . The pair distance  $r_{ij}$  is defined as the 2-norm of  $\vec{r}_{ij}$ :

$r_{ij} := \sqrt{(x_j - x_i)^2 + (y_j - y_i)^2 + (z_j - z_i)^2}$ . The potential energy of an atom pair is given by (cite ??):

$$E_{pot_{ij}} = 4\epsilon \left[ \left( \frac{\sigma}{r_{ij}} \right)^{12} - \left( \frac{\sigma}{r_{ij}} \right)^6 \right] \quad (1)$$

The parameters  $\epsilon$  and  $\sigma$  tune the potential strength and the resting distance respectively. To get the potential energy of the whole system, we have to sum over all atom pairs:

$$E_{pot} = \frac{1}{2} \sum_{ij, i \neq j} E_{pot_{ij}} \quad (2)$$

The force, which the  $i$ -th atom exerts on the  $k$ -th atom is the pair force:

$$\vec{f}_{ik} = \nabla_{\vec{r}_k} E_{pot_{ik}} = \frac{\partial E_{pot_{ik}}}{\partial r_{ik}} \cdot \nabla_{\vec{r}_k} r_{ik} \quad (3)$$

Using Eq. (1), the first part of Eq. (3) reads:

$$\frac{\partial E_{pot_{ik}}}{\partial r_{ik}} = \frac{24\epsilon(\sigma^6 r_{ik}^6 - 2\sigma^{12})}{r_{ik}^{13}} \quad (4)$$

The first line of the gradient of the pair distance can be calculated as follows:

$$\frac{dr_{ik}}{dx_k} = \frac{1}{2r_{ik}} \cdot 2(x_k - x_i) \cdot (1 - \delta_{ik}) \quad (5)$$

Applying the same principle from Eq. (5) to the other two coordinate axes and plugging it, together with the result from Eq. (4), into Eq. (3) yields:

$$\vec{f}_{ik} = \frac{24\epsilon(\sigma^6 r_{ik}^6 - 2\sigma^{12})}{r_{ik}^{14}} \cdot (1 - \delta_{ik}) \cdot \vec{r}_{ik} \quad (6)$$