

MATH 447 - Homework 5

Robert D. French

September 30, 2014

SECTION 3.1

PROBLEM 10

Prove that if $\lim(x_n) = 0$ and if $x > 0$, then there exists a natural number M such that $x_n > 0$ for all $n \geq M$.

PROOF. By the definition of limit, for each positive ϵ such an $M(\epsilon)$ can be found. Thus, take ϵ to be smaller than x , so that $x_n \in V_\epsilon(x) = (x - \epsilon, x + \epsilon)$ whenever $n \geq M$. Since $\epsilon < x$, $x - \epsilon \in \mathbb{P}$ so $x_n > 0$. ■

PROBLEM 14

Let $b \in \mathbb{R}$ satisfy $0 < b < 1$. Show that $\lim(nb^n) = 0$.

PROOF. We know that $n^{1/n} > 1 \forall n \in \mathbb{N}$. This implies that $n^{1/n} = 1 + k_n$ for some $k_n > 0$, so that we have $bn^{1/n} = b + bk_n$. Thus, $nb^n = (b + bk_n)^n$.

By the Binomial Theorem we know that

$$nb^n = (b + bk_n)^n = b^n + nb^{n-1}bk_n + \frac{1}{2}n(n-1)b^{n-2}b^2k_n^2 + \dots$$

which tells us that $nb^n \geq b^n + \frac{1}{2}n(n-1)b^n k_n^2$. Factoring out b^n and subtracting 1 yields

$$(n-1) \geq \frac{1}{2}n(n-1)k_n^2$$

so that we have $\frac{2}{n} \geq k_n^2$.

PROBLEM 17

HINT

If $n \geq 3$, then $0 < 2^n/n! \leq 2(\frac{2}{3})^{n-2}$.

PROOF. Let $n = 3$, then $\frac{2^3}{3!} = \frac{8}{6} \leq 2(\frac{2}{3}) = \frac{4}{3} = \frac{8}{6}$. Thus, we have established a basis for induction.

Now assume that $0 < \frac{2^{n-1}}{(n-1)!} \leq 2(\frac{2}{3})^{n-3}$. Multiplying by 2 yields $0 < \frac{2^n}{(n-1)!} \leq 2(\frac{2^{n-2}}{3^{n-3}})$. Since $n \geq 3$, we have $0 < \frac{1}{n} \leq \frac{1}{3}$ so that

$$0 < \frac{2^n}{(n-1)!} \cdot \frac{1}{n} = \frac{2^n}{n!} \leq 2 \cdot \left(\frac{2^{n-2}}{3^{n-3}}\right) \cdot \frac{1}{3} = 2\left(\frac{2}{3}\right)^{n-2}$$

whence $0 < \frac{2^n}{n!} \leq 2\left(\frac{2}{3}\right)^{n-2}$. Therefore the hypothesis holds by the Principal of Mathematical Induction. ■

MAIN PROBLEM

Show that $\lim(2^n/n!) = 0$.

PROOF. We know that $0 < \frac{2^n}{n!} \leq 2\left(\frac{2}{3}\right)^{n-2}$ when $n \geq 3$, so if we can show that $2\left(\frac{2}{3}\right)^{n-2} \rightarrow 0$, then $\frac{2^n}{n!} \rightarrow 0$ by the Squeeze Theorem.

Take $b = 2/3$ and $n = m - 2$, then by Exercise 3.1.14 we know $nb^n \rightarrow 0$. Since $2 \leq m - 2 \forall m > 3$, we have that $0 < 2\left(\frac{2}{3}\right)^{m-2} \leq (m-2)b^{m-2}$ so that $2\left(\frac{2}{3}\right)^{m-2} \rightarrow 0$. Thus, by the Squeeze Theorem, $2^n/n! \rightarrow 0$. ■

PROBLEM 18

If $\lim(x_n) = x > 0$, show that there exists a natural number K such that if $n \geq K$, then $\frac{1}{2}x < x_n < 2x$.

PROOF. Pick $\epsilon < \frac{1}{2}x$, then by the definition of limit $\exists K$ such that $x_n \in V_\epsilon(x) \forall n \geq K$. Since $V_\epsilon(x) = (x - \epsilon, x + \epsilon) = (x/2, 3x/2)$, we know $V_\epsilon(x) \subseteq (x/2, 2x)$, demonstrating that such a K exists. ■

SECTION 3.2

PROBLEM 7

If (b_n) is a bounded sequence and $\lim(a_n) = 0$, show that $\lim(a_nb_n) = 0$. Explain why Thm 3.2.3 cannot be used.

REMARK. We may not apply Theorem 3.2.3 directly because it requires that (b_n) is convergent, but we know only that (b_n) is bounded. However, we may employ Theorem 3.2.3 for certain features of our argument.

PROOF. Since (b_n) is bounded, we have

$$I_b = \inf(b_n) \leq b_j \leq \sup(b_n) = S_b \quad \forall j \in \mathbb{N}$$

which implies

$$I_b |a_j| = \inf(b_n) |a_j| \leq b_j \cdot |a_j| \leq \sup(b_n) \cdot |a_j| = S_b \cdot |a_j| \quad \forall j \in \mathbb{N}$$

Thus if $\inf(b_n) |a_j| \rightarrow 0$ and $\sup(b_n) |a_j| \rightarrow 0$, we will be confident that $b_j |a_j| \rightarrow 0$.

Since $\inf(b_n)$ and $\sup(b_n)$ are constants, and since we know $|a_j| \rightarrow 0$ due to the fact that $a_j \rightarrow 0$, we have that $\inf(b_n) |a_j| \rightarrow 0$ and $\sup(b_n) |a_j| \rightarrow 0$ by Theorem 3.2.3. Thus, $b_j \cdot |a_j| \rightarrow 0$.

Therefore, given $\epsilon > 0$, we know $\exists K(\epsilon)$ such that $|b_j \cdot |a_j| - 0| < \epsilon \quad \forall j \geq K(\epsilon)$.

What we want to show is that $|b_j \cdot a_j - 0| < \epsilon \quad \forall j \geq M(\epsilon)$. By the above work, we know $|b_j \cdot |a_j| - 0| < \epsilon$ which tells us that $|j \cdot j - 0| < \epsilon$ and therefore $\lim(a_n b_n) = 0$. ■

PROBLEM 9

Let $y_n := \sqrt{n+1} - \sqrt{n}$ for $n \in \mathbb{N}$. Show that $(\sqrt{n} y_n)$ converges. Find the limit.

PROOF.

PROBLEM 11

PART A

Find $\lim \left((3\sqrt{n})^{1/2n} \right)$

PROOF.

PART B

Find $(\sqrt{n^2 + 5n} - n)$

PROOF.

PROBLEM 12

If $0 < a < b$, determine $\left(\frac{a^{n+1} + b^{n+1}}{a^n + b^n} \right)$.

PROOF.

PROBLEM 13

If $a > 0$, $b > 0$, show that $\lim(\sqrt{(n+a)(n+b)} - n) = (a+b)/2$.

PROOF.

PROBLEM 14

PART A

Use the Squeeze Theorem to find $\lim(n^{1/n^2})$.

PROOF.

PART B

Use the Squeeze Theorem to find $\lim((n!)^{1/n^2})$.

PROOF.

PROBLEM 15

Show that if $z_n := (a^n + b^n)^{1/n}$ where $0 < a < b$, then $\lim(z_n) = b$.

PROOF.

PROBLEM 23

Show that if (x_n) and (y_n) are convergent sequences, then the sequences (u_n) and (v_n) defined by $u_n := \max\{x_n, y_n\}$ and $v_n := \min\{x_n, y_n\}$ are also convergent.

PROOF.

PROBLEM 24

Show that if (x_n) , (y_n) , (z_n) are convergent sequences, then the sequence (w_n) defined by $w_n := \text{mid}\{x_n, y_n, z_n\}$ is also convergent.

PROOF.