

Combinatorial structures with lots of symmetry

Robert Gray




University
of
St Andrews

Lisbon, October 2009


Graphs and symmetry

Definition

A graph Γ is a structure $(V\Gamma, \sim)$ where $V\Gamma$ is a set, and \sim is a symmetric irreflexive binary relation on $V\Gamma$.



Γ_1




Γ_2


Graphs and symmetry

Definition

A graph Γ is a structure $(V\Gamma, \sim)$ where $V\Gamma$ is a set, and \sim is a symmetric irreflexive binary relation on $V\Gamma$.



Γ_1




Γ_2

- Γ_1 has “more symmetry” than Γ_2 .


Graphs and symmetry

Definition

A graph Γ is a structure $(V\Gamma, \sim)$ where $V\Gamma$ is a set, and \sim is a symmetric irreflexive binary relation on $V\Gamma$.



Γ_1



Γ_2

- ▶ Γ_1 has “more symmetry” than Γ_2 .
- ▶ Imagine you are trapped inside the graph:
 - ▶ In Γ_1 the world looks the same from every vertex.
 - ▶ In Γ_2 the world looks different from each vertex.

Automorphisms

- ▶ An **isomorphism** $\phi : \Gamma_1 \rightarrow \Gamma_2$ of graphs is a bijection such that:

$$v \sim w \Leftrightarrow v\phi \sim w\phi \quad (\forall v, w \in \Gamma_1).$$

- ▶ An **automorphism** of a graph Γ is an isomorphism $\phi : \Gamma \rightarrow \Gamma$.
- ▶ **Aut(Γ)** - the full automorphism group of the graph Γ .

Automorphisms


- ▶ An **isomorphism** $\phi : \Gamma_1 \rightarrow \Gamma_2$ of graphs is a bijection such that:

$$v \sim w \Leftrightarrow v\phi \sim w\phi \quad (\forall v, w \in \Gamma_1).$$


- ▶ An **automorphism** of a graph Γ is an isomorphism $\phi : \Gamma \rightarrow \Gamma$.
- ▶ **Aut(Γ)** - the full automorphism group of the graph Γ .

Roughly speaking... the more symmetry a graph has the larger its automorphism group will be, and vice versa.

Vertex transitivity



Γ_1




Γ_2


Definition

A graph Γ is **vertex transitive** if $\text{Aut}(\Gamma)$ acts transitively on $V\Gamma$.

Vertex transitivity



Γ_1




Γ_2

Definition


A graph Γ is **vertex transitive** if $\text{Aut}(\Gamma)$ acts transitively on $V\Gamma$.

- ▶ Γ_1 is vertex transitive
 - ▶ $\text{Aut}(\Gamma) = \text{rotations} + \text{reflections}$ (Dihedral group)

Vertex transitivity



Γ_1



Γ_2

Definition

A graph Γ is **vertex transitive** if $\text{Aut}(\Gamma)$ acts transitively on $V\Gamma$.

- ▶ Γ_1 is vertex transitive
 - ▶ $\text{Aut}(\Gamma) = \text{rotations} + \text{reflections}$ (Dihedral group)
- ▶ Γ_2 is not vertex transitive
 - ▶ Even worse, $|\text{Aut}(\Gamma_2)| = 1$

Vertex transitive graphs

Question. Vertex transitive graphs are “nice”. Is there any chance we could classify them (i.e. explicitly describe them all)?

Vertex transitive graphs

Question. Vertex transitive graphs are “nice”. Is there any chance we could classify them (i.e. explicitly describe them all)?

Answer. No.

There are simply “too many” vertex transitive graphs for us to stand a chance of describing them all.

For example:

- ▶ G - group, A - generating set for G with $A = A^{-1}$
- ▶ the **Cayley graph** $\Gamma(G, A)$:
 - ▶ $V\Gamma(G, A) = G$
 - ▶ $x \sim y \Leftrightarrow xy^{-1} \in A$,

is vertex transitive.

(although not every vertex transitive graph is the Cayley graph of a group)

Symmetry properties for graphs

Examples

- ▶ Various symmetry properties have received attention:
 - ▶ vertex transitive
 - ▶ arc-transitive, k -arc-transitive (Tutte (1947))
 - ▶ distance-transitive (Biggs and Smith (1971))
 - ▶ homogeneous, k -homogeneous (Fraïssé (1953))

Symmetry properties for graphs

Examples

- ▶ Various symmetry properties have received attention:
 - ▶ vertex transitive
 - ▶ arc-transitive, k -arc-transitive (Tutte (1947))
 - ▶ distance-transitive (Biggs and Smith (1971))
 - ▶ homogeneous, k -homogeneous (Fraïssé (1953))

General problem

For a given symmetry property \mathcal{P} , classify those graphs Γ satisfying property \mathcal{P} .

Homogeneous graphs

Definition






A graph Γ is **homogeneous** if every isomorphism between finite induced subgraphs of Γ can be extended to an automorphism of Γ .

Homogeneous graphs

Definition

A graph Γ is **homogeneous** if every isomorphism between finite induced subgraphs of Γ can be extended to an automorphism of Γ .

Induced subgraphs.

Γ	Some induced subgraphs	
		
		


Homogeneous graphs

Definition


A graph Γ is **homogeneous** if every isomorphism between finite induced subgraphs of Γ can be extended to an automorphism of Γ .

Induced subgraphs.


However, the graph Γ :



has no induced subgraph isomorphic to:



Extending isomorphisms to automorphisms




The isomorphism

$$(v_1, v_3) \mapsto (v_4, v_2)$$

between finite induced subgraphs

Extending isomorphisms to automorphisms




The isomorphism

$$(v_1, v_3) \mapsto (v_4, v_2)$$

between finite induced subgraphs extends to the automorphism

$$\begin{pmatrix} v_1 & v_2 & v_3 & v_4 & v_5 \\ v_4 & v_3 & v_2 & v_1 & v_5 \end{pmatrix}$$

Extending isomorphisms to automorphisms



The isomorphism


$$(v_1, v_3) \mapsto (v_4, v_2)$$

between finite induced subgraphs extends to the automorphism

$$\begin{pmatrix} v_1 & v_2 & v_3 & v_4 & v_5 \\ v_4 & v_3 & v_2 & v_1 & v_5 \end{pmatrix}$$

Fact. The pentagon is a homogeneous graph.

Extending isomorphisms to automorphisms




The isomorphism between finite induced subgraphs

$$(v_1, v_3) \mapsto (v_1, v_4)$$

does not extend to the automorphism.

e.g. There is a path of length 2 from v_1 to v_3 , while there is no path of length 2 from v_1 to v_4 .

Extending isomorphisms to automorphisms



The isomorphism between finite induced subgraphs

$$(v_1, v_3) \mapsto (v_1, v_4)$$

does not extend to the automorphism.

e.g. There is a path of length 2 from v_1 to v_3 , while there is no path of length 2 from v_1 to v_4 .

So the hexagon is not a homogeneous graph.


Classification of finite homogeneous graphs

Gardiner classified the finite homogeneous graphs.

Theorem (Gardiner (1976))

A finite graph is homogeneous if and only if it is isomorphic to one of the following:

1. *finitely many disjoint copies of a **complete graph** K_r (or its complement, **complete multipartite graph**)*
2. *the **pentagon** C_5*
3. *the graph $K_3 \times K_3$ drawn below*



The random graph

Definition (The random graph R)

Constructed by [Rado \(1964\)](#). The vertex set is the natural numbers (including zero).

For $i, j \in \mathbb{N}$, $i < j$, then i and j are joined if and only if the i th digit in j in base 2, reading right-to-left, is 1.

Example


Since $88 = 8 + 16 + 64 = 2^3 + 2^4 + 2^6$ the numbers less than 88 that are adjacent to 88 are just $\{3, 4, 6\}$.

Of course, many numbers greater than 88 will also be adjacent to 88 (for example 2^{88}).

The random graph

Consider the following property of graphs:

(*) For any two finite disjoint sets U and V of vertices, there exists a vertex w adjacent to **every vertex in U** and to **no vertex in V** .



The random graph

Consider the following property of graphs:

(*) For any two finite disjoint sets U and V of vertices, there exists a vertex w adjacent to **every vertex in U** and to **no vertex in V** .

Theorem

There exists a countably infinite graph R satisfying property (), and it is unique up to isomorphism. The graph R is homogeneous.*

Existence. The random graph R defined above satisfies property (*).

The random graph

Consider the following property of graphs:

(*) For any two finite disjoint sets U and V of vertices, there exists a vertex w adjacent to **every vertex in U** and to **no vertex in V** .

Theorem

There exists a countably infinite graph R satisfying property (), and it is unique up to isomorphism. The graph R is homogeneous.*

Existence. The random graph R defined above satisfies property (*).

Uniqueness and homogeneity. Both follow from a **back-and-forth** argument. Property (*) is used to extend the domain (or range) of any isomorphism between finite substructures one vertex at a time.

Properties of the random graph

- ▶ **Universal**: it embeds every countable graph as an induced subgraph.
 - ▶ R is the unique countably infinite universal homogeneous graph.

Properties of the random graph

- ▶ **Universal**: it embeds every countable graph as an induced subgraph.
 - ▶ R is the unique countably infinite universal homogeneous graph.
- ▶ **Partition property**: for any partition $V\Gamma = X \cup Y$ either the subgraph induced by X , or the subgraph induced by Y , is again isomorphic to R .
 - ▶ Aside from the complete graph, and empty graph, R is the only countable graph with this property.

Properties of the random graph

- ▶ **Universal**: it embeds every countable graph as an induced subgraph.
 - ▶ R is the unique countably infinite universal homogeneous graph.
- ▶ **Partition property**: for any partition $V\Gamma = X \cup Y$ either the subgraph induced by X , or the subgraph induced by Y , is again isomorphic to R .
 - ▶ Aside from the complete graph, and empty graph, R is the only countable graph with this property.
- ▶ **Random**: if we choose a countable graph at random (edges independently with probability $\frac{1}{2}$), then with probability 1 it is isomorphic to R (Erdős and Rényi, 1963).

Properties of the random graph

- ▶ **Universal**: it embeds every countable graph as an induced subgraph.
 - ▶ R is the unique countably infinite universal homogeneous graph.
- ▶ **Partition property**: for any partition $V\Gamma = X \cup Y$ either the subgraph induced by X , or the subgraph induced by Y , is again isomorphic to R .
 - ▶ Aside from the complete graph, and empty graph, R is the only countable graph with this property.
- ▶ **Random**: if we choose a countable graph at random (edges independently with probability $\frac{1}{2}$), then with probability 1 it is isomorphic to R (Erdős and Rényi, 1963).
- ▶ $\text{Aut}(R)$ is an infinite simple group (Truss, 1985).

Homogeneous relational structures

Definition

A relational structure M is **homogeneous** if every isomorphism between finite substructures of M can be extended to an automorphism of M .

Homogeneous relational structures

Definition

A relational structure M is **homogeneous** if every isomorphism between finite substructures of M can be extended to an automorphism of M .

Relational structures

- ▶ a **relational structure** consists of a set A , and some relations R_1, \dots, R_m (can be unary, binary, ternary, ...)
- ▶ an (induced) **substructure** is obtained by taking a subset $B \subseteq A$ and keeping only those relations where all entries in the tuple belong to B
- ▶ an **isomorphism** is a “structure preserving” mapping (i.e. a bijection ϕ such that ϕ and ϕ^{-1} are both homomorphisms)

Examples of homogeneous structures

The countable **random graph** R

X - a pure set

- ▶ automorphism group is the full symmetric group where any partial permutation can be extended to a (full) permutation

(\mathbb{Q}, \leq) - the rationals with their usual ordering

- ▶ the automorphisms are the order-preserving permutations
- ▶ isomorphisms between finite substructures can be extended to automorphisms that are piecewise-linear

Connection with model theory

Common theme in **model theory**:

translation between “model theoretic terminology” and “permutation group theoretic terminology”

Connection with model theory

Common theme in **model theory**:

translation between “model theoretic terminology” and “permutation group theoretic terminology”

Example.

- (I) A structure M is **\aleph_0 -categorical** if all countable models of the first-order theory of M are isomorphic to M .
- (II) A permutation group on an infinite set Ω is called **oligomorphic**, if it has finitely many orbits of n -tuples, for all $n \geq 1$.

Connection with model theory

Common theme in **model theory**:

translation between “model theoretic terminology” and “permutation group theoretic terminology”

Example.

- (I) A structure M is **\aleph_0 -categorical** if all countable models of the first-order theory of M are isomorphic to M .
- (II) A permutation group on an infinite set Ω is called **oligomorphic**, if it has finitely many orbits of n -tuples, for all $n \geq 1$.

Theorem (Ryll-Nardzewski)

A countable structure M over a first-order language is \aleph_0 -categorical if and only if $\text{Aut}(M)$ is oligomorphic.

Connection with model theory

Common theme in **model theory**:

translation between “model theoretic terminology” and “permutation group theoretic terminology”

Example.

- (I) A structure M is **\aleph_0 -categorical** if all countable models of the first-order theory of M are isomorphic to M .
- (II) A permutation group on an infinite set Ω is called **oligomorphic**, if it has finitely many orbits of n -tuples, for all $n \geq 1$.

Theorem (Ryll-Nardzewski)

A countable structure M over a first-order language is \aleph_0 -categorical if and only if $\text{Aut}(M)$ is oligomorphic.

Homogeneous structures provide a rich source of \aleph_0 -categorical structures.

Homogeneous structures and Fraïssé's theorem

The notion of homogeneous structure goes back to the fundamental work of Fraïssé (1953).

- ▶ The **age** of a relational structure M is the class of isomorphism types of its finite substructures.

Fraïssé proved a theorem which gives a necessary and sufficient condition on a class \mathcal{C} of finite structures for it to be the age of a countable homogeneous structure M .

Homogeneous structures and Fraïssé's theorem

The notion of homogeneous structure goes back to the fundamental work of Fraïssé (1953).

- ▶ The **age** of a relational structure M is the class of isomorphism types of its finite substructures.

Fraïssé proved a theorem which gives a necessary and sufficient condition on a class \mathcal{C} of finite structures for it to be the age of a countable homogeneous structure M .

- ▶ The key condition is the **amalgamation property**:
 - ▶ two structures in \mathcal{C} with isomorphic substructures can be “glued together” so that the substructures are identified, inside a larger structure in \mathcal{C} .

If Fraïssé's conditions hold, then M is unique, \mathcal{C} is called a **Fraïssé class**, and M is called the **Fraïssé limit** of the class \mathcal{C} .

Countable homogeneous graphs

Examples

- ▶ The class of all finite graphs is a Fraïssé class. Its Fraïssé limit is the random graph R .

Countable homogeneous graphs

Examples

- ▶ The class of all finite graphs is a Fraïssé class. Its Fraïssé limit is the random graph R .
- ▶ The class of all finite graphs not embedding the complete graph K_n (for some fixed n) is a Fraïssé class. We call the Fraïssé limit the **countable generic K_n -free graph**.

Countable homogeneous graphs

Examples

- ▶ The class of all finite graphs is a Fraïssé class. Its Fraïssé limit is the random graph R .
- ▶ The class of all finite graphs not embedding the complete graph K_n (for some fixed n) is a Fraïssé class. We call the Fraïssé limit the **countable generic K_n -free graph**.

Theorem (Lachlan and Woodrow (1980))

*Let Γ be a countably infinite homogeneous graph. Then Γ is isomorphic to one of: the **random graph**, a disjoint union of **complete graphs** (or its complement), the **generic K_n -free graph** (or its complement).*

Classification results

For various other families, those members that are homogeneous have been completely determined.

Some classification results

	Finite	Countably infinite
Posets	(trivial)	Schmerl (1979)
Tournaments	Woodrow (1976)	Lachlan (1984)
Graphs	Gardiner (1976)	Lachlan & Woodrow (1980)
Digraphs	Lachlan (1982)	Cherlin (1998)

Weakening homogeneity

Symmetry properties

Strong		Weak
homogeneous (classification possible)	\rightsquigarrow	vertex transitive (classification impossible)

Idea. Consider natural weakenings of homogeneity and the resulting classification problems.

Question. How much can homogeneity be weakened before the corresponding classification problem becomes impossible?

Connected-homogeneous graphs

Definition


A graph Γ is **connected-homogeneous** if any isomorphism between *connected* finite induced subgraphs extends to an automorphism.

Connected-homogeneity...

1. is a natural weakening of homogeneity;
2. gives a class of graphs that lie between the (already classified) homogeneous graphs and the (not yet classified) distance-transitive graphs.

homogeneous \Rightarrow connected-homogeneous \Rightarrow distance-transitive

A finite connected-homogeneous graph




Isomorphisms between finite induced *connected* subgraphs, e.g.

$$(v_1, v_2, v_3) \mapsto (v_6, v_5, v_4),$$

clearly all extend to automorphisms.

So the hexagon is connected-homogeneous.

A finite connected-homogeneous graph



Isomorphisms between finite induced *connected* subgraphs, e.g.

$$(v_1, v_2, v_3) \mapsto (v_6, v_5, v_4),$$

clearly all extend to automorphisms.

So the hexagon is connected-homogeneous.

But we have already seen that it is not homogeneous.

Finite connected-homogeneous graphs

Theorem (Weiss (1976), Gardiner (1978))

A finite graph is connected-homogeneous if and only if it is isomorphic to a disjoint union of copies of one of the following:

1. *a finite homogeneous graph*
2. *bipartite “complement of a perfect matching”*
(obtained by removing a perfect matching from a complete bipartite graph $K_{s,s}$)
3. *cycle C_n*
4. *the graph $K_s \times K_s$*
5. *Petersen’s graph*
6. *the graph obtained by identifying *antipodal* vertices of the 5-dimensional cube Q_5*

Infinite connected-homogeneous graphs


The graph $X_{r,l}$

Take l copies of the complete graph K_r and amalgamate them at a single vertex.

Repeat the process, building a tree-like graph.

These graphs are all connected-homogeneous.

(They arise in [Macpherson's \(1982\)](#) classification of infinite locally-finite distance transitive graphs.)



Infinite connected-homogeneous graphs

Theorem (RG, Macpherson (2009))

A countable graph is connected-homogeneous if and only if it is isomorphic to the disjoint union of a finite or countable number of copies of one of the following:

1. *a finite connected-homogeneous graph;*
2. *a homogeneous graph;*
3. *the random bipartite graph;*
4. *bipartite infinite complement of a perfect matching;*
5. *the graph $K_{\aleph_0} \times K_{\aleph_0}$;*
6. *a treelike graph X_{κ_1, κ_2} with $\kappa_1, \kappa_2 \in (\mathbb{N} \setminus \{0\}) \cup \{\aleph_0\}$.*

Open problems

Digraphs

- ▶ [Cherlin \(1998\)](#) classified the countable homogeneous digraphs.
- ▶ There are 2^{\aleph_0} such graphs.

Problem 1. Classify the countably infinite connected-homogeneous digraphs.

Open problems

Digraphs

- ▶ Cherlin (1998) classified the countable homogeneous digraphs.
- ▶ There are 2^{\aleph_0} such graphs.

Problem 1. Classify the countably infinite connected-homogeneous digraphs.

Problem 2. Classify the **locally-finite** countably infinite connected-homogeneous digraphs.

- ▶ In recent joint work with **R. Möller** we have obtained a partial solution to Problem 2, for digraphs with more than one “end”.
- ▶ Our work relates the problem to the **highly-arc-transitive** digraphs of Cameron, Praeger and Wormald (1993).

Alternative ways of weakening homogeneity

Definition

A relational structure M is **set-homogeneous** if whenever two finite substructures U and V are isomorphic, there is an automorphism $g \in \text{Aut}(M)$ such that $Ug = V$.

Alternative ways of weakening homogeneity

Definition

A relational structure M is **set-homogeneous** if whenever two finite substructures U and V are isomorphic, there is an automorphism $g \in \text{Aut}(M)$ such that $Ug = V$.

- ▶ It is a concept originally due to [Fraïssé and Pouzet](#).
- ▶ [Ronse \(1978\)](#) and [Enomoto \(1981\)](#) showed that for finite graphs:

set-homogeneous \equiv homogeneous.

- ▶ [Droste, Giraudet, Macpherson and Sauer \(1994\)](#) considered infinite set-homogeneous graphs.
 - ▶ **Open problem.** Classify the countable set-homogeneous graphs.

Alternative ways of weakening homogeneity

Definition

A relational structure M is **set-homogeneous** if whenever two finite substructures U and V are isomorphic, there is an automorphism $g \in \text{Aut}(M)$ such that $Ug = V$.

- ▶ It is a concept originally due to [Fraïssé and Pouzet](#).
- ▶ [Ronse \(1978\)](#) and [Enomoto \(1981\)](#) showed that for finite graphs:

set-homogeneous \equiv homogeneous.

- ▶ [Droste, Giraudet, Macpherson and Sauer \(1994\)](#) considered infinite set-homogeneous graphs.
 - ▶ **Open problem.** Classify the countable set-homogeneous graphs.

Recent joint work with [D. Macpherson](#), [C. E. Praeger](#) and [G. Royle](#):

- ▶ We have classified the finite set-homogeneous digraphs.
 - ▶ This generalises work of Lachlan (1981).

References



G. L. Cherlin.

The classification of countable homogeneous directed graphs and countable homogeneous n -tournaments.

Mem. Amer. Math. Soc., 131(621):xiv+161, 1998.



H. Enomoto.

Combinatorially homogeneous graphs.

J. Comb. Theory Ser. B, 30:215–223, 1981.



D. M. Evans.

Examples of \aleph_0 -categorical structures.

In *Automorphisms of first-order structures*, OUP 1994.



A. Gardiner.

Homogeneous graphs.

J. Comb. Theory Ser. B 20, 94–102, 1976.



R. Gray and D. Macpherson

Countable connected-homogeneous graphs.

J. Combin. Theory Ser. B, 2009 (in press).



A. H. Lachlan and R. E. Woodrow.

Countable ultrahomogeneous undirected graphs.

Trans. Amer. Math. Soc., 262(1):51–94, 1980.