

**Measurement of the Deeply Virtual Neutral Pion
Electroproduction Cross Section at the Thomas
Jefferson National Accelerator Facility at 10.6
GeV**

by

Robert Johnston

Submitted to the Department of Physics
in partial fulfillment of the requirements for the degree of

Interdisciplinary PhD in Physics and Statistics

at the

MASSACHUSETTS INSTITUTE OF TECHNOLOGY

August 2023

© Massachusetts Institute of Technology 2023. All rights reserved.

Author
.....

Department of Physics
August, 2023

Certified by
.....

Richard Milner
Professor of Physics
Thesis Supervisor

Accepted by
.....

Lindley Winslow
Associate Department Head of Physics

**Measurement of the Deeply Virtual Neutral Pion
Electroproduction Cross Section at the Thomas Jefferson
National Accelerator Facility at 10.6 GeV**

by

Robert Johnston

Submitted to the Department of Physics
on August, 2023, in partial fulfillment of the
requirements for the degree of
Interdisciplinary PhD in Physics and Statistics

Abstract

Deeply virtual exclusive reactions provide unique channels to study both transverse and longitudinal properties of the nucleon simultaneously, allowing for a 3D image of nucleon substructure. This presentation will discuss work towards extracting an absolute cross section for one such exclusive process, deeply virtual neutral pion production, using 10.6 GeV electron scattering data off a proton target from the CLAS12 experiment in Jefferson Lab Hall B . This measurement is important as exclusive meson production has unique access to the chiral odd GPDs, and is also a background for other exclusive processes such as DVCS, making the determination of this cross section crucial for other exclusive analyses.

Thesis Supervisor: Richard Milner
Title: Professor of Physics

Acknowledgments

To Be Completed.

Contents

List of Figures	8
List of Tables	11
List of Equations	12
List of To Do Items	13
1 Introduction	15
1.1 Exploring Structure through Scattering	17
1.1.1 Scattering at Different Resolution Scales	17
1.1.2 Elastic Scattering and Form Factors	19
1.1.3 Inelastic Scattering and Parton Distribution Functions	23
1.2 Process Background	29
1.2.1 Wigner Functions, Generalized Parton Distributions	31
1.2.2 Deeply Virtual Exclusive Processes	35
1.2.3 Status of DV π^0 P Measurements	39
2 Experiment and Data Processing	45
2.1 Accelerator and Beamline	45
2.1.1 Thomas Jefferson National Accelerator Facility	47

2.1.2	CLAS12 Beamline	49
2.2	CLAS Detectors and Run Conditions	50
2.2.1	CLAS Detector System	50
2.2.2	Run Conditions	74
2.3	Reconstruction and Particle Identification	77
2.3.1	Decoding and Track Reconstruction	77
2.3.2	Particle Identification	77
2.3.3	Data Storage and Formatting	78
3	Simulations	81
3.1	Computing and Simulation Infrastructure	81
3.1.1	MIT Tier 2 and Supercomputing Clusters	81
3.1.2	Overview of Simulation Runs	81
3.2	Event Generation	82
3.2.1	Nonradiative Generator	84
3.2.2	Radiative Generator	84
3.3	Simulation Pipeline	84
3.3.1	Simulation Processing and JLab Computing	84
3.3.2	Geant4, GEMC, and CLAS Submissions	84
3.4	Simulation Enhancements with Normalizing Flows	84
3.4.1	Inverse Transforms and Autoregressive Flows	84
3.4.2	Exploration of Simulation Speedup with UNMAF	85
4	Cross Section Measurement	87
4.1	General Analysis Overview	88
4.1.1	Part 1	88

4.1.2 part 2	88
4.2 Data Pre-Processing	91
4.2.1 Experimental Data Pre-Processing	91
4.2.2 Simulation Data Pre-Processing	91
4.3 Event Selection	92
4.3.1 Exclusivity Cuts	92
4.3.2 Kinematic Fitting	99
4.4 Luminosity	99
4.5 Binning	102
4.6 Correction Factors	103
4.6.1 Acceptance Correction	103
4.6.2 Radiative Corrections	103
4.6.3 Binning Corrections	103
4.6.4 Overall Normalization Corrections	103
4.7 Uncertainty Estimation	103
4.8 Iterative Bayesian Unfolding	103
4.8.1 Acceptance Correction	103
4.8.2 Radiative Corrections	103
4.8.3 Binning Corrections	103
4.8.4 Overall Normalization Corrections	103
A Full Cross Section Data	115
B BSA Cross Check	117
C Derivation of phi math convention	119

List of Figures

1-1	J.J. Thomson's Plum Pudding Model	16
1-2	Elastic scattering diagram	19
1-3	Tree-level elastic scattering Feynman diagram	21
1-4	Fourier Transforms of Charge Distributions	23
1-5	Feynman Diagram for Inelastic Scattering	24
1-6	Scattering Cross Section vs. Energy Transfer	25
1-7	Proton Structure Function F_2^p	28
1-8	Parton Distribution Functions	29
1-9	Scattering Measurements at Different Scales	30
1-10	Distribution Relationship Cube	34
1-11	DVCS and DVMP Feynman Diagrams	35
1-12	Proton Pressure Distribution, Experiment	36
1-13	Proton Pressure Distribution, Theory	36
1-14	DVPiP Feynman Diagram	37
1-15	Diagram of Lepton-Hadron Plane Angle ϕ	38
1-16	Kinematic Reach Comparison	41
1-17	Analysis Flowchart	43
2-1	CLAS12 Detector System	46
2-2	CLAS12 Specification	46

2-3	CLAS12 Data Rates, Compared to Other Experiments	47
2-4	Low Threshold Cherenkov Counter	56
2-5	Silicon Vertex Tracker	69
2-6	74
2-7	75
2-8	76
2-9	The distribution for mass of two photons $M_{\gamma\gamma}$	79
2-10	The distribution for mass of two photons after exclusivity cuts	79
4-1	High-level data processing flow	90
4-2	Comparison of experiment (blue) and simulation (red) missing mass, energy, momentum, and invariant gamma-gamma mass distributions, before any smearing factors were added to the simulation data.	91
4-3	Comparison of experiment (blue) and simulation (red) missing mass, energy, momentum, and invariant gamma-gamma mass distributions, with smearing factors added to the simulation data proton and photon momenta.	92
4-4	MM^2 (epX) vs $\theta_X\pi$ 2D distribution.	93
4-5	Exclusive distributions for events with at least one electron, proton and two photons.	93
4-6	Exclusive distributions for events with at least one electron, proton and two photons.	94
4-7	Exclusive distributions for events with at least one electron, proton and two photons.	94
4-8	Exclusive distributions after tight $M_{\gamma\gamma}$ mass and transverse missing momenta cuts	95

4-9	$MM^2(epX)$ distributions for multiple $\theta_{X\pi}$ cut values.	96
4-10	The numbers of signal (red markers) and background (black markers) events as functions of $\theta_{X\pi}$ cut value for multiple Q^2 bins.	97
4-11	The numbers of signal (red markers) and background (black markers) events as functions of $\theta_{X\pi}$ cut value for multiple x_B bins.	97
4-12	Exclusive distributions after all exclusivity cuts	98
B-1	BSA Cross Check Results	118

List of Tables

1.1	GPDs Across Nucleon and Quark Polarizations	33
4.1	Terms of Luminosity Equation	101

List of Equations

1.1	Diffraction Limit	18
1.2	de Broglie Relationship	18
1.3	Rutherford Scattering Differential Cross Section	20
1.4	Mott Scattering Differential Cross Section	20
1.5	Elastic Scattering Differential Cross Section	21
1.8	Rosenbluth Scattering Cross Section	22
1.9	Outgoing Hadronic System Invariant Mass W	23
1.10	Bjorken X	25
1.11	Inelasticity y	25
1.12	Energy Transfer ν	26
1.13	Lorentz-invariant Rosenbluth Scattering Cross Section	26
1.14	Scattering Cross Section in Terms of $F_1(x_B, Q^2)$ and $F_2(x_B, Q^2)$	26
1.15	F_2 structure function in terms of PDFs	27
1.16	Wigner Quasi-probability Distributions	32
1.17	Wigner Operator	32
1.18	Reduced Wigner Distribution	32
1.21	DV π P Cross Section Decomposition	37
1.22	Virtual Photon Flux	38
4.5	DV π P Experimental Cross Section	89

List of To Do Items

1.14	input standard model, pion information	26
1.27	statement introducing idea of posing this as an inverse problem, introduce the experimental interpretation of the cross section	42

Chapter 1

Introduction

Humans have tried to understand the nature of the world around us for millennia, with discerning the structure of matter being a central effort in this quest. Famously, the Greek philosophers Leucippus and Democritus (\sim 5th century BCE) are credited with the concept of “atomism” - the belief that matter is composed of tiny indivisible particles called atoms (from the Greek $\alpha\tauομοσ$, roughly translating to “uncuttable” ([C.C.W. Taylor, 1999](#)). Even further back, there are Indian records from as early as the 8th century BCE conceptualizing the world as being built from tiny fundamental particles ([Thomas McEvilley, 2002](#)).

Scientific progress on this front stalled until the early 1800s, when chemists explored how different elements combined in to form compounds in specific, repeatable, small integer ratios. John Dalton formulated this idea as the Law of Multiple Proportions, which paved the way for early scientific atomic theory ([Britannica, 2010](#)). In 1897, J.J. Thomson discovered the first subatomic particle, the electron, by studying cathode rays([J. Thomson, 1901](#)). Accordingly, he devised a model of the atom which had electrons embedded in a ball of positively charged material, called the Thomson,




Figure 1-1: J.J. Thomson’s Plum Pudding Model of the atom ([Britannica, 2023](#))

or Plum Pudding, Model([Jaume Navarro, 1995](#)).

With the idea that the atom was a composite object, scientist began experimentation to study its exact structure. This was the start of what has been the 120 years of particle scattering studies to probe first atomic, then nuclear structure.

The rest of this chapter details the findings of previous scattering experiments and provides a background on Generalized Parton Distributions and Deeply Virtual Exclusive Processes, the topic of this work. Chapter 2 describes the experimental setup of the CLAS12 detector and data taking conditions. Chapter 3 discusses data analysis procedures to reconstruct particles and classify events from detector level information. Simulations and computational pipelines for this work are presented in chapter 4. The analysis procedure for combining experimental and simulated data into a differential cross section with correction factors is discussed in chapter 5.

Chapter 6 displays and discusses results and uncertainties. Chapter 7 summarizes this work and lays a path for finalizing the measurement. The appendices include numerous technical details and supplemental plots.

1.1 Exploring Structure through Scattering

The typical length scales for atoms and nucleons are 0.1 nm which is far smaller than the wavelength of human-visible light (~ 500 nm). As such, atomic and nuclear structure must be explored by forcing some interaction and then inferring the structure from the observed results. Thomson's atomic model was famously tested in the early 1900s by Ernest Rutherford's research group, wherein α particles were fired at thin metal targets, and the scattering behaviour was observed ([Geiger and Marsden, 1909](#)) ([Rutherford, 1911](#)).

The results were not consistent with Thomson's model, but instead indicated that there was a very small, dense, positively charged nucleus at the center of every atom. Further experiments by Rutherford would lead to the discovery of the proton around 1920 ([Rutherford, 1919](#)). Puzzles about the nucleus remained, including a consistent description of isotopes, until 12 years later when James Chadwick suggested the existence of the neutron ([Chadwick, 1932](#)). With electrons and the two nucleons discovered, it seemed as though the indivisible constituents of the atom were finally realized, but future experiments showed a much more complex, sub-nuclear structure.

1.1.1 Scattering at Different Resolution Scales

The diffraction limit for microscopic (compared to telescopic) systems can be approximated by equation [1.1](#), where n is the index of refraction, θ is a measure of the

device aperture, λ is the wavelength of the probe, and d is the minimum resolvable length scale. Thus, the wavelength of a probe sets a fundamental lower limit on the achievable resolution of a microscopic imaging system - roughly, at small enough distances, the probe's waves interfere, prohibiting resolution at or below that scale.

$$d = \frac{\lambda}{2n \sin \theta} \quad (1.1)$$

For visible light microscope systems, $\lambda \sim 500$ nm, and so the minimum resolvable feature size is approximately $d \sim 250$ nm. Techniques exist to extend the resolution size by approximately an order of magnitude, e.g. expansion microscopy ([Chen, Tillberg, and Boyden, 2015](#)) or Near-Field Scanning Optical Microscopy ([Ma et al., 2021](#)), but non-visible-light probes are needed for scales below ~ 10 nm.

In particular, the de Broglie relationship [1.2](#) ([Broglie, 1924](#)) states that the wavelength λ of a particle is inversely proportional to its momentum p , with h being Planck's constant.

$$\lambda = \frac{h}{p} \quad (1.2)$$

With this relationship, we can see that by increasing a particle's momentum, its effective wavelength is reduced. This is the fundamental principle which allows electron microscopes to image matter at a resolution of $\sim 10\text{-}0.1$ nm ([Franken et al., 2020](#)), corresponding to electron momenta of $\sim 1\text{-}100$ keV. At this scale, viruses, cells, molecular structures, and even atoms can be imaged ([Williams and Carter, 2009](#)), with striking results commonly published online. Other probes could be used to circumvent the diffraction limit, such as high energy (low-wavelength) photons or

high momentum (low de Broglie wavelength) protons or neutrons, but electrons are an ideal candidate in this regime as they are easy to produce, steer, interact with, and detect.

To move beyond imaging at the atomic scale ($\sim 1 \text{ \AA}$) to the nuclear scale ($\sim 1 \text{ fm}$) requires probes that are 100,000 times more powerful. Electrons are still an ideal probe due to their (apparent) lack of internal structure, but rather than a room sized microscope, an entire accelerator facility is needed to achieve high enough energies and luminosities for sub-nuclear scale resolution.

1.1.2 Elastic Scattering and Form Factors

Imaging with electrons (or other non-visible-light probes) at any energy scale is commonly understood in terms of scattering cross sections, σ , with dimensions of area and interpreted as the probability for a certain interaction to occur. Typical elastic scattering cross sections for transitions metals with 100 keV incident electrons as in electron microscopy are $\sim 10^{-22} \text{ m}^2$ (Williams and Carter, 2009). In contrast, the cross sections to be discussed in this thesis are on the order of tens of nanobarn (10^{-36} m^2), or 14 orders of magnitude smaller.




Figure 1-2: Elastic scattering diagram

The scattering cross section for a probe (such as an electron) incident on a target, can be calculated at lowest order by considering a fixed (no recoil), point-like (no structure), radially symmetric Coulomb potential (e.g., a proton) with a non-relativistic incident charged particle. The resulting equation was used by Rutherford's group in the discovery of the nucleus, and for an electron beam of energy E_{beam} is given by (1.3), where α is the fine structure constant.

$$\frac{\theta}{2} \left(\frac{d\sigma}{d\Omega} \right)_{Ruth} = \frac{\alpha^2}{16E_{beam}^2 \sin^4(\theta/2)} \quad (1.3)$$

To probe smaller resolution scales, it is necessary to increase the energy of the beam, and eventually the probe must be treated relativistically. This correction term is provided by the Mott scattering cross section, given by (1.4), which still assumes a fixed, point-like target, with only Coulomb interactions.

$$\left(\frac{d\sigma}{d\Omega} \right)_{Mott} = \frac{\alpha^2}{4E^2 \sin^4(\theta/2)} \cos^2 \frac{\theta}{2} = \left(\frac{\alpha}{2E \sin^2(\theta/2)} \cos \frac{\theta}{2} \right)^2 = 4 \cos^2 \frac{\theta}{2} \left(\frac{d\sigma}{d\Omega} \right)_{Ruth} \quad (1.4)$$

At higher incident electron energies (and thus finer spatial resolutions), the proton's finite size must be accounted for, as well as the momentum transferred to it. The tree-level Feynman diagram for elastic electron-proton scattering is show in Fig. 1-3. The incoming electron e exchanges a virtual photon with the proton p , resulting in a momentum transfer of $q = p_{e'} - p_e$.




Figure 1-3: Tree-level elastic scattering Feynman diagram

The momentum transfer q sets the resolution scale for these processes, but it is convenient to work with the negative square of this value, defined as $Q^2 = -q^2$. With this term, we can express the relativistic differential cross section for the scattering of electrons off a resting, point-like proton as in (1.5), where m_p is the mass of the proton.

$$\frac{d\sigma}{d\Omega} = \frac{\alpha^2}{4E_{beam}^2 \sin^4(\theta/2)} \frac{E_{e'}}{E_{beam}} \left(\cos^2 \frac{\theta}{2} + \frac{Q^2}{2m_p^2} \sin^2 \frac{\theta}{2} \right) \quad (1.5)$$

Compared with Mott Scattering, there are two differences in this formula: The $\frac{E_{e'}}{E_{beam}}$ term in the scattering cross section comes from the electron losing energy to the proton's final state kinetic energy (no longer fixed), and the term proportional to $\sin^2(\theta/2)$ is due to a purely magnetic spin-spin interaction.

If the proton were a point, then (1.5) would agree with experiment for all electron scattering energies. Instead, deviations are observed as we increase the beam energy.

To account for this structure, we need to include two form factors, $G_E(Q^2)$ - related to the distribution of charge, and $G_M(Q^2)$, related to the distribution of magnetism inside the proton. In the low- Q^2 limit, these form factors are the Fourier transforms of the charge and magnetic moment distributions as in (1.6) and 1.7, reducing to the charge and the magnetic moment of the proton in the $Q^2 = 0$ limit.

$$G_E(Q^2) \approx G_E(q^2) = \int e^{iq \cdot r} \rho(r) d^3r \quad G_E(0) = \int \rho(r) d^3r = 1 \quad (1.6)$$

$$G_M(Q^2) \approx G_E(q^2) = \int e^{iq \cdot r} \mu(r) d^3r \quad G_M(0) = \int \mu(r) d^3r = 2.79 \quad (1.7)$$

Including these form factors in our cross section gives us the full elastic scattering cross section, as shown in (1.8).

$$\frac{d\sigma}{d\Omega} = \frac{\alpha^2}{4E_1^2 \sin^4(\theta/2)} \frac{E_3}{E_1} \left(\frac{G_E^2 + \tau G_M^2}{1 + \tau} \cos^2 \frac{\theta}{2} + 2\tau G_M^2 \frac{Q^2}{2m_p^2} \sin^2 \left(\frac{\theta}{2} \right) \right) \quad (1.8)$$

Where $\tau = \frac{Q^2}{4m_p^2}$.




Figure 1-4: Samples of charge distributions and their corresponding form factors $F(\mathbf{q}^2)$, from ([M. Thomson, 2013](#))

By the 1960s, elastic scattering had been studied sufficiently well as to measure the proton form factors up to several GeV in Q^2 . The observed results were consistent with a proton having a ‘dipole’ form factor, as shown in Fig. 1-4. Investigating proton structure at finer spatial resolutions requires increasing the beam energy, but eventually the the elastic scattering cross section becomes negligible and instead the interactions are sufficiently energetic so as to create additional particles.

1.1.3 Inelastic Scattering and Parton Distribution Functions

Elastic scattering can be defined as interactions where the target stays intact; specifically, the variable W is the invariant mass of the outgoing struck target (1.9), where elastic scattering satisfies the condition $W^2 = m_p^2$. If $W^2 > m_p^2$, we instead have inelastic scattering, written as $e p \rightarrow e' X$, where X stands for some outgoing hadronic system, as shown in the Feynman diagram in Fig. 1-5.

$$W^2 \equiv (p_p + q)^2 = (p_p + (p_e - p_{e'}))^2 \quad (1.9)$$




Figure 1-5: Feynman Diagram for Inelastic Scattering

The cross section for inelastic scattering has several peaks at various proton resonances, as indicated in the top sketch of Fig. 1-6. Continuing to higher energy transfers we reach the ‘Deep Inelastic Scattering’ (DIS) regime, defined by kinematics as $Q^2 > 1 \text{ GeV}^2$ and $W^2 > 4 \text{ GeV}^2$. Note that in the DIS process, the proton is smashed apart, yielding many subparticles. Other high-energy inelastic processes where the proton is left intact will be discussed in section 1.2.1.

Since we remove the constraint that the mass of the final state is the proton mass, we now have one extra degree of freedom, i.e., we need at least 2 variables to describe scattering here. Convenient choices are the squared four-momentum transfer of the virtual photon Q^2 and Bjorken X , defined in (1.10). x_B is a measure of elasticity: $x_B = 1$ for elastic scattering. It is useful in that it can also be interpreted as the fraction of proton momentum carried by the struck quark in the infinite momentum frame.




Figure 1-6: Sketch of cross section as a function of electron energy transfer for inclusive electron scattering off a proton (top) and a nucleus (bottom), from ([T. W. Donnelly et al., 2017](#))

$$x_B \equiv \frac{Q^2}{2p_p \cdot q} = \frac{Q^2}{Q^2 + W^2 - m_p^2} \quad (1.10)$$

Another useful quantity is y , which is a measure of the inelasticity of the scattering. It is the fractional energy lost by the electron in the scattering process, where $y=0$ is for perfectly elastic collisions, and is given by (1.11)

$$y \equiv \frac{p_p \cdot q}{p_p \cdot p_e} = \frac{\nu}{E_{beam}} = 1 - \frac{E_{e'}}{E_{beam}} \quad (1.11)$$

Where ν is the energy transferred in the collision (1.12).

$$\nu = \frac{Q^2}{2 * x_B * m_p} \quad (1.12)$$

With these definitions, we can write the differential cross section for inelastic scattering. Note that the general formula for the differential cross section for elastic scattering, (1.8) can be re-written in explicitly Lorentz-invariant form as in (1.13).

$$\frac{d\sigma}{d\Omega} = \frac{4\pi\alpha^2}{Q^4} \left[\left(1 - y - \frac{m_p^2 y^2}{Q^2}\right) \frac{G_E^2 + \tau G_M^2}{1 + \tau} + y^2 \frac{G_M^2}{2} \right] \quad (1.13)$$

This equation can be generalized to extend to inelastic scattering by replacing the terms corresponding to the combinations of form factors G_E and G_M with more structure functions $F_1(x_B, Q^2)$ and $F_2(x_B, Q^2)$, which describe proton structure as a function of both independent variables. This results in the differential cross section given by (1.14) .

$$\frac{d\sigma}{d\Omega} = \frac{4\pi\alpha^2}{Q^4} \left[\left(1 - y - \frac{m_p^2 y^2}{Q^2}\right) \frac{F_2(x_B, Q^2)}{x_B} + y^2 F_1(x_B, Q^2) \right] \quad (1.14)$$

input standard model, pion information

Experiments in the 1960s on DIS indicated that the structure functions $F_1(x_B, Q^2)$ and $F_2(x_B, Q^2)$ were nearly independent of Q^2 , a feature known as Bjorken scaling ([Bjorken and Paschos, 1969](#)). This indicated scattering was occurring off of point-like constituents - current experiment results provide a constraint on the maximum radius of these constituent to be at most 10–18 m ([M. Thomson, 2013](#)).

Secondly, DIS results indicated the two structure functions could be expressed as $F_2(x_B) = 2 * x_B * F_1(x_B)$, named the Callan-Gross relation. This relationship can be explained if the electron is scattering off of spin-half point-like particles inside the proton, which combined with Bjorken scaling to give strong evidence for the

existence of quarks inside the proton, and gave motivation for the development of the parton model ([Feynman, 1969](#)).

The parton model connects the experimentally measurable structure functions to Parton Distribution functions which describe the distribution of proton longitudinal momentum amongst its constituents. Specifically, a PDF $q_i(x_B)$ describes the probability density of finding a parton carrying a longitudinal momentum fraction in the interval $(x_B, x_B + dx_B)$. The relationship between PDFs and structure functions is given by [\(1.15\)](#), where Q_i is the charge of each quark.

$$F_2^p(x_B) = x_B \sum_i Q_i^2 q_i(x_B) \quad (1.15)$$

Structure functions have been studied in great detail over a very large kinematic range across Q^2 and x_B , the results of which are shown in [Fig. 1-7](#).




Figure 1-7: The proton structure function F_2^p , measured at various experiments as listed, all with $W^2 > 3.5\text{GeV}^2$. F_2^p values have been multiplied by 2^{ix} for visual purposes, from ([Zyla et al., 2020](#))

The global experimental results can be combined with theoretical QCD frameworks such as the DGLAP evolution equations ([Altarelli and Parisi, 1977](#)). to plot PDFs for various constituents of the proton, as shown in Fig. 1-8.




Figure 1-8: Quark and gluon distribution functions from NNLO NNPDF3.0 global analysis at $\mu^2 = 10 \text{ GeV}^2$ (left) and $\mu^2 = 10^4 \text{ GeV}^2$ (right), from (Zyla et al., 2020)

All of these major scattering scales explored through the 20th century are summarized in Fig. 1-9, spanning roughly four orders of magnitude in length scale. While steady increases in resolving power have been made, the focus of this work (red triangle in figure) is not to image even finer scale parton dynamics, but rather to understand the multidimensional structure of the nucleon. In particular, while PDFs allow for a 1 dimensional mapping of the inner workings of a proton, even more information can be gleaned from more complex scattering reactions. Efforts are now directed towards so called *proton tomography* - 3D imaging of nucleon structure - which is the focus of this analysis.

1.2 Process Background

The term *tomography* is derived from the Greek word *tomos*, which translates to *slice* or *section*. In the medical field, CT scans (computed tomography) combine




Figure 1-9: Scattering experiments performed at different energy scales reveal different information about proton structure. This work (red triangle) focuses on multi-dimensional structure mapping in the valence quark regime. Figure modified from ([Klein, 2005](#))

many 2-D images from X-ray scans to generate a three-dimensional reconstruction of bodily organs. Building on this, proton tomography harnesses many nuclear reactions to reconstruct multi-dimensional mappings of partons' spatial and momentum distributions inside nucleons.

1.2.1 Wigner Functions, Generalized Parton Distributions

In classical mechanics, a particle can be completely described by its position in its six-dimensional phase space (three spatial and three momentum coordinates). An ensemble of such particles can be most completely understood through its phase space distribution function, which contains the probability of finding a particle in a particular region in phase space. In quantum systems, a pure phase space distribution is not well defined because of Heisenberg's uncertainty principle. However, in 1932 Eugene Wigner introduced a formalism that addressed this ([Wigner, 1932](#)), yielding functions provide the most comprehensive representation achievable for quantum systems.

Wigner Quasi-probability Distributions

Wigner Quasi-probability Distributions, commonly referred to as simply Wigner functions or Wigner distributions, are defined as in [\(1.16\)](#). This can be integrated over x (p) to yield momentum (space) density, but for arbitrary (x,p) the distribution can take negative values, and so violates probability axioms and thus is a quasi-probability distribution rather than a full probability distribution. Wigner distributions are useful outside of particle physics, notably in signal processing, with more details available in [\(Hillery et al., 1984\)](#).

$$W(x, p) = \frac{1}{\pi \hbar} \int_{-\infty}^{\infty} \psi^*(x + \eta) \psi(x - \eta) e^{2ip\eta/\hbar} d\eta \quad (1.16)$$

The corresponding generalization to relativistic quark and gluon phase space distributions is covered in (Ji, 2004) to yield a Wigner Operator (1.17) which can be used to obtain the reduced quantum phase-space quark distributions in the nucleon (1.18).

$$\hat{W}_{\Gamma}(\vec{r}, k) = \int_{-\infty}^{\infty} e^{ik \cdot \eta} \bar{\Psi}(\vec{r} - \eta/2) \Gamma \Psi(\vec{r} - \eta/2) d^4\eta \quad (1.17)$$

$$W_{\Gamma}(\vec{r}, \vec{k}) = \int \frac{dk^-}{4\pi^2} \frac{1}{2} \int \frac{d^3\vec{q}}{8\pi^3} e^{-i\vec{q} \cdot \vec{r}} \langle \vec{q}/2 | \hat{W}_{\Gamma}(\vec{r}, k) | -\vec{q}/2 \rangle \quad (1.18)$$

Here we have integrated over $k^- = (k^0 - k^z)/\sqrt{2}$, the light-cone energy, since it is difficult to measure in high-energy processes.

Generalized Parton Distributions

No known experiments currently exist that are able to directly measure this distribution (nor is it known if it is possible). Fortunately, in recent decades theorists have been able to link experimental observables to further reduced forms of (1.18). Specifically, integration can be performed over spatial coordinates to yield Transverse Momentum Distributions (TMDs) which are outside the scope of this work. Alternatively, integration can be performed over momentum coordinates to yield Generalized Parton Distributions (GPDs), which encode transverse spatial as well as

longitudinal momentum distributions of partons inside the nucleon. As shown in (Ji, 2004), at leading-twist (twist-2), iterating through all choices of the Dirac matrix for quark distributions Γ yields 8 distinct GPDs. They are generally expressed in terms of parton momentum fraction x , skewness $\xi = \frac{-q^2}{q \cdot P} \sim \frac{x_B}{2-x_B}$, and momentum transfer t .

4 correspond to helicity conserving (chiral even) processes and 4 correspond to helicity flipping (chiral odd) processes: H , E , \tilde{H} , and \tilde{E} for chiral even, and H_T , E_T , \tilde{H}_T , and \tilde{E}_T ($\bar{E}_T = 2^* \tilde{H}_T + E_T$ is commonly used). Table 1.1 summarizes the GPDs with respect to polarization states.

Nucleon		Quark Polarization		
Polarization		U	L	T
U	H	*	$\tilde{E}_T = 2^* \tilde{H}_T + E_T$	
L	*	\tilde{H}		\tilde{E}_T
T	E	\tilde{E}		H_T, \tilde{H}_T

Table 1.1: GPDs Across Nucleon and Quark Polarizations. * forbidden by parity.

GPDs can be understood by considering further integrations and forward limits - in the same way that the nucleon charge must be recovered when integrating over PDFs or Form Factors, these functions themselves are recovered when appropriately integrating over GPDs, as in Fig. 1-10. Specifically, first moments of the GPDs H and E are related to the Dirac and Pauli form factors F_1 and F_2 respectively:




Figure 1-10: Relation cube, from ([M. Burkardt and Pasquini, 2016](#)).

$$\int dx H(x, \xi, t) = F_1(t) \quad (1.19)$$

$$\int dx E(x, \xi, t) = F_2(t) \quad (1.20)$$

GPDs have been shown ([Ji, 1997](#)) to encode spin distributions and also can be interpreted as describing the transverse spatial distribution of quarks ([Matthias Burkardt, 2007](#)). Further, GPDs relation to energy-momentum tensor form factors (EMTs) allows access to EMT densities, which describe the distribution of energy, momentum, and pressure inside the nucleon. It was originally not known how to measure GPDs, but were realized to be experimentally accessible through deeply virtual exclusive processes (DVEP) relying on factorization theorems ([Collins and Freund, 1999](#)), ([Bauer et al., 2002](#)).

1.2.2 Deeply Virtual Exclusive Processes

Deeply virtual exclusive processes are nuclear reactions occurring with high photon virtuality ($Q^2 \gg m_p^2$) in the DIS regime ($W^2 \gg m_p^2$), with thresholds normally set to require $Q^2 > 1 \text{ GeV}^2$ and $W > 2 \text{ GeV}$. The processes involve the scattering of a virtual photon off a nucleon target, yielding either a real photon or mesons, along with an intact final state nucleon and incident particle (e.g. from an electron or muon beam). These processes are in stark contrast to DIS, where the nucleon target is shattered into many pieces, and can instead be thought of as a hard yet precise peaceful process. Different reactions have been shown to have different GPD dependencies, and thus provide different windows into sub-nucleon mechanics.

Deeply Virtual Compton Scattering and Meson Production

Fig. 1-11 illustrates diagrams for Deeply Virtual Compton Scattering (DVCS) and Deeply Virtual Meson Production (DVMP).




Figure 1-11: Feynman diagrams for DVCS (left) and DVMP (right), from (Kubarovsky, 2011). DA refers to the appropriate meson distribution amplitude.

DVCS is widely regarded as the “cleanest” channel and has already been leveraged to provide great insights into the structure of the nucleon. For example, Fig. 1-12 shows the pressure distribution inside a proton from DVCS data (V. D. Burkert, Elouadrhiri, and Girod, 2018), which has since been further investigated by theorists to generate similar mappings through LQCD as in Fig. 1-13




Figure 1-12: Proton Pressure Distribution from DVCS data, from (V. D. Burkert, Elouadrhiri, and Girod, 2018).




Figure 1-13: Proton Pressure Distribution from Lattice QCD, from (Shanahan and Detmold, 2019).

DVCS is primarily sensitive to non-quark spin flip GPDs (e.g. H , E), as are DVMP for vector mesons, such as the ρ . On the other hand, pseudoscalar meson production, such as pions, are sensitive to the transversity, or chiral-odd, GPDs. Additionally, some meson processes, such as ϕ due to its strange quark content, are particularly sensitive to gluon GPDs. This work focuses on DVMP with the production of a neutral pion, π^0 , which in addition to accessing chiral-odd GPDs, is an important background for other processes such as DVCS due to sample contamination from π^0 decay (Lee, 2022).

Deeply Virtual Neutral Pion Production

The Feynman diagram for Deeply Virtual Neutral Pion Production (DV π^0 P or DVPiP) is shown in Fig. 1-14.




Figure 1-14: DV π^0 P Feynman Diagram

The unpolarized cross-section for DV π^0 P can be decomposed into longitudinal and transverse structure functions as in (1.21), the formalism for which is covered in (Donnachie and Shaw, 1978), (Dreschsel and Tiator, 1992), and (T. Donnelly, Jeschonnek, and Van Orden, 2023).

$$\frac{d^4\sigma_{ep \rightarrow ep'\pi^0}}{dQ^2 dx_B dt d\phi_\pi} = \Gamma(Q^2, x_B, E) \frac{1}{2\pi} \left\{ \left(\frac{d\sigma_T}{dt} + \epsilon \frac{d\sigma_L}{dt} \right) + \epsilon \cos(2\phi) \frac{d\sigma_{TT}}{dt} + \sqrt{2\epsilon(1+\epsilon)} \cos(\phi) \frac{d\sigma_{LT}}{dt} \right\} \quad (1.21)$$

Here Γ is the virtual photon flux as in (1.22) and represents the number of virtual photons per scattered electron (Amaldi, Fubini, and Furlan, 1979), ϵ is the ratio of transverse to longitudinally polarized photons (1.23), ϕ is the angle between the lepton and hadron planes, as illustrated in Fig. 1-15, and the structure functions

can be expressed as convolutions of GPDs as shown in (1.24)-(1.27) as discussed in (Bedlinskiy et al., 2014).

$$\Gamma(Q^2, x_B, E) = \frac{\alpha}{8\pi} \frac{Q^2}{m_p^2 E^2} \frac{1 - x_B}{x_B^3} \frac{1}{1 - \epsilon} \quad (1.22)$$

$$\epsilon = \frac{1 - y - \frac{Q^2}{4E^2}}{1 - y + \frac{y^2}{2} + \frac{Q^2}{4E^2}} \quad (1.23)$$




Figure 1-15: Diagram of Lepton-Hadron Plane Angle ϕ

$$\frac{d\sigma_L}{dt} = \frac{4\pi\alpha}{kQ^2} \left\{ (1 - \xi^2) |\langle \tilde{H} \rangle|^2 - 2\xi^2 \Re [\langle \tilde{H} \rangle^* \langle \tilde{E} \rangle] - \frac{t'}{4m^2} \xi^2 |\langle \tilde{E} \rangle|^2 \right\} \quad (1.24)$$

$$\frac{d\sigma_T}{dt} = \frac{2\pi\alpha\mu_\pi^2}{kQ^4} \left\{ (1 - \xi^2) |\langle \textcolor{red}{H}_T \rangle|^2 - \frac{t'}{8m^2} |\langle \bar{\textcolor{blue}{E}}_T \rangle|^2 \right\} \quad (1.25)$$

$$\frac{d\sigma_{LT}}{dt} = \frac{4\pi\alpha\mu_\pi}{\sqrt{2}kQ^3} \xi \sqrt{1 - \xi^2} \frac{\sqrt{-t'}}{2m} \Re \left\{ \langle \textcolor{red}{H}_T \rangle^* \langle \tilde{\textcolor{blue}{E}} \rangle \right\} \quad (1.26)$$

$$\frac{d\sigma_{TT}}{dt} = \frac{4\pi\alpha\mu_\pi^2}{kQ^4} \frac{-t'}{16m^2} \langle \bar{\textcolor{blue}{E}}_T \rangle^2 \quad (1.27)$$

The terms involved in these expressions are:

- $t' = t - t_0$ where $t_0 = \frac{-4m^2\xi^2}{1-\xi^2}$
- Skewness $\xi = \frac{x_B}{2-x_B}$
- The bracket $\langle \tilde{F} \rangle$ is the convolution of a GPD and an appropriate subprocess amplitude: $\langle \tilde{F} \rangle = \sum_\lambda \int_{-1}^1 d\bar{x} H_{0\lambda,0\lambda}(\bar{x}, \xi, Q^2, t=0) \tilde{F}(\bar{x}, \xi, Q^2, t)$
 - λ is the unobserved helicities of the partons participating in the subprocess
- Phase space factor $k = 16\pi \left(W^2 - m^2 \right) \sqrt{\Lambda(W^2, -Q^2, m^2)}$
 - $\Lambda(W^2, -Q^2, m^2)$ is the Källén function: $W^4 + Q^4 + m^4 + 2W^2Q^2 + 2Q^2m^2 - 2W^2m^2$
- Reduced pion mass $\mu_{\pi^0} = \frac{m_{\pi^0}^2}{m_u + m_d}$
 - m_u and m_d are respective masses of up and down quarks

1.2.3 Status of DV π^0 P Measurements

With theoretical advancements occurring in the mid 1990s to early 2000s, the first analyses of experimental measurements of DV π^0 P have only been released in the

past decade.

Summary of Existing Measurements

The earliest of such measurements were taken at the Thomas Jefferson National Accelerator Facility (JLab) with a ~ 6 GeV electron beam. Two of the four experimental halls - Hall A ([Fuchey et al., 2011](#)) and Hall B ([Bedlinskiy et al., 2014](#)) produced cross-section results. Hall A houses a small acceptance precision spectrometer and recorded data in several kinematic bins. Hall B housed a large acceptance spectrometer, yielding cross-section measurements over a large kinematic regime.

Recent upgrades at JLab have nearly doubled the beam energy to 10.6 GeV, and both detector halls have accumulated data allowing for the measurement of this process. Both halls repeated data taking at this higher energy, with Hall A recently releasing updated cross-section values across three fixed x_B bins (0.36, 0.48 and 0.6) and over a range of Q^2 values from 3 to 9 GeV 2 ([Dlamini et al., 2021](#)).

This work expands on these results by covering a much larger kinematic regime, as well as having much higher statistics compared to the 6 GeV Hall B result. A kinematic overlap plot in Q^2 , x_B is shown in Fig. [1-16](#) summarizes these differences. It is also noted that the CERN COMPASS collaboration ([Alexeev et al., 2020](#)) has measured this process using a 160 GeV muon beam, obtaining results over a much lower x_B range.




Figure 1-16: Kinematic reach plot between this work and other results: CLAS6 ([Bedlinskiy et al., 2014](#)), COMPASS ([Alexeev et al., 2020](#)), Hall A 6 GeV ([Fuchey et al., 2011](#)), and Hall A 12 GeV ([Dlamini et al., 2021](#)). The reach shown for Hall A are approximate areas around their reported bin centers.

Overview of This Analysis

This work details the analysis of data taken at the JLab CLAS12 experiment to measure the deeply virtual neutral pion electroproduction cross-section .

Chapter 2 describes the experimental setup. Chapter 3 discusses the computational and simulational infrastructure built and used as an integral part in estimating correction factors and performing an accurate measurement. Chapter 4 explains the specific analysis procedures and estimation methods used to arrive at cross-section values. Chapter ?? presents further physics analysis, made possible with the extracted cross-section values. Fig. 1-17 broadly summarizes the analysis flow; elec-

tronic readers can conveniently click on boxes for hyperlinks to relevant sections.

statement introducing idea of posing this as an inverse problem, introduce the experimental interpretation of the cross section




Figure 1-17: Analysis Overview Flowchart

Chapter 2

Experiment and Data Processing

The CLAS12 detector is a large angle spectrometer that generally covers angles from 5 to 130 degrees, spanned by two main detector subsystems - the Forward Detector and the Central Detector.

Overview of Jefferson Lab Why is CLAS12 particularly suited for this measurement?

2.1 Accelerator and Beamline

CLAS12 acceptances and resolutions are also superior to that of CLAS6. Main differences are:

- RGK has outbending torus vs inbending CLAS6 data
- the distance between the target and the PCal has increased, the FTCal extends to lower angles, and the gap between FTCal and PCal is much smaller than between IC and EC
- proton polar angle was limited to 60 deg in the e1dvcs dataset if my memory is correct

volker clas12 exp [V. Burkert et al., 2020](#)

in beam dump area, link to own fraday cup paper [Johnston et al., 2019](#)




Figure 2-1: CLAS12 Detector System

Capability	Quantity	Status
Coverage & Efficiency	Tracks (FD) Tracks (CD)	$5^\circ < \theta < 35^\circ$ $35^\circ < \theta < 125^\circ$
	Momentum (FD & CD)	$p > 0.2 \text{ GeV}$
	Photon angle (FD) Photon angle (FT)	$5^\circ < \theta < 35^\circ$ $2.5^\circ < \theta < 4.5^\circ$
	Electron detection (HTCC) Efficiency	$5^\circ < \theta < 35^\circ, 0^\circ < \phi < 360^\circ$ $\eta > 99\%$
	Neutron detection (FD) Efficiency	$5^\circ < \theta < 35^\circ$ $\leq 75\%$
	Neutron detection (CD) Efficiency	$35^\circ < \theta < 125^\circ$ 10%
	Neutron Detection (BAND) Efficiency	$155^\circ < \theta < 175^\circ$ 35%
Resolution	Momentum (FD)	$\sigma_p/p = 0.5 - 1.5\%$
	Momentum (CD)	$\sigma_p/p < 5\%$
	Pol. angles (FD)	$\sigma_\theta = 1 - 2 \text{ mrad}$
	Pol. angles (CD)	$\sigma_\theta = 10 - 20 \text{ mrad}$
	Azim. angles (FD)	$\sigma_\phi < 1 \text{ mrad}/\sin \phi$
	Azim. angles (CD)	$\sigma_\phi < 1 \text{ mrad}$
Operation	Timing (FD)	$\sigma_T = 60 - 110 \text{ ps}$
	Timing (CD)	$\sigma_T = 80 - 100 \text{ ps}$
	Energy (σ_E/E) (FD)	$0.1/\sqrt{E} \text{ (GeV)}$
	Energy (σ_E/E) (FT)	$0.03/\sqrt{E} \text{ (GeV)}$
DAQ	Luminosity	$L = 10^{35} \text{ cm}^{-2}\text{s}^{-1}$
Magnetic Field	Data Rate	20 kHz, 800 MB/s., L.T. 95%
	Solenoid	$B_0 = 5 \text{ T}$
	Torus	$\int B dl = 0.5 - 2.7 \text{ Tm at } 5^\circ < \theta < 25^\circ$

Figure 2-2: CLAS12 Specification

	Front End DAQ Rate	Event Size	L1 Trigger Rate	Bandwidth to mass Storage	
JLab	GlueX	3 GB/s	15 kB	200 kHz	300 MB/s
LHC	CLAS12	0.1 GB/s	20 kB	10 kHz	100 MB/s
LHC	ALICE	500 GB/s	2,500 kB	200 kHz	200 MB/s
LHC	ATLAS	113 GB/s	1,500 kB	75 kHz	300 MB/s
LHC	CMS	200 GB/s	1,000 kB	100 kHz	100 MB/s
BNL	LHCb	40 GB/s	40 kB	1000 kHz	100 MB/s
BNL	STAR	50 GB/s	1,000 kB	0.6 kHz	450 MB/s
BNL	PHENIX	0.9 GB/s	~60 kB	~ 15 kHz	450 MB/s

* Jeff Landgraf Private Comm. 2/11/2010
 ** CHEP2006 talk Martin L. Purschke, current capability is 800MB/s peak, 500MB/s sustained (priv. comm. 2/14/2010)
 CHeP2007 talk Sylvin Chapelin

June 3, 2014

The GlueX Detector in Hall-D - David Lawrence - JLab

31

Figure 2-3: CLAS12 Data Rates, Compared to Other Experiments

2.1.1 Thomas Jefferson National Accelerator Facility

2.1.1.1 Electron Source

2.1.1.2 Polarimeters

As stated, for RGA, the fact that the beam is polarized is not useful, but it is true and is measured by Moller Polarimeters.

Polarimetry: Good for beam energies between 100 MeV and 50 GeV. Polarized beam electrons are scattered from other polarized electrons in a target, usually magnetized foils. Only a small fraction of all the target electrons are polarized, so this method has a small analyzing power. Analyzing power is exactly calculable in QED. At high beam energies, analyzing power and scattering probability both become independently of beam energy. Maximum analyzing power is about 80versely polarized target can be used to measure transverse beam polarization, but analyzing

ing power is only about 10tron at half beam energy, so magnets are used to bend these electrons out to detectors. These detectors can be, for example lead glass total absorption cherenkov counters. Since the two electrons are corellated, can use things like time coincidence to reduce background, although for low duty factor accelerators only one electron is required as statistics would otherwise be too low. A main background to this process is Mott scattering with the electron radiating off energy after scattering, appearing as a Moller electron

The scattering target is either iron or vanadium permendur (iron-cobalt alloy). Only 2 of 26 electrons in iron have their spins oriented, leading to a total analyzing power of only 6 percent and transverse analyzing power of only 1 actually are corresponds to an uncertainty in analyzing power. There are 'easy' and 'hard' magnetization schemes - easy does a soft magnetization, while hard uses a several tesla magnet to saturate the target. In principle, uncertainties on magnetization in the hard scheme can be removed by using the Kerr magneto-optic effect, but this has not ever been implemented. An important correction is due to the Levchuk effect, where due to momentum differences between electrons in different shells, electrons scattered off of polarized electrons are more likely to be detected than off of unpolarized electrons. Specifically, inner electrons are unpolarized and have a large average momenta, so when struck they can fall outside the 113 TOC acceptance of the Moller detectors, while the outer electrons, which are polarized, have a small average momentum, and behave as expected. This is up to a 15 measurements, and is currently a work in progress.

2.1.1.3 Accelerator

For entry into CLAS12, the beamline specs are as follows:

Beam current: up to 50 nA

Beam energy spread: 10^{-4}

Beam size: Less than 0.4 mm

Beam stability: Less than 0.1 mm

Beam halo: 10^{-4}

Beam polarization: up to 80%

2.1.2 CLAS12 Beamline

2.1.2.1 Liquid Hydrogen Target

Rasterization of some kind

The hydrogen target in RGA is cooled to 20 K using a He4 evaporation fridge.

Can be polarized by dynamic nuclear polarization, driven by a 140 GHz microwave source, can reach 90% polarization for protons, 40% for deuterons (both longitudinally polarized). The polarization can be measured by a Q-meter based NMR. 2.5 cm diameter target, extended 5 cm long.

RGA does not use a polarized target. The beam is polarized, but the target is not, so polarization is not helpful for extracting the 5-fold differential cross section (but it would be if the target was also polarized, and is useful for BSA measurements).

2.1.2.2 Faraday Cup and Beam Dump

Luminosity in CLAS12 is measured from the Faraday Cup and using reference reactions such as elastic scattering. We don't use the Faraday Cup event by event, but we do use it run by run. For beam current measurements, beam position monitors upstream are used - but this is for monitoring on-line, not for analysis. Can manage 175 Watts - 17 nA at 10 GeV. Is used to calibrate beam current, needs a blocker in at higher currents.

2.2 CLAS Detectors and Run Conditions

2.2.1 CLAS Detector System

2.2.1.1 Forward Detector

Overview:

2.2.1.2 HTCC

Used basically as an electron trigger. Composed of 60 lightweight ellipsoidal mirrors, that focus Cherenkov light onto eight 5-inch phototubes (48 channels on entire HTCC). Working gas is CO₂ at STP ($n=1.0005$, $\theta_{max} = 1.7$ degrees, covers 5 to 35 degrees, 2π in azimuth. Active area 2.4 meters in diameter. Electron signal threshold is 15 MeV, charged pion threshold is 5 GeV. 99.9% electron detection efficiency vs. pions. 15 feet in diameter, 6 feet long. Mirror thickness is 0.1 g/cm². Kaons have no signal, as they would need 16 GeV to generate a signal. Uses Winston cones to increase collection efficiency. 20 photoelectrons per electron in HTCC, 25% quantum efficiency. The PMTs have 14 dynodes, gain of about 10⁷. HTCC material budget

0.135 g/cm²

Reconstruction Software Status

Reconstruction software:

All major software development completed, or in the final stage (open pull requests)

Extensive validation from RG-A and RG-B

Pass2 vs. Pass1:

Large increase in charged particle reconstruction efficiency (10% per track or more)

Resolution improvements due to alignment (both FD and CD) and removal of tracking biases and energy loss correction (CD)

Improvements to neutral reconstruction in ECAL (handling of overlapping clusters) and FT (improved calibrations)

Several updates to EB to improve track-hit matching and provide more information for analyses

Alignment of trackers:

CVT alignment based on Kalman Filter Alignment algorithm (see S. Paul *et al.* (CLAS), arXiv:2208.05054, submitted to NIM)

New implementation of DC alignment

DC-CVT alignment based on beam spot method

For more details:

Offline software status and plan: Raffaella De Vita – Tuesday

New CVT tracking finding: Véronique Ziegler – Tuesday

CALCOM status update and pass-2 preparations: Daniel Carman

Update on DC calibrations: Florian Hauenstein – Thursday

DC alignment: Trevor Reed – Thursday

RG-B Spring 2019

AI Conventional

Count


$M_X(ep \rightarrow e^+\pi^+\pi^-X)$ (GeV)

p (GeV)

CD PID after energy loss, showing pion, proton and deuteron bands

[Raffaella De Vita] [Daniel Carman]

Nov 2o22
Patrick Achenbach



2.2.1.3 Torus

Outbending allows for lower Q2 measurements, inbending allows for slightly higher Q2 measurements.

6 coil torus, 4k amps, 3.5 Tesla torodial field, supercritical LHe cooled. 14.2 Megajoules stored energy. 2 Henries of inductance. Field strongest at small angles, weakest at large angles. Inbending vs out bending: I have been wondering about this as well. All I know is that inbending and outbending have different acceptances. So, I guess some channels prefers inbending while the others do outbending? I'm not sure though. FX claims outbending results have better quality for these days.




Figure 6: Combined solenoid and torus magnetic fields. The color code shows the total magnetic field of both the solenoid and torus at full current. The open boxes indicate the locations and dimensions of the active detector elements.

2.2.1.4 Drift Chambers


There are 3 layers of drift chambers, each with 6 sections. Each chamber has 2 superlayers of 6 layers by 112 wires, for a total of 24,192 wires. (Structure is 112 wires * 6 layers * 2 superlayers * 18 DC sections = 24,192 wires). Physical wire sectioning looks like:

(IIIIII)-(IIIIII)—(IIIIII)-(IIIIII)—(IIIIII)-(IIIIII) x 6 sectors

Where each "I" is a layer of 112 wires.

Spatial resolution is $300 \mu\text{ m}$, angular coverage 5-40 degrees. Momentum resolution $\Delta p/p < 1\%$, angular resolution is 1 mrad in theta, $1\text{mrad}/\sin\theta$ for phi. The Drift Chambers are located 2, 3, and 4 meters from the gas mixture is 90/10 Argon/CO₂. Time resolution = ?

DC specifics: 30 micron diameter tungsten sense wires, 80 micron Cu-Be field wires, 140 micron Cu-Be guard wires. 20 g tension on sense, 62 g tension of field, 180 g on guard. Max sag calculated to be on order of 10 microns.



55

TOC

Flow: 1-17

2.2.1.5 LTCC

6 sectors, perfluorobutane (C_4F_{10}) $\rightarrow n = 1.0013 \rightarrow \theta_{max} = 3$ degrees. Electron threshold 9 MeV, pion threshold 2.7 GeV, Kaon threshold 9.4 GeV. Allows for good pion/kaon discrimination from 3.5 GeV to 9 GeV.

Each section has 108 mirrors, 36 winston cones, and 36 PMTs. Mirror is aluminium with MgF_2 coating. Kevlar support structure. Perflourobutane is 100% transparent above 220 nm light.



Figure 2-4: Low Threshold Chrenkov Counter

2.2.1.6 RICH

Provides PID in the range of 3-8 GeV, replacing one sector of LTCC (right middle sector). Pion/Kaon rejection factor > 500 , Kaon/Proton rejection factor > 100 . Covers 5 to 25 degrees in theta, uses aerogel ($n=1.05 \rightarrow \theta_{max} = 18$ degrees). Pion

threshold 460 MeV, Kaon threshold 1.6 GeV. Read out by 64 channel photomultipliers. $\beta_{min}=0.95$, $\gamma_{min} = 3.3$

What is angular resolution?

Reminder of relevant equation:

$$\cos \theta = \frac{1}{\beta n} \longrightarrow \theta = \arccos \frac{1}{\beta n} \quad (2.1)$$

$$\gamma = \frac{1}{\sqrt{1 - \beta^2}} \longrightarrow \beta = 1 - \frac{1}{1 - \gamma^2} \quad (2.2)$$

	momentum	$\sim \gamma$	θ
pion	3	21	17.4
kaon	3	6	12
pion	5	36	17.6
kaon	5	10.1	15.9
pion	8	57.1	17.7
kaon	8	16	17.1

Detector	Scintillator	PMT
FTOF - 1a	BC-408	Phillips XP2262, EMI 9954A
FTOF - 1b	BC-404	Hama. R9779
FTOF - 2	BC-408	EMI 4312KB
PCAL	FNAL	Hama. R6095
ECAL	BC-412	Philips XP2262, EMI 9954
CND	EJ-200	Hama. R10533
CTOF	BC-408	Hama. R2083
HTCC	N/A	ET 9823QKB
LTCC	N/A	200 Photonis XP 4500B
LTCC	N/A	16 Photonis XP 4508 (Quartz Window)

*ET stands for Electron Tube, a company. Could not find a spec sheet for this PMT type. ** Could not find spec sheets for either HTCC, or LTCC PMTs.

Scintillator	Detectors	Principal Use/Features	L.O.	WME	R/D Time	L.A. Length
BC-404	FTOF-1b	Fast Counting	68	408	0.7/1.8	140
BC-408	FTOF-1b,2,CTOF	TOF - Large Area	64	425	0.9/2.1	210
BC-412	ECAL	Large Area	60	434	1.0/3/3	210
EJ-200	CND	Long attenuation, fast	64	425	0.9/2.1	380

L.O - Light Output - % Anthracene WME - Wavelength Maximum of Emitted Photons R/D Time - Rise / Decay time (ns) L.A. Length - light attenuation length (cm)

All scintillators have a PVT (Polyvinyltoluene) base.

EJ-200: 200 – 10K photons per 1 MeV.

Thermal effects: EJ-200 loses 5% of its light output between 20 degrees C and 60 degrees C. No change between -60 to 20 degrees C.

PMT	Det.	TS/A	WVE	PHTC	DNY	Anode	Time Resp.
Hama. R6095	PCAL	28/25	300/420/650	BA/BSG	B&L/11/2.1	1500/0.1	4/30/3
Hama. R9779	FTOF-1b	51/46	300/420/650	BA/BSG	LF/8/0.5	1750 /0.1	1.8/20/0.25
Hama. R10533	CND	51/46	300/420/650	BA/BSG	LF/10/4.2	1000/0.1	2/24
Hama. R2083	CTOF	51/46	300/420/650	BA BSG	LF/8/2.5	3000/0.2	0.7/16/
Phillips XP2262	FT1a ECAL						
EMI 9954A	FT1a ECAL						
EMI 4312KB	FTOF-2						

No spec sheets could be found for the PMTs used in the TFOT1a, ECAL, or FTOF-2. Typical dark currents for all PMTs are 100 nA.

Tube size / photocathode area (diameter in mm) Wavelength short / peak / long (nm) Photocathode / window material (BA = Bialkali, BSG = Borosilicate Glass) Dynode structure / stages / gain LF = Linear-focused, B&L = Box and line / / Gain - Gain $\times 10^6$ Anode to Cathode Voltage / Anode Current - Volts / mA Rise / transit time/time spread in ns

[R10533 PMT](#) [R2083 PMT](#) [R9779 PMT](#) [R6095 PMT](#)

[CND Scintillator](#)

[BC Scint Specs from Saint Gobain](#)

[CND PMT](#)

[CTOF PMT Spec sheet](#) Bialkali photocathode, 8 dynodes, 2.5×10^6 typical gain Linear-focused dynode structure Window material Borosilicate glass peak wavelength 420 nm, range from 300 to 650 nm. Max anode to cathode voltage of 3500 V, made anode current of 0.2 mA. Dark current around 100 nA.

2.2.1.7 FTOF

Used for PID, three layer system - 1a, 1b, and 2. Has a design resolution of 60 ps to 160 ps. Average scintillation rate 250 kHz. Pion/Kaon separation up to 2.8 GeV, Kaon Proton separation up to 4.8 GeV, pion proton separation up to 5.4 GeV.


6 meters away from target. Time resolution of 80 ps less than 36 degrees, 150 ps greater than 36 degrees. PMTs are shielded from CLAS12 torus. 6 sectors, plastic scintillator, double sided PMT readout. 3 panels - 1a - 23 counters, 1b - 62 counters, 2 - 5 counters.


15cm wide x 5 cm deep x 33 cm up to 376 cm long.

20-30 cm up to 15x5 130 ps

350-400 cm 6x6 60 ps

370 to 430 cm, 22x5cm, 150 ps





The timing resolution minimums are for being close to the beam axis where particles are moving faster, and farther out from the beamline (larger theta) particles are moving slower so a less resolved time difference is acceptable.

2.2.1.7.1 1a Coverage is 50% at 5 degrees to 85% at 35 degrees

Dimensions: L 32.3 cm to 376.1 cm, wxh = 15x5 cm

Material BC-408

PMTS: EMI 9954A, Phillips XP2262

Time resolution 90 - 160 ps small bar to big bars

2.2.1.7.2 1b Coverage is 50% at 5 degrees to 85% at 35 degrees

Dimensions: L 17.3 cm to 407.9 cm, wxh = 6x6 cm

Material BC-404 (first half) and BC-408

PMTS: Hamamatsu R9779

Time resolution 60-110 ps small bar to big bars

2.2.1.7.3 2 Coverage is 85% at 35 degrees to 90% at 45 degrees

Dimensions: L 371.3 cm to 426.2 cm, wxh = 22x5 cm

Material BC-408

PMTS: EEMI 4312KB

Time resolution 140 - 165 ps small bar to big bars

For more FTOF specifications, look [here](#)

FTOF two panels: Official answer from CLAS12 FTOF NIM paper: "For tracks that pass through both arrays the combined time information (described in Ref. [10]) is used and results in a 20

I guess this is the right answer though: 1a is recycled one from CLAS while 1b is new one.

2.2.1.8 PCal and ECal

ECal from CLAs could only contain showers with $E < 5$ GeV. Above 5.5 GeV, couldn't resolve neutral pion gamma gamma angle, so needed PCAL. PCAL is 7 meters from target, ECAL is 7.5 M from target. EC segmentation 10 cm, PCAL finer segmentation. PCal 5.5 radiation lengths. 20.5 radiation lengths total. Both are sampling calorimeters, with PB and scintillator layers. The CLAS ECAL was resused and a new PCAL was installed in front of it. Primarily used for identification of electrons, photons, gamma gamma decays from pions, and neutrons. They are sampling calorimeters with six moduels. Each module has a triangular shape with 54 (15/15/24 - PCAL/ECALinner/ECALouter) layers of 1 cm htick scintilla-

tors segmented into 4.5/10 cm (PCAL/ECAL) wide strips and sandwiched between 2.2 mm thick lead sheets. The total thickness is about 20.5 radiation lengths.


Scintillator layers are grouped into three readout views with 5/5/8 PCAL/ECinner/ECouter, layers per view providing several cm resolution of energy clusters. Light from each scintillator readout group is routed to PMTs via flexible optical fibers.


Overall performance:

Energy resolution of 10%, position resolution of 2 cm, time resolution of 500 ps.

Are these the real statistics? Because they seem like BS.

BC-408






2.2.1.9 PCAL

50% coverage at 5 degrees, 85% coverage at 35 degrees. 15 scintillators, 14 lead layers, per module. 1200 scintillator strips, $1 \times 4.5 \text{ cm}^2$ up to 432 cm long, with two holes along the strip, and 0.25 mm TiO₂ coating (reflective coating)


Lead sheets are 2.2 mm thick. Readout by fibers into 1 inch PMTs, Hamamatsu

R6095. Light yield is 11-12 photo-electrons per MeV.

PCAL scintillator was manufactured at the FNAL-NICADD Extrusion Line Facility. Polystyrene base was Dow STYRON 663 W, primary dopant is 2,5-diphenyloxazole (PPO, 1% by weight) - this is the organic scintillator, peaks at 385 nm:



The Secondary dopant is 1,4 bis (5-phenyloxazol- 2-yl) benzene (POPOP, 0.03% by weight) - also scintillator, peaks at 410 nm.



A reflective surface coating of polystyrene with 12% TiO₂ with 0.25 mm nominal thickness was co-extruded.

Cast plastic scintillator costs about \$50 per kg, while extruded scintillator is significantly lower in price - about \$10 per kg.

[Interesting write up on FNAL Scintillator extrusion](#)

[PCal Technical Report](#) [ECal Technical Report](#)

2.2.1.10 ECAL

50% coverage at 5 degrees, 85% coverage at 35 degrees. 39/38 scintillators / lead layers per module. 216 readout channels per module, 1200 strips per module. Strips are 1x10to12cm² by up to 441 cm long, BC-412 (plastic scintillator with high light output, longest light attenuation length, [cheap!](#)). Lead sheets are 2.4 mm thick. Read out by fiber into 2 inch PMTs, Phillips XP2262 and EMI 9954. 3-4 photoelectrons/MeV deposited energy.

2.2.1.11 Central Detector

Overview: The Central Detector spans roughly 35 to 125 degrees, and contains 4 sub-detectors, all in a 5 Tesla solenoidal field. The 5 detectors are: SVT, MMVT, CTOF, and CND. Low t-data are very important for the meson exclusive physics. The GPD interpretation works only in the region $-t/Q^2 < 1$. From this point of view the central detector will not only increase the total statistics by a factor more than 2 but will add the valuable data with low t.

2.2.1.12 SVT

The Silicon Vertex Tracker (SVT) covers from 35 to 125 degrees in θ . Has 8 layers (4 concentric rings) with 10, 14, 18, and 24 sectors respectively, double sided. 2π angular coverage. Read out with ASICs- FSSR2s. Designed to operate at 10^{35} luminosity,

momentum resolution of $\sim 5\%$ for 1 GeV particles with $\theta = 90$ degrees. 42 cm long, 4 cm wide, 0.4 cm thick. Spatial resolution of $50 \mu\text{m}$, momentum resolution $\sim 5\%$, theta resolution 10 mrad, phi resolution 5 mrad. 33,792 total readout channels. Sensor thickness is $320 \mu\text{m}$, readout pitch $156 \mu\text{m}$. Supported by rohacell and carbon fiber backing to reduce material budget, at $\sim 1\%$ of a radiation length.





Figure 2-5: Silicon Vertex Tracker



2.2.1.13 MMVT

Composed of two parts: a **Barrel Tracker** and a **Forward Tracker**. PCB is 200 μm thick, 0.3% of a radiation length. 20 MHz sampling frequency. Time resolution of 10 ns. 500 μm strip pitch.

Advantages of MMVT for CLAS12 :

Price: much cheaper compared to SVT. For large area, the price become rapidly prohibitive. Material: Since it is a gasesous detector, it is good for the material budget. Physics Requirements: Not as good spatial resolution as SVT, but can resolve polar angle better. Optimal performance is actually achieved with a combination of both detectors are used.


Overall momentum uncertainty (σ_p/p) = 1.6%. $\sigma_\theta = 1.4$ mrad. $\sigma_\phi = 2.6$ mrad. $\sigma_z = 270$ mm.

2.2.1.14 Barrel Tracker

18 cylindrical detectors arranged in 6 layers. Covers 35 to 125 degrees. 15,000 readout elements. Gas Mixture 90% Argon, 10% isobutane. 3 mm drift gap. 5 kV/cm field. 75% mesh transparency.

2.2.1.15 Forward Tracker


6 circular, flat detectors from 6 to 29 degrees in θ . Improves vertex resolution by a factor of up to 10x compared to just the drift chambers along. 6,000 readout elements. 80% neon, 10% ethane, 10% Carbon Tetrafluoride. 5 mm drift gap. 1kV/cm field. 100% mesh transparency.



2.2.1.16 CTOF


Central for PID purposes. Divides into 48 1 meter long plastic scintillators with double sided PMT readout. PMTs are in the 0.1 T fringe field region and enclosed

in magnetic shielding. 65 picosecond timing resolution. 35 to 125 degrees, 2π in polar angle. 3 cm x 3 cm scintillator planks. Pion/Kaon separation up to 0.64 GeV, Kaon/proton separation up to 1 GeV, pion proton separation up to 1.25 GeV.



2.2.1.17 CND

Detects 0.2-1 GeV neutrons. 3 layers, 48 paddles per layer. Plastic scintillator, 3 cm x 3cm, 0.7 meters long. Neutron detection efficiency $\sim 10\%$. 130 picosecond timing resolution, 2 degrees angular resolution (polar and azimuth).



2.2.1.18 Solenoid

5 Tesla super conducting magnet, uniform field ($\Delta B/B = 10^{-4}$). Weakest at small angles, strongest at large angles. Opening polar angle of 40 degrees. Momentum range of interest 0.3 to 1.3 GeV. 18 Megajoules stored energy. 85 cm in diameter, 4.2 Kelvin operation.

2.2.1.19 Forward Tagger

2.2.1.20 BAND

BAND not impportant.

2.2.2 Run Conditions

CLAS12 runs with "open trigger", which means different sub-experiments can define their own triggering logic. There is a standard electron trigger, based off of hits in HTCC, ECal, and FTOF.

Only about 50% of the electron triggers recorded with an inbending torus polarity are actually electrons. For outbending torus polarity, the electron trigger purity is as high at 70%.

Detector Specs

20 kHz Level 1 trigger rate, 1 GB/s.




Figure 2-6





Figure 2-7

Target Assembly

Side view



Top view




Figure 2-8

Data taken is RGA taken in Fall 2018

Mention configurations and combinatorics

2.3 Reconstruction and Particle Identification

here we talk about CLAS PID

2.3.1 Decoding and Track Reconstruction

2.3.2 Particle Identification

**photon cuts: pid 22, status > 2000 (in FD or CD, not ftagger) momentum greater than 400 MeV each

**proton cuts: pid 2212

**electron cuts: pid==1 and status < 0(negative particle)

For this analysis all final state particles should be detected. After π^0 decay we are going to have 4 particles: electron, proton and two photons. The particle identification methods are applied to select the exclusive event with at least one electron, proton and two photons.

2.3.2.1 Electron

2.3.2.2 Proton

2.3.2.3 Photon

2.3.2.4 Pion

Basic event builder cuts are utilized, then additional cuts are made that are common with the RGA Analysis note ([overleaf link](#) and developed by Sangbaek Lee (sang-

baek@mit.edu - [github code here](#). For this analysis, both the central detector and forward detector are utilized for proton tracking. The forward tagger is also utilized for photon identification.

2.3.2.5 Neutral pion

In addition to individual particle PID procedures the cut on the mass of two photons is applied:

- $0.07 < M_{\gamma\gamma} < 0.2 \text{ GeV}$

The pion is more thoroughly constrained by the exclusivity cuts, described in the next section.

2.3.3 Data Storage and Formatting

2.3.3.1 Data Location and Availability

2.3.3.2 File Formatting and Conversion

Mention that same transformations are used for rec events and filtering




Figure 2-9: The distribution for mass of two photons $M_{\gamma\gamma}$




Figure 2-10: The distribution for mass of two photons after exclusivity cuts

Chapter 3

Simulations

3.1 Computing and Simulation Infrastructure

Motivation for massive simulations, inverse problem scope Discuss scale of data - how many simulations needed, how much data is produced, how long it takes to run simulations, cost and energy involved. overview: event generation - physics simulation - reconstruction - other

3.1.1 MIT Tier 2 and Supercomputing Clusters

OSG MIT Tier 2 etc

3.1.2 Overview of Simulation Runs

Simulation hours: As for your question, if there is constant pressure then yes, MIT baseline would be constant.

However: need much more than 20K jobs / day to be at constant pressure. The dedicated resources + high priority can go up to 20K cores at once and if a job last

4 hours that's $24/4 = 6*20K = 120K$ jobs / day to be at constant pressure! Please do not submit that many jobs ;-)

You can check the monitoring page for more details. In particular you can select 1 month timeframe, and log scale. You will see holes in queue. And if you select just 'MIT' on the dedicated graph, you'll see the attached picture, where the 'holes' in the pressure are well shown and the number of MIT cores under pressure is about 3K.

Simulations are necessary in order to extract correction factors.

GEMC was used to process generated events through the CLAS12 fall 2018 RG-A configuration. Specifically, a generator based off the GK model and CLAS6 data - aao_norad¹.

3.2 Event Generation

Maid model discussion Xiaqing said that the following: [Dreschsel and Tiator, 1992](#) contains the formalism for the MAID model

You don't need much details about generator, it is based on GK model with Valery's fit to CLAS6 data.

NON RADIATED, INBENDING // Pi0 leptoproduction in Goloskokov-Kroll (GK) model. The code is currently being tested and implemented in PARTONS framework with additional features. If you plan to use this work in a publication, please use and reference the most recent version of PARTONS in <http://partons.cea.fr>

Andrey Kim and Nick Markov have the pi0 generator. It has my parametrization for $W > 2$ GeV and MAID for $W < 1.7$ GeV.

¹https://github.com/drewkenjo/aao_norad

My model will for sure work for 12 GeV. It actually very close even for the COMPASS pi0 data (180 GeV muon beam).

There is reasonable coincidence between my model and MAID in the point W=1.7 GeV, not ideal but good enough for the MC.

I think actually that my parametrization has to work in the region W<2 GeV but I am not sure that MAID is doing good job due to the absence the experimental data at W 1.7 GeV.

Hi Bobby, Please take a look at README: [norad](#) It has instructions how to compile, run and configure the program. Please don't hesitate to ask questions!

For event generator we have aao_rad can generate radiatied pi0 events in resonance region use 2007 model Put parameterization from valerly's paper, can cover up to whatever Q2 range covered in the paper, beyond that we put some general Q2 behaviour For Exclurad we have similar model, in end may have to iterate a few times to improve the model Exlclurad specifically for resonance region, theoretically should be correct, input probably needs to be updated, can put Valery's new parameterization to cover higher range. Should not be a real issue to implement it because same thing was done for AAORad. High q2 cannot be covered because parameterization only goes to CLAS6 range FX: the cruicail thing is to fold in the radiative corrections with acceptance and efficiens. Best mothod is to use fast monte carlo

aao_rad and aao_norad are event generators for exclusive pi0 and pi+ channels with/without radiative effects. They are written in Fortran. The program was initially developed by Volker Burkert long time ago for the resonance region, then has been evolved for many years and recently extended to DIS region even though lots of things need to be done. Try this to see whether it works.

Amplitudes And Observables (AAO)

3.2.1 Nonradiative Generator

Include generator plots, specifics of layout

3.2.2 Radiative Generator

Include generator plots, specifics of layout, plots showing W cut offs, etc

3.3 Simulation Pipeline

3.3.1 Simulation Processing and JLab Computing

volatile work To cancel all jobs: scancel -u robertej

To view all jobs: squeue -u robertej

3.3.2 Geant4, GEMC, and CLAS Submissions

talk about own work on portals returns to vol, filtering and converting as covered in chapter 2

3.4 Simulation Enhancements with Normalizing Flows

3.4.1 Inverse Transforms and Autoregressive Flows

Introduce theory of MAFs

3.4.2 Exploration of Simulation Speedup with UNMAF

Discuss actual work performed

Chapter 4

Cross Section Measurement

Official Repo: <https://github.com/robertej19/clas12DVPiP>

For each kinematic bin the differential cross section can be written as:

$$\sigma = \frac{N_{meas}}{L\epsilon} \frac{1}{\delta} \quad (4.1)$$

Where $\frac{N_{meas}}{L}$ is the number of events from experiment normalized by the integrated luminosity before acceptance and radiative corrections. $\epsilon = \frac{N_{rec}^{RAD}}{N_{gen}^{RAD}}$ is the acceptance correction and δ is the radiative correction.

δ can be obtained by using the following:

$$\delta = \frac{N_{gen}^{RAD}}{N_{gen}^{NORAD}} \quad (4.2)$$

δ and ϵ need to be properly calculated, but for a first pass we will ignore them so we have just

4.1 General Analysis Overview

4.1.1 Part 1

Cross-sections are theoretically interpreted as the probability for a specific interaction to occur. They can be experimentally estimated by measuring the occurrence frequency relative to the total possible interaction opportunities. In general, the cross-section σ can be expressed as (4.3) the number of measured events of interest N_{meas} divided by the number of total interaction opportunities \mathcal{L} . \mathcal{L} is known as luminosity and is a product of only experimental parameters, such as the number of particles present and the experiment duration.

4.1.2 part 2

$$\sigma = \frac{N_{exp}}{\mathcal{L}} \quad (4.3)$$

Measuring the complete cross-section at once is not feasible, so instead estimates are made of the differential cross-section (4.4), which instead evaluates the probability for a specific interaction to occur in a differential region of phase space, $\frac{d\sigma}{d\Omega}$. Infinitesimal measurements are not possible, so events are counted over some small discretized generalized volume $\Delta\Omega$.

$$\frac{d\sigma}{d\Omega} = \frac{N_{exp}}{\mathcal{L}\Delta\Omega} \quad (4.4)$$

In practice, a number of correction terms need to be included to account for differences between experiment and theory. These correction terms, combined with the specifics of this analysis, yield the full experimental expression of the cross-section (4.5).

$$\frac{d^4\sigma_{ep \rightarrow ep' \pi^0}}{dQ^2 dx_B dt d\phi_\pi} = \frac{N(Q^2, x_B, t, \phi_\pi)}{\mathcal{L}_{int} \Delta Q^2 \Delta x_B \Delta t \Delta \phi_\pi} \frac{1}{\epsilon_{acc} \delta_{RC} \delta_{Norm} Br(\pi^0 \rightarrow \gamma\gamma)} \quad (4.5)$$

The terms on the right-hand side of this equation are:

- $N(Q^2, x_B, t, \phi_\pi)$ - Number of events recorded in a given Q^2, x_B, t, ϕ_π bin.
- \mathcal{L}_{int} - Integrated luminosity
- $\Delta Q^2 \Delta x_B \Delta t \Delta \phi_\pi$ - These are the bin sizes or intervals for the variables Q^2, x_B, t , and ϕ_π .
- ϵ_{acc} - Acceptance correction, which is a combination of detector efficiency and geometrical acceptance, determined through simulations.
- δ_{RC} - Radiative correction factor
- δ_{Norm} - Overall normalization factor
- $Br(\pi^0 \rightarrow \gamma\gamma)$ - Branching ratio of the decay of a neutral pion (π^0) into two photons ($\gamma\gamma$), which is most recently measured at 98.8131% [Husek, Goudzovski, and Kampf, 2019](#)




Figure 4-1: High-level data processing flow

4.2 Data Pre-Processing

4.2.1 Experimental Data Pre-Processing

4.2.1.1 Momentum Corrections

Mom Corr

4.2.1.2 Energy Loss Corrections

4.2.2 Simulation Data Pre-Processing

4.2.2.1 Simulation:Experiment Resolution Matching

The standard aao simulations result in missing mass distributions that are too optimistic compared to experimental data. Observe the discrepancies between simulated and experimental distributions in figure 4-2.




Figure 4-2: Comparison of experiment (blue) and simulation (red) missing mass, energy, momentum, and invariant gamma-gamma mass distributions, before any smearing factors were added to the simulation data.

To improve the matching between simulation and experiment, gaussian smearing factors were added after reconstruction to the simulated dataset. These factors were tuned by Sangbaek Lee to have optimal matching across all missing mass spectra combinations (figure 4-3). Once these factors were determined, the simulations were used to extract an acceptance correction.




Figure 4-3: Comparison of experiment (blue) and simulation (red) missing mass, energy, momentum, and invariant gamma-gamma mass distributions, with smearing factors added to the simulation data proton and photon momenta.

4.3 Event Selection

Overview words about event selection

4.3.1 Exclusivity Cuts

After the selection of events with at least one electron, proton and two photons, it is time to take a look at the exclusive distributions. The Fig. 4-4 shows 2D distribution of MM^2 (epX) vs $\theta_{X\pi}$, where MM^2 (epX) is a missing mass squared of (epX) system and should have a peak near 0.0182 GeV^2 , and $\theta_{X\pi}$ is an angle

between expected and reconstructed pion. The bright spot on the figure corresponds to the exclusive $ep \rightarrow ep\pi^0$ events. In order to reduce the background exclusivity cuts need to be developed based on the conservation of energy and momentum. The relevant 1D exclusive distributions are shown on the Fig. 4-5 and 4-6.




Figure 4-4: MM^2 (epX) vs $\theta_{X\pi}$ 2D distribution.




Figure 4-5: Exclusive distributions for events with at least one electron, proton and two photons.




Figure 4-6: Exclusive distributions for events with at least one electron, proton and two photons.

4.3.1.1 Tight $M_{\gamma\gamma}$ ass and transverse missing momenta cuts

The first step is to use tighter $\gamma\gamma$ mass cut: $0.096 < M_{\gamma\gamma} < 0.168$ GeV, and take a look at the missing transverse momentum distributions (see Fig. 4-7). From momentum conservation law we expect transverse momentum to be zero, so we can apply cuts on Δp_x and Δp_y to further improve exclusive channel selection. The cuts $|\Delta p_x| < 0.2$ and $|\Delta p_y| < 0.2$ correspond roughly to 4-5 σ .




Figure 4-7: Exclusive distributions for events with at least one electron, proton and two photons.

The exclusive distributions after tight $M_{\gamma\gamma}$ mass and transverse missing momenta

cuts are shown on Fig. 4-8 and display much stronger signal peaks on top of reduced background.




Figure 4-8: Exclusive distributions after tight $M_{\gamma\gamma}$ mass and transverse missing momenta cuts .

4.3.1.2 $\theta_{X\pi}$ ut determination

The cut on angle between expected and reconstructed pion is used in order to further reduce background. To choose the value of the $\theta_{X\pi}$ cut the $MM^2(epX)$ distribution is analyzed at multiple $\theta_{X\pi}$ cut values and fit using gaussian+polynomial function as shown on Fig. 4-9. From the fit we can estimate the number of good exclusive events (gaussian) and the number of background events (polynomial) and their dependence on $\theta_{X\pi}$ cut. Fig. 4-10 and 4-11 show the numbers of signal and background events

as functions of $\theta_{X\pi}$ cut value for multiple bins in Q^2 and x_B . These plots show that the cut $\theta_{X\pi} < 2^\circ$ allows to select the most number of good events with the least background, and relaxing it beyond 2° does not gain us any good exclusive events but increases background.




Figure 4-9: $MM^2(epX)$ distributions for multiple $\theta_{X\pi}$ cut values.




Figure 4-10: The numbers of signal (red markers) and background (black markers) events as functions of $\theta_{X\pi}$ cut value for multiple Q^2 bins.




Figure 4-11: The numbers of signal (red markers) and background (black markers) events as functions of $\theta_{X\pi}$ cut value for multiple x_B bins.

4.3.1.3 Final exclusivity cuts

The list of final exclusive cuts is following:

- $\Delta p_x < 0.2 \text{ GeV}$
- $\Delta p_y < 0.2 \text{ GeV}$
- $\theta_{X\pi} < 2^\circ$
- $0.096 < M_{\gamma\gamma} < 0.168 \text{ GeV}$
- $MM^2(epX) < 0.5 \text{ GeV}^2$

Exclusive distributions after all exclusivity cut except $MM^2(epX) < 0.5 \text{ GeV}$ are shown on Fig. 4-12




Figure 4-12: Exclusive distributions after all exclusivity cuts .

To arrive at a DVEP candidate event, we do the following

Code flow:

Consider a directory with n hipo files. For each hipo file, do the following.

Read each file event by event, and do the following

Check that the event has the proper databanks, and if not, go to teh next event.

Get a list of all the electrons*, protons*, and photons* in the event

*= links to most up to date PID methods

for every electron in the event (always only one, at least in the skims, but not held to be one) do the following For every proton in the event, do the following

Calculate some basic quantities and fill histograms

for every permutation of pairs of photons in the event, do the following

calculate various kinematic quantities, and pass to see if creates a viable pion* and a viable DVEP event*

if so, fill relevant histograms and count as a DVEP event, otherwise skip to next event

viable pion: pion mass between 100 and 180 MeV pion momentum greater than 1.5 GeV angle (theta) between each photon and the electron to be greater than 8 degrees

viable DVEP event: Q^2 greater than 1 W greater than 2 difference between theta of missing 4-momentum and reconstructed pion less than 2 degrees difference between missing X px and py 300 MeV each or less Difference in missing mass squared between pion and X less than 1 GeV ** make sure this is right difference in missing energy and X less than 1 GeV **make sure this is right

4.3.2 Kinematic Fitting

Instead of rigid cuts, we can use maximum likelihood estimators - include notes from Janet's class

4.4 Luminosity

The strategy to calculate the luminosity is as follows:

- For each run, retrieve a measure of how much beam passed through the target, I believe in the case of CLAS12 using the Faraday cup to measure beam charge - sum the beam charge over all runs being considered and include any relevant corrections factors - multiply this by target length, density, etc. to get the integrated luminosity
 - use this value to calculate cross sections.

Compare integrated luminosity of CLAS6 to CLAS12 (in 2011 analysis note)

Implementation: The bank `REC::Event` has an object `beamCharge`, in nanoCoulombs, which is described in the DST as “beam charge integrated from the beginning of the run to the most recent reading of the gated Faraday Cup scaler in `RAW::scaler`, with slope/offset conversion to charge from CCDB. Note, this value will be zero in each file until the first scaler reading in that file.”. This is the (un?)gated beam charge.

This can be accessed via:

According to [this](#) we might need to use tag=1 RAW::scaler::fcupgated instead of REC::Event::beamCharge

The beam charge needs to be converted to integrated luminosity, which can be done as follows:

Luminosity: Events are not necessarily time ordered, need to take largest value minus smallest value

Luminosity is calculated according to equation 4.6

$$\mathcal{L} = \frac{N_A l \rho Q_{FCUP}}{e} \quad (4.6)$$

The terms in equation 4.6 are as tabulated in table 4.4. The accumulated charge on the Faraday cup is calculated by taking difference between the maximum and minimum values of beamQ for each run, and then summing these values. The luminosity determined for the fall 2018 inbending run was 5.5E+40 cm⁻² and the fall 2018 outbending run was 4.65E+40 cm⁻²

Quantity		CLAS12 Value
Avogadro's Number	N_A	6×10^{23}
Electron Charge	e	1.6×10^{-19}
Target Length	l	5 cm
Target Density	ρ	0.07 g/cm ³ (LH2)
Charge on Faraday Cup	Q_{FCUP}	In data

Table 4.1: Terms of Luminosity Equation

We can calculate the luminosity L through the following equation

$$L = \frac{N_A l \rho Q_{FCUP}}{e} \quad (4.7)$$

Where N_A is Avogadro's constant, l is the length of the target, ρ is the density of the target (liquid hydrogen), Q_{FCUP} is the charge collected on the Faraday cup, and e is the charge of the electron. The values of these quantities are (ignoring uncertainties on experimental quantities for the time being):

$$N_A = 6.02214 \times 10^{23}$$

$$l = 5 \text{ cm}$$

$$\rho = 0.07 \text{ g/cm}^3$$

$$e = 1.602 \times 10^{-19} \text{ Coulombs}$$

Q_{FCUP} - this must be measured and obtained from analysis. Typical runs at CLAS12 have an accumulated beam charge of tens to hundreds of thousands of nanoCoulombs.

4.5 Binning

Words about binning

4.6 Correction Factors

4.6.1 Acceptance Correction

4.6.2 Radiative Corrections

4.6.3 Binning Corrections

4.6.4 Overall Normalization Corrections

4.7 Uncertainty Estimation

4.8 Iterative Bayesian Unfolding

4.8.1 Acceptance Correction

4.8.2 Radiative Corrections

4.8.3 Binning Corrections

4.8.4 Overall Normalization Corrections

Goldstein's straightforward results are elucidated in their 2012 work [Gary R Goldstein, J Osvaldo Gonzalez Hernandez, and Liuti, 2012](#).

McAllister provides a comprehensive study on elastic particles in his 1956 publication [McAllister and Hofstadter, 1956](#).

A flexible observables approach by Goldstein was explained in their 2011 research [Gary R. Goldstein, J. Osvaldo Gonzalez Hernandez, and Liuti, 2011](#).

Brodsky addresses light-cone scattering in a 2001 paper [Brodsky, Markus Diehl, and Hwang, 2001](#).

Abt presents their measurement at HERA in a 1993 study [Abt et al., 1993](#).

Derrick's 1995 work discusses measurement data in detail [Derrick et al., 1995](#).

Diehl's 2005 paper provides an in-depth look at protons [M. Diehl and Sapeta, 2005](#).

Patrignani's 2018 paper provides a review of physics [Patrignani and et al. \(Particle Data Group\), 2018](#).

Qiu's 2023 work discusses single distributions [Qiu and Yu, 2023](#).

Zhu's 2015 work explores aspects of microscopy [Zhu and Dürr, 2015](#).

Gockeler's 2007 paper explores transverse simulations [Göckeler et al., 2007](#).

Bibliography

- [1] I. Abt et al. “Measurement of the proton structure function F2 (x, Q2) in the low-x region at HERA”. In: *Nuclear Physics B* 407.3 (Oct. 1993), pp. 515–535. ISSN: 05503213. DOI: [10.1016/0550-3213\(93\)90090-C](https://doi.org/10.1016/0550-3213(93)90090-C) (cit. on p. 104).
- [2] M.G. Alexeev et al. “Measurement of the cross section for hard exclusive π^0 muoproduction on the proton”. In: *Physics Letters B* 805 (June 2020), p. 135454. ISSN: 03702693. DOI: [10.1016/j.physletb.2020.135454](https://doi.org/10.1016/j.physletb.2020.135454). URL: <https://www.sciencedirect.com/science/article/pii/S0370269320302586/pdf> (cit. on pp. 40, 41).
- [3] G. Altarelli and G. Parisi. “Asymptotic freedom in parton language”. In: *Nuclear Physics B* 126.2 (Aug. 1977), pp. 298–318. ISSN: 05503213. DOI: [10.1016/0550-3213\(77\)90384-4](https://doi.org/10.1016/0550-3213(77)90384-4). URL: <https://www.sciencedirect.com/science/article/pii/0550321377903844> (cit. on p. 28).
- [4] E. Amaldi, S. Fubini, and G. Furlan. *Pion-electroproduction*. Vol. 83. Berlin, Heidelberg: Springer Berlin Heidelberg, 1979, pp. 1–10. ISBN: 978-3-540-08998-8. DOI: [10.1007/BFb0048208](https://doi.org/10.1007/BFb0048208) (cit. on p. 37).
- [5] Christian W. Bauer et al. “Hard scattering factorization from effective field theory”. In: *Physical Review D* 66.1 (July 2002), p. 014017. ISSN: 0556-2821.

DOI: [10.1103/PhysRevD.66.014017](https://doi.org/10.1103/PhysRevD.66.014017). URL: <https://arxiv.org/pdf/hep-ph/0202088.pdf> (cit. on p. 34).

- [6] I. Bedlinskiy et al. “Exclusive Neutral Pion electroproduction at W>2 GeV with CLAS”. In: *Physical Review C - Nuclear Physics* 90.2 (2014). ISSN: 1089490X. DOI: [10.1103/PhysRevC.90.025205](https://doi.org/10.1103/PhysRevC.90.025205). URL: <https://journals.aps.org/prc/abstract/10.1103/PhysRevC.90.025205> <https://arxiv.org/pdf/1405.0988.pdf> (cit. on pp. 38, 40, 41).
- [7] J. D. Bjorken and E. A. Paschos. “Inelastic Electron-Proton and gamma-Proton Scattering and the Structure of the Nucleon”. In: *Physical Review* 185.5 (Sept. 1969), pp. 1975–1982. ISSN: 0031-899X. DOI: [10.1103/PhysRev.185.1975](https://doi.org/10.1103/PhysRev.185.1975). URL: <https://journals.aps.org/pr/abstract/10.1103/PhysRev.185.1975> (cit. on p. 26).
- [8] The Editors of Encyclopaedia Britannica. *law of multiple proportions*. 2010. URL: <https://www.britannica.com/science/law-of-multiple-proportions> (cit. on p. 15).
- [9] The Editors of Encyclopaedia Britannica. *Thomson atomic model*. 2023. URL: <https://www.britannica.com/science/Thomson-atomic-model> (cit. on p. 16).
- [10] Stanley J. Brodsky, Markus Diehl, and Dae Sung Hwang. “Light-cone wavefunction representation of deeply virtual Compton scattering”. In: *Nuclear Physics B* 596.1-2 (Feb. 2001), pp. 99–124. ISSN: 05503213. DOI: [10.1016/S0550-3213\(00\)00695-7](https://doi.org/10.1016/S0550-3213(00)00695-7). URL: <https://www.sciencedirect.com/science/article/abs/pii/S0550321300006957> <https://arxiv.org/pdf/hep-ph/0009254.pdf> (cit. on p. 104).

- [11] Louis de Broglie. “A tentative theory of light quanta”. In: *The London, Edinburgh, and Dublin Philosophical Magazine and Journal of Science* 47.278 (Feb. 1924), pp. 446–458. ISSN: 1941-5982. DOI: [10.1080/14786442408634378](https://doi.org/10.1080/14786442408634378). URL: <https://www.tandfonline.com/doi/full/10.1080/14786442408634378?scroll=top&needAccess=true&role=tab> (cit. on p. 18).
- [12] M. Burkardt and B. Pasquini. “Modelling the nucleon structure”. In: *The European Physical Journal A* 52.6 (June 2016), p. 161. ISSN: 1434-6001. DOI: [10.1140/epja/i2016-16161-7](https://doi.org/10.1140/epja/i2016-16161-7). URL: <https://arxiv.org/pdf/1510.02567.pdf> (cit. on p. 34).
- [13] Matthias Burkardt. “GPDs with ζ Not Equal to 0”. In: *ArXiv* (Nov. 2007). URL: <https://arxiv.org/pdf/0711.1881.pdf> (cit. on p. 34).
- [14] V. D. Burkert, L. Elouadrhiri, and F. X. Girod. “The pressure distribution inside the proton”. In: *Nature* 557.7705 (May 2018), pp. 396–399. ISSN: 0028-0836. DOI: [10.1038/s41586-018-0060-z](https://doi.org/10.1038/s41586-018-0060-z). URL: <https://www.nature.com/articles/s41586-018-0060-z> (cit. on p. 36).
- [15] V.D. Burkert et al. “The CLAS12 Spectrometer at Jefferson Laboratory”. In: *Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment* 959 (Apr. 2020), p. 163419. ISSN: 01689002. DOI: [10.1016/j.nima.2020.163419](https://doi.org/10.1016/j.nima.2020.163419). URL: [https://www.jlab.org/div_dept/physics_division/dsg/publications2/CLAS12%20NIM%20A959%20\(2020\)%20163419.pdf%20http://eprints.gla.ac.uk/210295/1/210295.pdf](https://www.jlab.org/div_dept/physics_division/dsg/publications2/CLAS12%20NIM%20A959%20(2020)%20163419.pdf%20http://eprints.gla.ac.uk/210295/1/210295.pdf) (cit. on p. 45).
- [16] C.C.W. Taylor. *The atomists, Leucippus and Democritus: fragments, a text and translation with a commentary*. University of Toronto Press, 1999, pp. 157–158. ISBN: 0-8020-4390-9 (cit. on p. 15).

- [17] J. Chadwick. “Possible Existence of a Neutron”. In: *Nature* 129.3252 (Feb. 1932), pp. 312–312. ISSN: 0028-0836. DOI: [10.1038/129312a0](https://doi.org/10.1038/129312a0) (cit. on p. 17).
- [18] Fei Chen, Paul W. Tillberg, and Edward S. Boyden. “Expansion microscopy”. In: *Science* 347.6221 (Jan. 2015), pp. 543–548. ISSN: 0036-8075. DOI: [10.1126/science.1260088](https://doi.org/10.1126/science.1260088). URL: <https://www.science.org/doi/10.1126/science.1260088> (cit. on p. 18).
- [19] John C. Collins and Andreas Freund. “Proof of factorization for deeply virtual Compton scattering in QCD”. In: *Physical Review D* 59.7 (Feb. 1999), p. 074009. ISSN: 0556-2821. DOI: [10.1103/PhysRevD.59.074009](https://doi.org/10.1103/PhysRevD.59.074009). URL: <https://arxiv.org/pdf/hep-ph/9801262.pdf> (cit. on p. 34).
- [20] M. Derrick et al. “Measurement of the proton structure function F2 from the 1993 HERA data”. In: *Zeitschrift für Physik C Particles and Fields* 65.3 (Sept. 1995), pp. 379–398. ISSN: 0170-9739. DOI: [10.1007/BF01556128](https://doi.org/10.1007/BF01556128) (cit. on p. 104).
- [21] M. Diehl and S. Sapeta. “On the analysis of lepton scattering on longitudinally or transversely polarized protons”. In: *The European Physical Journal C* 41.4 (June 2005), pp. 515–533. ISSN: 1434-6044. DOI: [10.1140/epjc/s2005-02242-9](https://doi.org/10.1140/epjc/s2005-02242-9). URL: <https://link.springer.com/article/10.1140/epjc/s2005-02242-9%20https://arxiv.org/pdf/hep-ph/0503023.pdf> (cit. on p. 104).
- [22] M. Dlamini et al. “Deep Exclusive Electroproduction of $\pi^+ \pi^-$ at High Q^2 in the Quark Valence Regime”. In: *Physical Review Letters* 127.15 (Oct. 2021), p. 152301. ISSN: 0031-9007. DOI: [10.1103/PhysRevLett.127.152301](https://doi.org/10.1103/PhysRevLett.127.152301). URL: <https://arxiv.org/pdf/2107.05001.pdf>

//journals.aps.org/prl/pdf/10.1103/PhysRevLett.127.152301 (cit. on pp. 40, 41).

- [23] A. Donnachie and G. Shaw. “Generalized Vector Dominance”. In: *Electromagnetic Interactions of Hadrons*. Boston, MA: Springer US, 1978, pp. 169–194. DOI: [10.1007/978-1-4757-0713-7_\]3](https://doi.org/10.1007/978-1-4757-0713-7_3) (cit. on p. 37).
- [24] T. William Donnelly et al. *Foundations of Nuclear and Particle Physics*. Cambridge University Press, Feb. 2017. ISBN: 9780521765114. DOI: [10.1017/9781139028264](https://doi.org/10.1017/9781139028264) (cit. on p. 25).
- [25] T.W. Donnelly, Sabine Jeschonnek, and J.W. Van Orden. “General tensor structure for electron scattering in terms of invariant responses”. In: *Annals of Physics* 448 (Jan. 2023), p. 169174. ISSN: 00034916. DOI: [10.1016/j.aop.2022.169174](https://doi.org/10.1016/j.aop.2022.169174) (cit. on p. 37).
- [26] D Dreschsel and L Tiator. “Threshold pion photoproduction on nucleons”. In: *Journal of Physics G: Nuclear and Particle Physics* 18.3 (Mar. 1992), pp. 449–497. ISSN: 0954-3899. DOI: [10.1088/0954-3899/18/3/004](https://doi.org/10.1088/0954-3899/18/3/004). URL: <https://iopscience.iop.org/article/10.1088/0954-3899/18/3/004/pdf> (cit. on pp. 37, 82).
- [27] Richard P. Feynman. “Very High-Energy Collisions of Hadrons”. In: *Physical Review Letters* 23.24 (Dec. 1969), pp. 1415–1417. ISSN: 0031-9007. DOI: [10.1103/PhysRevLett.23.1415](https://doi.org/10.1103/PhysRevLett.23.1415). URL: <https://journals.aps.org/prl/abstract/10.1103/PhysRevLett.23.1415%20https://authors.library.caltech.edu/3871/1/FEYprl169.pdf> (cit. on p. 27).
- [28] Linda E. Franken et al. “A Technical Introduction to Transmission Electron Microscopy for Soft-Matter: Imaging, Possibilities, Choices, and Technical De-

- velopments”. In: *Small* 16.14 (Apr. 2020), p. 1906198. ISSN: 1613-6810. DOI: [10.1002/smll.201906198](https://doi.org/10.1002/smll.201906198). URL: <https://onlinelibrary.wiley.com/doi/full/10.1002/smll.201906198> (cit. on p. 18).
- [29] E. Fuchey et al. “Exclusive neutral pion electroproduction in the deeply virtual regime”. In: *Physical Review C* 83.2 (Feb. 2011), p. 025201. ISSN: 0556-2813. DOI: [10.1103/PhysRevC.83.025201](https://doi.org/10.1103/PhysRevC.83.025201). URL: <https://journals.aps.org/prc/abstract/10.1103/PhysRevC.83.025201%20https://arxiv.org/pdf/1003.2938.pdf> (cit. on pp. 40, 41).
- [30] Hans Geiger and Ernest Marsden. “On a Diffuse Reflection of the alpha-Particles”. In: *Proceedings of the Royal Society of London Series A* 82.557 (1909), pp. 495–500 (cit. on p. 17).
- [31] M. Göckeler et al. “Transverse Spin Structure of the Nucleon from Lattice-QCD Simulations”. In: *Physical Review Letters* 98.22 (May 2007), p. 222001. ISSN: 0031-9007. DOI: [10.1103/PhysRevLett.98.222001](https://doi.org/10.1103/PhysRevLett.98.222001). URL: <https://arxiv.org/pdf/hep-lat/0612032.pdf%20https://journals.aps.org/prl/abstract/10.1103/PhysRevLett.98.222001> (cit. on p. 104).
- [32] Gary R Goldstein, J Osvaldo Gonzalez Hernandez, and Simonetta Liuti. “Easy as π 0: on the interpretation of recent electroproduction results”. In: *Journal of Physics G: Nuclear and Particle Physics* 39.11 (Nov. 2012), p. 115001. ISSN: 0954-3899. DOI: [10.1088/0954-3899/39/11/115001](https://doi.org/10.1088/0954-3899/39/11/115001). URL: <https://inspirehep.net/literature/1086573%20https://arxiv.org/pdf/1201.6088.pdf> (cit. on p. 103).
- [33] Gary R. Goldstein, J. Osvaldo Gonzalez Hernandez, and Simonetta Liuti. “Flexible parametrization of generalized parton distributions from deeply virtual Compton scattering observables”. In: *Physical Review D* 84.3 (Aug. 2011),

- p. 034007. ISSN: 1550-7998. DOI: [10.1103/PhysRevD.84.034007](https://doi.org/10.1103/PhysRevD.84.034007). URL: <https://journals.aps.org/prd/abstract/10.1103/PhysRevD.84.034007> 20<https://arxiv.org/pdf/1012.3776.pdf> (cit. on p. 103).
- [34] M. Hillery et al. “Distribution functions in physics: Fundamentals”. In: *Physics Reports* 106.3 (Apr. 1984), pp. 121–167. ISSN: 03701573. DOI: [10.1016/0370-1573\(84\)90160-1](https://doi.org/10.1016/0370-1573(84)90160-1). URL: <http://ursula.chem.yale.edu/~batista/classes/v572/Scully.pdf> (cit. on p. 31).
- [35] Tomáš Husek, Evgueni Goudzovski, and Karol Kampf. “Precise Determination of the Branching Ratio of the Neutral-Pion Dalitz Decay”. In: *Physical Review Letters* 122.2 (Jan. 2019), p. 022003. ISSN: 0031-9007. DOI: [10.1103/PhysRevLett.122.022003](https://doi.org/10.1103/PhysRevLett.122.022003). URL: <https://arxiv.org/pdf/1809.01153.pdf> (cit. on p. 89).
- [36] Jaume Navarro. *A History of the Electron: J.J. and G.P. Thomson*. Cambridge University Press, 1995. ISBN: 978-1-107-00522-8. (Cit. on p. 16).
- [37] Xiangdong Ji. “Gauge-Invariant Decomposition of Nucleon Spin”. In: *Physical Review Letters* 78.4 (Jan. 1997), pp. 610–613. ISSN: 0031-9007. DOI: [10.1103/PhysRevLett.78.610](https://doi.org/10.1103/PhysRevLett.78.610). URL: <https://journals.aps.org/prl/abstract/10.1103/PhysRevLett.78.610> 20<https://arxiv.org/pdf/hep-ph/9603249.pdf> (cit. on p. 34).
- [38] Xiangdong Ji. “GENERALIZED PARTON DISTRIBUTIONS”. In: *Annual Review of Nuclear and Particle Science* 54.1 (Dec. 2004), pp. 413–450. ISSN: 0163-8998. DOI: [10.1146/annurev.nucl.54.070103.181302](https://doi.org/10.1146/annurev.nucl.54.070103.181302). URL: <https://www.annualreviews.org/doi/10.1146/annurev.nucl.54.070103.181302> (cit. on pp. 32, 33).

- [39] R. Johnston et al. “Realization of a large-acceptance Faraday Cup for 3 MeV electrons”. In: *Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment* 922 (Apr. 2019), pp. 157–160. ISSN: 01689002. DOI: [10.1016/j.nima.2018.12.080](https://doi.org/10.1016/j.nima.2018.12.080). URL: <https://arxiv.org/abs/1811.12196><https://www.sciencedirect.com/science/article/abs/pii/S0168900218318965> (cit. on p. 46).
- [40] Max Klein. *Resolving the Inner Structure of Matter The H1 Experiment at HERA*. Tech. rep. Workshop at Podgorica - 25 years of the Science Faculty, 2005 (cit. on p. 30).
- [41] Valery Kubarovskiy. “Deeply Virtual Exclusive Reactions with CLAS”. In: *Nuclear Physics B - Proceedings Supplements* 219-220 (Oct. 2011), pp. 118–125. ISSN: 09205632. DOI: [10.1016/j.nuclphysbps.2011.10.080](https://doi.org/10.1016/j.nuclphysbps.2011.10.080). URL: <https://www.sciencedirect.com/science/article/abs/pii/S0920563211008218/pdf?md5=2efd6bc3a69acdce62fc3d0bfa1be5c5&pid=1-s2.0-S0920563211008218-main.pdf> (cit. on p. 35).
- [42] Sangbaek Lee. “Measurement of the Deeply Virtual Compton Scattering Cross Section from the Proton at 10.6 GeV using the CLAS12 Detector”. PhD thesis. Cambridge: Massachusetts Institute of Technology, Sept. 2022 (cit. on p. 36).
- [43] Xuezhi Ma et al. “6 nm super-resolution optical transmission and scattering spectroscopic imaging of carbon nanotubes using a nanometer-scale white light source”. In: *Nature Communications* 12.1 (Nov. 2021), p. 6868. ISSN: 2041-1723. DOI: [10.1038/s41467-021-27216-5](https://doi.org/10.1038/s41467-021-27216-5) (cit. on p. 18).
- [44] R. W. McAllister and R. Hofstadter. “Elastic Scattering of 188-MeV Electrons from the Proton and the Alpha Particle”. In: *Physical Review* 102 (Mar. 1956), pp. 851–856 (cit. on p. 103).

- [45] C. Patrignani and et al. (Particle Data Group). *Review of Particle Physics*. Tech. rep. lawrence berkeley national laboratory, 2018, pp. 441–455. DOI: [10.1088/1674-1137/40/10/100001](https://doi.org/10.1088/1674-1137/40/10/100001). URL: <https://pdg.lbl.gov/> (cit. on p. 104).
- [46] Jian-Wei Qiu and Zhite Yu. “Single diffractive hard exclusive processes for the study of generalized parton distributions”. In: *Physical Review D* 107.1 (Jan. 2023), p. 014007. ISSN: 2470-0010. DOI: [10.1103/PhysRevD.107.014007](https://doi.org/10.1103/PhysRevD.107.014007). URL: <https://journals.aps.org/prd/pdf/10.1103/PhysRevD.107.014007> [20https://arxiv.org/pdf/2210.07995.pdf](https://arxiv.org/pdf/2210.07995.pdf) (cit. on p. 104).
- [47] Ernest Rutherford. “Collision of alpha Particles with Light Atoms. IV. An Anomalous Effect in Nitrogen”. In: *Philosophical Magazine Series 6* 37 (1919), pp. 581–587. URL: <https://www.tandfonline.com/doi/abs/10.1080/14786431003659230> (cit. on p. 17).
- [48] Ernest Rutherford. “The Scattering of Alpha and Beta Particles by Matter and the Structure of the Atom”. In: *Philosophical Magazine Series 6* 21 (1911), pp. 669–688 (cit. on p. 17).
- [49] P. E. Shanahan and W. Detmold. “Pressure Distribution and Shear Forces inside the Proton”. In: *Physical Review Letters* 122.7 (Feb. 2019), p. 072003. ISSN: 0031-9007. DOI: [10.1103/PhysRevLett.122.072003](https://doi.org/10.1103/PhysRevLett.122.072003). URL: <https://journals.aps.org/prl/abstract/10.1103/PhysRevLett.122.072003> [20https://arxiv.org/pdf/1810.07589.pdf](https://arxiv.org/pdf/1810.07589.pdf) (cit. on p. 36).
- [50] Thomas McEvilley. *The Shape of Ancient Thought: Comparative Studies in Greek and Indian Philosophies*. New York: Allworth Press, 2002. ISBN: 978-0-8020-4390-0 (cit. on p. 15).

- [51] J.J. Thomson. “On Bodies Smaller than Atoms”. In: *The Poular Science Monthly* (1901), pp. 323–325. URL: <https://books.google.com/books?id=3CMDAAAAMBAJ&pg=PA323#v=onepage&q&f=false> (cit. on p. 15).
- [52] Mark Thomson. *Modern Particle Physics*. Cambridge University Press, Sept. 2013, pp. 160–204. ISBN: 9781107034266. DOI: [10.1017/CBO9781139525367](https://doi.org/10.1017/CBO9781139525367) (cit. on pp. 23, 26).
- [53] E. Wigner. “On the Quantum Correction For Thermodynamic Equilibrium”. In: *Physical Review* 40.5 (June 1932), pp. 749–759. ISSN: 0031-899X. DOI: [10.1103/PhysRev.40.749](https://doi.org/10.1103/PhysRev.40.749). URL: https://lab409.chem.ccu.edu.tw/~yach932/CH3_Reference/51.PhysRev.40.749.pdf (cit. on p. 31).
- [54] David B. Williams and C. Barry Carter. *Transmission Electron Microscopy*. Boston, MA: Springer US, 2009. ISBN: 978-0-387-76500-6. DOI: [10.1007/978-0-387-76501-3](https://doi.org/10.1007/978-0-387-76501-3). URL: <https://link.springer.com/book/10.1007/978-0-387-76501-3> (cit. on pp. 18, 19).
- [55] Yimei Zhu and Hermann Dürr. “The future of electron microscopy”. In: *Physics Today* 68.4 (Apr. 2015), pp. 32–38. ISSN: 0031-9228. DOI: [10.1063/PT.3.2747](https://doi.org/10.1063/PT.3.2747). URL: <https://physicstoday.scitation.org/doi/10.1063/PT.3.2747> (cit. on p. 104).
- [56] P A Zyla et al. “Review of Particle Physics”. In: *Progress of Theoretical and Experimental Physics* 2020.8 (Aug. 2020). ISSN: 2050-3911. DOI: [10.1093/ptep/ptaa104](https://doi.org/10.1093/ptep/ptaa104). URL: <https://pdg.lbl.gov/2020/reviews/rpp2020-rev-structure-functions.pdf> (cit. on pp. 28, 29).

Appendix A

Full Cross Section Data

To be completed

Appendix B

BSA Cross Check

As an additional cross check, Bobby calculated a $DV\pi^0P$ beam spin asymmetry and compared to Andrey Kim's results. This check will not comment on any acceptance, luminosity, or virtual photon flux factor calculations, but does validate exclusive event selection criteria. By examining figure B-1 we can see that agreement is reasonable, especially considering Bobby's calculation does not have sideband subtraction included.

Fig B-1 shows an overlay comparison of Andrey Kim's results (black datapoints, red fit line) and Bobby's results (red datapoints, orange fit line)




Figure B-1: BSA Cross Check Results

Appendix C

Derivation of phi math convention

Thus: $\phi = \arccos((\mathbf{v3l} \cdot \mathbf{v3h}) / (\text{mag } \mathbf{v3l} \text{ mag } \mathbf{v3h}))$

$$\phi = \cos^{-1} \left(\frac{(\mathbf{p}_e \times \mathbf{p}_{e'}) \cdot (\mathbf{p}_{p'} \times \mathbf{p}_{\gamma^*})}{\|\mathbf{p}_e \times \mathbf{p}_{e'}\| \|\mathbf{p}_{p'} \times \mathbf{p}_{\gamma^*}\|} \right)$$

if $\text{dot}(\mathbf{p}_e \times \mathbf{p}_{e'}, \mathbf{p}_{p'})$ is greater than 0, then do $360 - \phi = \phi$. If we expand the above out, we get: $-\mathbf{p}_{p'} \cdot \mathbf{e}_z \mathbf{e}_y + \mathbf{p}_y \cdot \mathbf{e}_z \mathbf{e}_x$ is greater than zero which we can reduce to $-\mathbf{p}_{p'} \cdot \mathbf{e}_y + \mathbf{p}_y \cdot \mathbf{e}_x$ is greater than zero

By inspecting table below, we can see what this really amounts to, is the trento convention saying that we take the angle by measuring counterclockwise from the proton vector to the electron vector.

$360 - \theta$ when $P_y'e'_x > P_xe'_y$

P_y	P_x	e_y	e_x	L	R	θ	$ e'_x $	$ e'_y $
-	-	-	-	+	+		-	-
-	-	-	+	-	+	$< 180^\circ$	-	+
-	-	t	-	+	-	$> 180^\circ$	-	+
-	-	+	7	-	-		$> 180^\circ$	
-	+	-	-	+	-			
-	+	-	+	-	+			
-	+	+	+	-	+			
-	+	t	7	-	+			

$P_x > e$



$-e'_x > -e'_y$ $e'_x < e'_y$ $|e'_x| > |e'_y|$ shear in

